

Common Distributions

Binomial distribution

$$f(x|n, p) = \binom{n}{x} p^x (1-p)^{n-x}$$

$$E(x) = np$$

$$\text{var}(x) = np(1-p)$$

Binomial divided by number of trials:

$$E(x/n) = p$$


$$\text{var}(x/n) = p(1-p)/n$$

Copyright Dick Startz

1

Copyright Dick Startz

2



Copyright Dick Startz

3

Copyright Dick Startz

4

Write a Matlab program that plots two figures (one figure for small n and one for a large n), each comparing two cdfs. For both figures, one cdf is the number of heads out of n coin flips with a probability of heads equals to $p = 1/2$. The other cdf is for a normal distribution with mean np and standard deviation $\sqrt{np(1-p)}$. The Matlab function `binocdf` will give you the former and `normcdf` will give you the latter.

Average of standard normals

$$x \sim N(0,1)$$

$$\bar{x} \sim N\left(0, \frac{1}{n}\right)$$

$$p(\sqrt{n}\bar{x} < -1.96 \cup \sqrt{n}\bar{x} > 1.96) = 0.05$$

$$I_{\sqrt{n}\bar{x} < -1.96 \cup \sqrt{n}\bar{x} > 1.96} \sim \text{Bernoulli}$$

Suppose we draw n independent random variables $x_i \sim N(0,1)$ and generate the average \bar{x} . We know that $\Pr\left(\left|\bar{x}/\sqrt{1/n}\right| > 1.96\right) = 0.05$. Suppose we were to generate m such samples and count up the number of times that $\left|\bar{x}/\sqrt{1/n}\right| > 1.96$. Call this count divided by m , α .

- What is the standard deviation of α ?
- Suppose we wanted there to be a 90 percent chance that the estimated α is between 0.04 and 0.06. How large must m be? You may approximate the distribution of α by the normal distribution.
- Write a simulation program that sets $n = 10$ and uses a variety of values of m , some smaller than your value given in part (b) and some larger. Then run this for a number of times saving α . Make a plot that shows the theoretical standard deviation of α (from part (a)) as a function of m as well as the empirical standard deviation of α from the simulations. Effectively, do a Monte Carlo of doing a Monte Carlo.

Copyright Dick Startz

6

Copyright Dick Startz

7

Poisson


$$f(x) = \frac{e^{-\lambda} \lambda^x}{x!}, x = 0, 1, \dots$$

$$E(x) = \lambda$$

$$\text{var}(x) = \lambda$$

$$f(x) = \frac{\lambda}{x} f(x-1),$$

$$x = 1, 2, \dots$$



Copyright Dick Startz

10

Zero-inflated Poisson

$$f(x=0) = \pi + (1-\pi)e^{-\lambda}$$

$$f(x|x > 0) = (1-\pi) \frac{e^{-\lambda} \lambda^x}{x!}$$

$$E(x) = (1-\pi)\lambda$$

$$\text{var}(x) = \lambda(1-\pi)(1+\lambda\pi)$$

Copyright Dick Startz

11

Negative binomial

$$f(x) = \binom{x-1}{r-1} p^r (1-p)^{x-r}, x = r, r+1, \dots$$

$$E(x) = \frac{r}{p}$$

$$\text{var}(x) = r \frac{1-p}{p^2}$$

n.b. The Poisson is a special case of the negative binomial with

$$\lambda = \frac{r}{1-p}, r \rightarrow \infty, p \rightarrow 1$$

Copyright Dick Startz

12


Geometric density

$$f(x) = p(1-p)^{x-1}, x = 1, 2, \dots$$

$$E(x) = \frac{1}{p}$$

$$\text{var}(x) = \frac{1-p}{p^2}$$

where p is the probability of success in one trial



Copyright Dick Startz

13

Geometric is memoryless

- Note that the probability of no success in n periods is $(1-p)^n$

For $s > t$

$$f(x > s | x > t) = \frac{P(x > s \cap x > t)}{P(x > t)} = \frac{P(x > s)}{P(x > t)}$$


$$\begin{aligned} f(x > s | x > t) &= \frac{(1-p)^s}{(1-p)^t} = (1-p)^{s-t} \\ &= f(x > s-t) \end{aligned}$$

Copyright Dick Startz

14

Discrete uniform

$$f(x) = \frac{1}{n}, x = 1, 2, \dots, n$$



Copyright Dick Startz

15

Continuous Uniform

$$f(x|a, b) = \begin{cases} \frac{1}{b-a}, & \text{if } x \in [a, b] \\ 0, & \text{otherwise} \end{cases}$$

$$E(x) = \frac{b+a}{2}$$

$$\text{var}(a) = \frac{(b-a)^2}{12}$$

Copyright Dick Startz

16

Pareto distribution


$$f(x|\alpha, \beta) = \frac{\beta \alpha^\beta}{x^{\beta+1}}, \alpha < x < \infty, \alpha > 0, \beta > 0$$

$$E(x) = \frac{\beta \alpha}{\beta - 1}, \beta > 1$$

$$\text{var}(x) = \frac{\beta \alpha^2}{(\beta - 1)^2 (\beta - 2)}, \beta > 2$$

Copyright Dick Startz

17



Copyright Dick Startz

18

Gamma

$$f(x|\alpha, \beta) = \frac{1}{\Gamma(\alpha)\beta^\alpha} x^{\alpha-1} e^{-\frac{x}{\beta}}, 0 < x < \infty, \alpha > 0, \beta > 0$$

$$\Gamma(\alpha) = \int_0^\infty t^{\alpha-1} e^{-t} dt$$

$$\Gamma(\alpha + 1) = \alpha \Gamma(\alpha)$$

$$\Gamma(n) = (n-1)! \text{ for integer } n$$


$$E(x) = \alpha \beta$$

$$\text{var}(x) = \alpha \beta^2$$

α is called the shape parameter. β is called the scale parameter.

Copyright Dick Startz

19



Copyright Dick Startz

20

χ_p^2 (chi-squared)

- χ_p^2 is gamma with $\beta = 2$ (and $p = 2\alpha$).

$$f(x|p) = \frac{1}{\Gamma\left(\frac{p}{2}\right) 2^{\frac{p}{2}}} x^{\frac{p}{2}-1} e^{-\frac{x}{2}}, 0 < x$$

$$\begin{aligned} x &\sim \chi_p^2 \\ E(x) &= p \\ \text{var}(x) &= 2p \end{aligned}$$

Copyright Dick Startz

21

Write a Matlab program that draws a simulated $\chi^2(4)$ density and an analytic density on the same figure. To simulate the χ^2 generate 4 standard normals (`randn(4,1)`), square them and add them up. Do this a lot of times and then draw the histogram using the Matlab function `histogram`. Then draw the analytic expression using the `chi2pdf` function with 4 degrees of freedom.

Copyright Dick Startz

22

Exponential

$$\begin{aligned} f(x) &= \lambda e^{-\lambda x}, 0 < x < \infty \\ E(x) &= 1/\lambda \\ \text{var}(x) &= 1/\lambda^2 \end{aligned}$$

$$F(x) = 1 - e^{-\lambda x}$$

Copyright Dick Startz

24

Exponential and discounting

- Suppose we are going to receive x dollars in t years discounted at rate r , then the present value would be

$$PV = xe^{-rt}$$

- Payment date is uncertain with exponential distribution

Copyright Dick Startz

25

Exponential and discounting cont.

$$\begin{aligned} EPV &= \int_0^{\infty} xe^{-rt} f(t) dt \\ EPV &= \int_0^{\infty} xe^{-rt} \lambda e^{-\lambda t} dt \\ &= x \frac{\lambda}{\lambda + r} \int_0^{\infty} (\lambda + r) e^{-(\lambda + r)t} dt \\ &= x \frac{\lambda}{\lambda + r} \times 1 = \frac{x}{1 + r/\lambda} \end{aligned}$$

Copyright Dick Startz

26

Gaussian or Normal

$$f(x|\mu, \sigma^2) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{1}{2\sigma^2}(x-\mu)^2}$$

$$E(x) = \mu$$


$$\text{var}(x) = \sigma^2$$

$$x \sim N(\mu, \sigma^2)$$

The normal is symmetric around μ and has a maximum at μ .

Copyright Dick Startz

27



Copyright Dick Startz

28

Standard normal

$$\phi(x) \equiv f_N(x|0,1)$$

$$\Phi(x) \equiv F_N(x|0,1)$$


- It is useful that

$$f(x|\mu, \sigma^2) = \frac{1}{\sigma} \phi\left(\frac{x-\mu}{\sigma}\right)$$

$$F(x|\mu, \sigma^2) = \Phi\left(\frac{x-\mu}{\sigma}\right)$$

Copyright Dick Startz

29



Copyright Dick Startz

30


Log normal

If $\log x \sim N(\mu, \sigma^2)$, then
 $x \sim \ln N(\mu, \sigma^2)$ is distributed
lognormal.

$$f(x) = \frac{1}{x\sqrt{2\pi\sigma^2}} e^{-\frac{1}{2\sigma^2}(\log x - \mu)^2}$$

$$E(x) = e^{\mu + \frac{\sigma^2}{2}}$$

$$\text{var}(x) = (e^{\sigma^2} - 1)e^{(2\mu + \sigma^2)}$$



Copyright Dick Startz

31

Example of proportional errors

- Cobb-Douglas with errors


$$Y = AK^\beta L^{1-\beta} \varepsilon$$

- If $\varepsilon \sim \ln N(\cdot)$, then

$$\log Y = \log A + \beta \log K + (1 - \beta) \log L + \log \varepsilon$$

Copyright Dick Startz

32



Copyright Dick Startz

33

Location and scale families

Theorem 3.5.1 Let $f(x)$ be any pdf and let μ and $\sigma > 0$ be any given constants, then the function

$$g(x|\mu, \sigma) = \frac{1}{\sigma} f\left(\frac{x - \mu}{\sigma}\right)$$

is a pdf and

$$G(x|\mu, \sigma) = F\left(\frac{x - \mu}{\sigma}\right)$$

is a cdf.

Copyright Dick Startz

34

Exponential families

- A pdf is an *exponential family* if it can be expressed as

$$f(x|\theta) = h(x)c(\theta) \exp\left(\sum_{i=1}^k w_i(\theta) t_i(x)\right)$$

Includes: normal, gamma, beta, binomial, and Poisson

Copyright Dick Startz

35

Logistic distribution

$$f(x) = \frac{e^{-\frac{x-\mu}{\sigma}}}{\sigma \times \left(1 + e^{-\frac{x-\mu}{\sigma}}\right)^2}$$

$$E(x) = \mu$$

$$\text{var}(x) = \frac{\sigma^2 \pi^2}{3}$$

$$F(x) = \frac{1}{1 + e^{-\frac{x-\mu}{\sigma}}}$$

Copyright Dick Startz

36

Standard logistic distribution

$$F(x) = \frac{1}{1 + e^{-\frac{x-\mu}{\sigma}}}$$

Standard logistic, $\mu = 0, \sigma = 1$ which gives

$$F(x) = \frac{1}{1 + e^{-x}}$$

An application of the standard logistic is if we think something happens if $x < x_0$ and otherwise doesn't happen. Then the *odds* of it happening are

$$\frac{F(x_0)}{1 - F(x_0)} = e^{x_0}, \log \frac{F(x_0)}{1 - F(x_0)} = x_0$$

Copyright Dick Startz

37

Logistic regression on UW law school admissions

Dependent Variable: ADMIT
Method: ML - Binary Logit (Newton-Raphson / Marquardt steps)
Date: 06/16/15 Time: 09:32
Sample: 1 1643 IF LSAT>99 AND LSAT<200 AND GPA<=4
Included observations: 1567
Convergence achieved after 4 iterations
Coefficient covariance computed using observed Hessian

Variable	Coefficient...	Std. Error	z-Statistic	Prob.
C	-10.02057	0.771661	-12.98571	0.0000
GPA	2.637745	0.218239	12.08650	0.0000
McFadden R-square...	0.100332	Mean dependent var	0.296107	
S.D. dependent var	0.456685	S.E. of regression	0.428441	
Akaike info criterion	1.095703	Sum squared resid	287.2737	
Schwarz criterion	1.102540	Log likelihood	-856.4831	
Hannan-Quinn criter.	1.096244	Deviance	1712.966	
Restr. deviance	1903.996	Restr. log likelihood	-951.9992	
LR statistic	191.0322	Avg. log likelihood	-0.546575	
Prob(LR statistic)	0.000000			
Obs with Dep=0	1103	Total obs	1567	
Obs with Dep=1	464			

Copyright Dick Startz

38

Extreme value (Gumbel) Distribution

$$f(x) = \frac{1}{\beta} e^{-\left(\left(\frac{x-\mu}{\beta}\right) + e^{-\left(\frac{x-\mu}{\beta}\right)}\right)}$$

$$F(x) = e^{-e^{-\left(\frac{x-\mu}{\beta}\right)}}$$

$$\text{mode} = \mu$$

$$\text{median} = \mu - \beta \log(\log(2))$$


$$\text{mean} = \mu + \beta\gamma$$

where $\gamma \approx 0.5772$ is the Euler-Mascheroni constant.

$$\text{var} = \frac{\pi^2}{6} \beta^2$$

Copyright Dick Startz

39



Copyright Dick Startz

40

The pdf of the extreme value distribution can be

written $f(x) = \frac{1}{\beta} e^{-\left(\frac{x-\mu}{\beta} + e^{-\left(\frac{x-\mu}{\beta}\right)}\right)}$, where
 $mean(x) = \mu + \beta\gamma$, $\gamma \approx 0.5772$ and $median(x) = \mu - \log(\log(2))$. Generate a bunch of samples of standard normals each with 120 observations. Find the maximum value for each sample. Find the mean and median of the distribution of the maxima. Now solve for the values of μ and β implied by the mean and median. Plot the empirical density (use the histogram function) of the maxima along with the theoretical pdf implied by the values of μ and β .