

Bivariate Random Variables

Median

$$\text{median}(F_X) = x \text{ s.t. } F_X(x) = \frac{1}{2}$$

Examples:

$$x \sim N(\mu, \sigma^2) \Rightarrow \text{median}(x) = \mu$$

$$x \sim \text{exponential}(\lambda) \Rightarrow \text{median}(x) = \frac{\log(2)}{\lambda}$$

Copyright Dick Startz

1

Copyright Dick Startz

2


Assignment

- Generate 1,000 samples of 100 iid exponential(λ) variables with $\lambda = 2$, $f(x) = \lambda e^{-\lambda x}$, $0 < x < \infty$.
- Plot the distribution of the 25th, 50th, and 75th percentiles of your samples.

Copyright Dick Startz


3

Median wage and salary income vs years of education
for those reporting positive income
CPS March 2014




Copyright Dick Startz


5



6




7



9

Boxplot



Sometimes width represents proportion of sample, or square root of proportion of sample. Sometimes box is shaded or notched to give confidence intervals for the median

Copyright Dick Startz

10


Bivariate pdf

- An n -dimensional random vector is a function from a sample space S into \mathbb{R}^n .
- A joint pdf (or pmf) maps a random vector into \mathbb{R}^1 . In particular, a bivariate pdf maps a sample in 2-space (maybe \mathbb{R}^2) into \mathbb{R} . We have a joint pdf of the form

$$f_{X,Y}(x, y)$$

Copyright Dick Startz

11



Copyright Dick Startz

12

Moments

$$E(x) = \iint x f(x, y) dx dy$$

Etc.

Copyright Dick Startz

13

Marginal distributions

Discrete:

$$f_X(x) = \sum_Y f_{X,Y}(x, y)$$

Continuous

$$f_X(x) = \int_{-\infty}^{\infty} f_{X,Y}(x, y) dy$$

Copyright Dick Startz

14

Joint and marginals for two dice

$f(x, y)$	1	2	3	4	5	6	$f(x)$
1	1/36	1/36	1/36	1/36	1/36	1/36	1/6
2	1/36	1/36	1/36	1/36	1/36	1/36	1/6
3	1/36	1/36	1/36	1/36	1/36	1/36	1/6
4	1/36	1/36	1/36	1/36	1/36	1/36	1/6
5	1/36	1/36	1/36	1/36	1/36	1/36	1/6
6	1/36	1/36	1/36	1/36	1/36	1/36	1/6
$f(y)$	1/6	1/6	1/6	1/6	1/6	1/6	

Copyright Dick Startz

15

Assignment

- Suppose you have a pair of honest dice. What is the probability of a natural (7 or 11)? What is the probability of craps (2, 3, or 12)?

Copyright Dick Startz

16

Joint CDF

$$F(x, y) = P(X \leq x, Y \leq y)$$

$$F(x, y) = \int_{-\infty}^x \int_{-\infty}^y f(x, y) dy dx$$

- Consider the joint uniform

$$f(x, y) = \begin{cases} 1, & 0 \leq x \leq 1, 0 \leq y \leq 1 \\ 0, & \text{otherwise} \end{cases}$$

$$\begin{aligned} F(0.1, 0.5) &= \int_0^{0.1} \int_0^{0.5} 1 dy dx = \int_0^{0.1} [y]_0^{0.5} dx \\ &= \int_0^{0.1} .5 dx = .05 \end{aligned}$$

Copyright Dick Startz

18

Independent Random Variables

Let (X, Y) be a bivariate random vector with joint pdf $f(x, y)$ and marginal pdfs $f_X(x)$ and $f_Y(y)$. Then X and Y are *independent random variables* if for all (x, y)

$$f(x, y) = f_X(x) \cdot f_Y(y)$$

Copyright Dick Startz

19

Independent Random Variables

$$\begin{aligned}
 f_X(x) &= \int_{-\infty}^{\infty} f_{X,Y}(x,y) dy \\
 f_X(x) &= \int_{-\infty}^{\infty} [f_X(x) \cdot f_Y(y)] dy \\
 &= f_X(x) \cdot \int_{-\infty}^{\infty} f_Y(y) dy = f_X(x) \cdot 1
 \end{aligned}$$

Copyright Dick Startz

20


Bivariate independent normal pdf

- $x_1 \sim N(0,1)$
- $x_2 \sim N(0,1)$

$$\begin{aligned}
 f(x_1, x_2) &= \left[\frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}x_1^2} \right] \times \left[\frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}x_2^2} \right] \\
 &= (2\pi)^{-\frac{n=2}{2}} e^{-\frac{1}{2}(x_1^2 + x_2^2)}
 \end{aligned}$$

Copyright Dick Startz

21



Copyright Dick Startz

22


Conditional density

Let (X, Y) be a continuous bivariate random vector with joint pdf $f(x, y)$ and marginal pdfs $f_X(x)$ and $f_Y(y)$. For any x such that $f_X(x) > 0$, the *conditional pdf of Y given $X = x$* is

$$f(y|x) = \frac{f(x, y)}{f_X(x)}$$

Copyright Dick Startz

23



Copyright Dick Startz

24

A conditional density is a density

$$\int_{-\infty}^{\infty} f(y|x) dy = \int_{-\infty}^{\infty} \frac{f(x, y)}{f_X(x)} dy$$

$$= \frac{1}{f_X(x)} \int_{-\infty}^{\infty} f(x, y) dy$$

$$\frac{1}{f_X(x)} \times f_X(x) = 1$$

Copyright Dick Startz

25

Bayes theorem again

$$f(y|x) = \frac{f(x, y)}{f_X(x)}$$

$$f(x|y) = \frac{f(x, y)}{f_Y(y)}$$

$$f(y|x) = \frac{f(x|y) \times f_Y(y)}{f_X(x)}$$

Copyright Dick Startz

26

Law of total probability

$$P(y) = \sum_x P(y|x)P(x)$$

$$f_Y(y) = \int f(y|x)f_X(x)dx$$

Copyright Dick Startz

27

Conditional moments

$$E(y|x) = \int_{-\infty}^{\infty} y \cdot f(y|x) dy$$

$$\text{var}(y|x) = \int_{-\infty}^{\infty} (y - E(y|x))^2 \cdot f(y|x) dy$$

Copyright Dick Startz

28

Independence and conditional distribution

$$f(y|x) = \frac{f(x, y)}{f_X(x)} = \frac{f_X(x) \cdot f_Y(y)}{f_X(x)} = f_Y(y)$$

Copyright Dick Startz

29

Law of Iterated Expectations

Theorem 4.4.3 If X and Y are random variables then

$$E(E(Y|X)) = E(Y)$$

Provided the expectations exist.

Copyright Dick Startz

30

Proof

Note that $E(y|x)$ is a function of x .

$$\begin{aligned} E(E(Y|X)) &= \int E(y|x) f_X(x) dx \\ &= \int \left[\int y \cdot f(y|x) dy \right] f_X(x) dx \\ &= \int \left[\int y \cdot f(y, x) / f_X(x) dy \right] f_X(x) dx \\ &= \int \int y \cdot f(y, x) dy dx \\ E(y) &= \int \int y \cdot f(y, x) dy dx \end{aligned}$$

Substitute in $f(y|x) = f(y, x)/f_X(x)$

Copyright Dick Startz

31

Law of iterated iterations

$$E(E(y|x_1, x_2) | x_1) = E(y|x_1)$$

Hansen calls this “The smaller information set wins.”

Forecast errors are mean zero

$$y^e = E(y|x)$$

$$\varepsilon = y - y^e$$

$$E(\varepsilon|x) = E(y|x) - E(E(y|x)|x)$$

$$E(y|x) - E(y|x) = 0$$

By the Law of Iterated Expectations $E(\varepsilon) = 0$.

Note: Random walk of stock prices. If theory says $p_t = E(p_{t+1}|I_t)$, then

$$p_{t+1} = E(p_{t+1}|I_t) + \varepsilon$$

$$p_{t+1} = p_t + \varepsilon$$

Copyright Dick Startz

32

Copyright Dick Startz

33

Dependent Variable: CLOSE
Method: Least Squares
Date: 08/28/16 Time: 11:42
Sample (adjusted): 1/04/1950 7/10/2015
Included observations: 16485 after adjustments

Variable	Coefficient	Std. Error	t-Statistic	Prob.
C	0.059183	0.080990	0.730741	0.4649
CLOSE(-1)	1.000139	0.000113	8887.713	0.0000
R-squared	0.999791	Mean dependent var	472.9167	
Adjusted R-squared	0.999791	S.D. dependent var	542.7898	
S.E. of regression	7.840210	Akaike info criterion	6.956530	
Sum squared resid	1013192.	Schwarz criterion	6.957465	
Log likelihood	-57337.19	Hannan-Quinn criter.	6.956838	
F-statistic	78991436	Durbin-Watson stat	2.123942	
Prob(F-statistic)	0.000000			

Copyright Dick Startz

34

$$\frac{\log p_{t+1} - \log p_t}{p_t} \approx \log p_{t+1} - \log p_t = \varepsilon_t$$

Dependent Variable: LNP
Method: Least Squares
Date: 09/07/17 Time: 12:02
Sample (adjusted): 1/04/1950 7/10/2015
Included observations: 16485 after adjustments

Variable	Coefficient	Std. Error	t-Statistic	Prob.
C	0.000555	0.000313	1.775517	0.0758
LNP(-1)	0.999951	5.66E-05	17674.34	0.0000
R-squared	0.999947	Mean dependent var	5.363609	
Adjusted R-squared	0.999947	S.D. dependent var	1.336568	
S.E. of regression	0.009709	Akaike info criterion	-6.431437	
Sum squared resid	1.553717	Schwarz criterion	-6.430501	
Log likelihood	53013.12	Hannan-Quinn criter.	-6.431128	
F-statistic	3.12E+08	Durbin-Watson stat	1.944971	
Prob(F-statistic)	0.000000			

Copyright Dick Startz

35

Conditioning theorem

$$E(x|x) = x$$

$$E(g(x)|x) = g(x)$$

$$E(g(x)y|x) = g(x) E(y|x)$$

$$E(g(x)y) = E(g(x) E(y|x))$$

Copyright Dick Startz

36

Forecast error is uncorrelated with
functions of variables used in the
forecast

$$E(h(x)(y - E(y|x))) = 0$$

Proof:

$$= E(E[h(x)(y - E(y|x))|x])$$

By the law of iterated expectations.

$$= E(h(x) E[(y - E(y|x))|x])$$

By conditioning theorem

$$E[(y - E(y|x))|x] = 0$$

Copyright Dick Startz

37

ls log(gdpc96) c log(gdpc96(-1)) log(gdpc96(-2))

fit(d) yfit

show log(gdpc96)-yfit log(gdpc96(-1))

log(gdpc96(-2))

Dependent Variable: LOG(GDPC96)
Method: Least Squares
Date: 10/11/15 Time: 11:38
Sample (adjusted): 1947Q3 2015Q2
Included observations: 272 after adjustments

Variable	Coefficient...	Std. Error	t-Statistic	Prob.
C	0.020863	0.007576	2.753776	0.0063
LOG(GDPC96(-1))	1.351018	0.056859	23.76073	0.0000
LOG(GDPC96(-2))	-0.352819	0.056722	-6.220118	0.0000

R-squared 0.999808 Mean dependent var 8.766165
Adjusted R-squared 0.999806 S.D. dependent var 0.634578
S.E. of regression 0.008835 Akaike info criterion -6.609316
Sum squared resid 0.020995 Schwarz criterion -6.569546
Log likelihood 901.8670 Hannan-Quinn criter. -6.593350
F-statistic 698961.9 Durbin-Watson stat 2.061133
Prob(F-statistic) 0.000000

Copyright Dick Startz

38

Covariance Analysis: Ordinary

Date: 10/11/15 Time: 11:42

Sample: 1947Q3 2015Q2

Included observations: 272

Balanced sample (listwise missing value deletion)

Covariance Correlation	LOG(GDPC96)-YFIT...	LOG(GDPC96(-1))	LOG(GDPC96(-2))...
LOG(GDPC96)-YFIT	7.72E-05 1.000000		
LOG(GDPC96(-1))	3.72E-14 6.67E-12	0.403226 1.000000	
LOG(GDPC96(-2))	-5.21E-14 -9.31E-12	0.404156 0.999890	0.405177 1.000000

Copyright Dick Startz

39

Prove the following theorem

Theorem: If you want to minimize expected square error given the variable x , conditional expected value is right way to forecast. (Note that $E(y|x)$ is a function of x .)

$$y^e = \underset{y^e}{\operatorname{argmin}} E((y - y^e)^2)$$

$$y^e = E(y|x)$$

Copyright Dick Startz

40

Mean independence

Y is mean-independent of X iff

$$E(y|x) = E(y) \forall x \text{ s.t. } f_X(x) \neq 0$$

- Note: Y and X independent imply mean independence, but the implication does not necessarily follow the other way around. There can be information about higher moments without information about the mean.

Copyright Dick Startz

42

Conditional variance identity

Theorem 4.4.7 (Conditional variance identity)

For any two random variables X and Y ,

$$\operatorname{var}(X) = E(\operatorname{var}(X|Y)) + \operatorname{var}(E(X|Y))$$

Provided that the expectations exist.

Copyright Dick Startz

43

Covariance

$$\operatorname{cov}(x, y) = E[(x - E(x))(y - E(y))]$$

$$\operatorname{cov}(x, y)$$

$$= \iint (x - E(x))(y - E(y))f(x, y)dx dy$$

$$\operatorname{cov}(x, y) = E(xy) - E(x)E(y)$$

$$\operatorname{cov}(a + bx, c + dy) = bd \cdot \operatorname{cov}(x, y)$$

Copyright Dick Startz

44

Variance rules

$$\begin{aligned}\text{var}(x + y) &= E([(x - E(x)) + (y - E(y))]^2) \\ &= E((x - E(x))^2) + E((y - E(y))^2) \\ &\quad + 2 E((x - E(x)) \times (y - E(y)))\end{aligned}$$

$$\begin{aligned}\text{var}(ax + by) \\ &= a^2 \text{var}(x) + b^2 \text{var}(y) + 2ab \cdot \text{cov}(x, y) \\ \text{var}(x + y) &= \text{var}(x) + \text{var}(y) + 2 \text{cov}(x, y)\end{aligned}$$

Copyright Dick Startz

45

Correlation

$$\rho_{x,y} = \frac{\text{cov}(x, y)}{\sqrt{\text{var}(x)} \sqrt{\text{var}(y)}}$$

$$\rho_{x,y} = \frac{\sigma_{x,y}}{\sigma_x \sigma_y}$$

Copyright Dick Startz

46

Perfect affine relation

$$y = a + bx$$

$$\begin{aligned}\text{var}(y) &= b^2 \text{var}(x) \\ \text{cov}(y, x) &= b \text{var}(x)\end{aligned}$$

$$\rho = \frac{b \text{var}(x)}{\sqrt{\text{var}(x)} \sqrt{b^2 \text{var}(x)}} = \pm 1$$

Copyright Dick Startz

47

Independence and correlation

$$\text{independence} \Rightarrow \text{uncorrelated}$$

$$\text{mean independence} \Rightarrow \text{uncorrelated}$$

$$\text{uncorrelated} \nRightarrow \text{independence}$$

Copyright Dick Startz

48

Numerical example

$$\begin{bmatrix} r_1 \\ r_2 \end{bmatrix} \sim \left(\begin{bmatrix} \bar{r}_1 \\ \bar{r}_2 \end{bmatrix}, \begin{bmatrix} \sigma_1^2 & \sigma_{12} \\ \sigma_{12} & \sigma_2^2 \end{bmatrix} \right)$$

$$r_p = \omega r_1 + (1 - \omega) r_2$$

$$\begin{bmatrix} r_1 \\ r_2 \end{bmatrix} \sim \left(\begin{bmatrix} .10 \\ .15 \end{bmatrix}, \begin{bmatrix} .1^2 & 0.004 \\ 0.004 & .2^2 \end{bmatrix} \right)$$


$$r_p \sim (\omega \bar{r}_1 + (1 - \omega) \bar{r}_2, \omega^2 \sigma_1^2 + (1 - \omega)^2 \sigma_2^2 + 2\omega(1 - \omega)\sigma_{12})$$

Copyright Dick Startz

49


Copyright Dick Startz

50



Copyright Dick Startz

51



Copyright Dick Startz

52

Weight λ in risk-free


- Expected return

$$\lambda r_f + (1 - \lambda) \bar{r}_p$$
- Standard deviation

$$(1 - \lambda) \sigma_p$$

Copyright Dick Startz

53



Copyright Dick Startz

54

Capital market line

- Goes through the points
 $(r_f, 0)$ and (\bar{r}_m, σ_m)
- So the equation of the capital market line is

$$r = r_f + \frac{\bar{r}_m - r_f}{\sigma_m} \sigma$$

- Sharpe ratio:

$$\frac{\bar{r}_m - r_f}{\sigma_m}$$

Copyright Dick Startz

55

Individual security, weight ω

$$\begin{bmatrix} r_i \\ r_m \end{bmatrix} \sim \left(\begin{bmatrix} \bar{r}_i \\ \bar{r}_m \end{bmatrix}, \begin{bmatrix} \sigma_i^2 & \sigma_{im} \\ \sigma_{im} & \sigma_m^2 \end{bmatrix} \right)$$

$$W \sim \left(\frac{\omega \bar{r}_i + (1 - \omega) \bar{r}_m}{\omega^2 \sigma_i^2 + (1 - \omega)^2 \sigma_m^2 + 2\omega(1 - \omega) \sigma_{im}} \right)$$

Copyright Dick Startz

56

$$W \sim \left(\frac{\omega \bar{r}_i + (1-\omega) \bar{r}_m}{\omega^2 \sigma_i^2 + (1-\omega)^2 \sigma_m^2 + 2\omega(1-\omega) \sigma_{im}} \right)$$

$$\frac{\partial E(W)}{\partial \omega} = \bar{r}_i - \bar{r}_m$$

$$\frac{\partial \text{std}(W)}{\partial \omega} = \frac{2\omega \sigma_i^2 - 2(1-\omega) \sigma_m^2 + 2(1-2\omega) \sigma_{im}}{2 \text{std}(W)}$$

$$\lim_{\omega \rightarrow 0} \frac{\partial \text{std}(W)}{\partial \omega} = \frac{-\sigma_m^2 + \sigma_{im}}{\sigma_m}$$

$$\frac{d E(W)}{d \text{std}(W)} = \frac{\bar{r}_i - \bar{r}_m}{\frac{-\sigma_m^2 + \sigma_{im}}{\sigma_m}} = \frac{\bar{r}_m - r_f}{\sigma_m}$$

Copyright Dick Startz

57

$$\frac{d E(W)}{d \text{std}(W)} = \frac{\bar{r}_i - \bar{r}_m}{\frac{-\sigma_m^2 + \sigma_{im}}{\sigma_m}} = \frac{\bar{r}_m - r_f}{\sigma_m}$$

$$\bar{r}_i - \bar{r}_m = \frac{\bar{r}_m - r_f}{\sigma_m} \times \frac{-\sigma_m^2 + \sigma_{im}}{\sigma_m}$$

• Define

$$\beta = \frac{\sigma_{im}}{\sigma_m^2}$$

$$\bar{r}_i = \bar{r}_m + (\bar{r}_m - r_f)(\beta - 1)$$

$$\bar{r}_i - r_f = \beta(\bar{r}_m - r_f)$$

Copyright Dick Startz

58

Security market line

- β is the regression coefficient of r_i on r_m .
- The variance of security i is irrelevant. This is much more general than this particular model. Diversifiable risk should not be priced.
- You will also see

$$\bar{r}_i - r_f = \alpha + \beta(\bar{r}_m - r_f)$$

Copyright Dick Startz

59

Cauchy-Schwarz Inequality

$$(x \cdot y)^2 \leq (x \cdot x)(y \cdot y)$$

where the dots are inner products.

For example

$$|x_1 y_1 + x_2 y_2|^2 \leq (x_1^2 + x_2^2)(y_1^2 + y_2^2)$$

$$(E(XY))^2 \leq E(X^2) E(Y^2)$$

Let

$$X = x - \mu_x, Y = y - \mu_y$$

$$\left(E((x - \mu_x)(y - \mu_y)) \right)^2 \leq E((x - \mu_x)^2) \times E((y - \mu_y)^2)$$

$$|\text{cov}(x, y)| \leq \sigma_x \sigma_y$$

Copyright Dick Startz

60

Minkowski's inequality

Let X and Y be any two random variables. Then for $1 \leq p < \infty$,

$$(E[|X + Y|^p])^{\frac{1}{p}} \leq (E[|X|^p])^{\frac{1}{p}} + (E[|Y|^p])^{\frac{1}{p}}$$

Copyright Dick Startz

61

Conditional expectations inequality

For $r \geq 1$,

$$E(|E(y|x)|^r) \leq E(|y|^r)$$

Copyright Dick Startz

62

Hierarchical models and mixture distributions


A random variable X is said to have a mixture distribution if the distribution of X depends on a quantity that also has a distribution.

Example: Mixture of normals

$$f(x) = p \cdot \left[\frac{1}{\sqrt{2\pi\sigma_1^2}} e^{-\frac{1}{2\sigma_1^2}(x-\mu_1)^2} \right] + (1-p) \cdot \left[\frac{1}{\sqrt{2\pi\sigma_2^2}} e^{-\frac{1}{2\sigma_2^2}(x-\mu_2)^2} \right]$$

Copyright Dick Startz

63



Copyright Dick Startz

64

Jacobian

$$\mathbf{f}(\mathbf{x}): \mathbb{R}^n \rightarrow \mathbb{R}^m$$

$$\mathbf{J} = \frac{d\mathbf{f}}{d\mathbf{x}} = \begin{bmatrix} \frac{\partial \mathbf{f}}{\partial x_1} & \cdots & \frac{\partial \mathbf{f}}{\partial x_n} \end{bmatrix} = \begin{bmatrix} \frac{\partial f_1}{\partial x_1} & \cdots & \frac{\partial f_1}{\partial x_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial f_m}{\partial x_1} & \cdots & \frac{\partial f_m}{\partial x_n} \end{bmatrix}$$

$$|\mathbf{J}|$$

Copyright Dick Startz

65

Transformation pdf

- **Theorem 2.1.5** Let X have continuous pdf $f_X(x)$ on \mathcal{X} and let $Y = g(X)$, where $g(\cdot)$ is a monotone function and that $g^{-1}(\cdot)$ has a continuous derivative on \mathcal{Y} . Then the pdf of Y is given by

$$f_Y(y) = \begin{cases} f_X(g^{-1}(y)) \left| \frac{d}{dy} g^{-1}(y) \right|, & y \in \mathcal{Y} \\ 0, & \text{otherwise} \end{cases}$$

Copyright Dick Startz

66

Bivariate transformation

Let (X, Y) be a bivariate random vector with joint pdf $f_{X,Y}(x, y)$ and consider the one-to-one functions

$$\begin{aligned} u &= g_u(x, y) \\ v &= g_v(x, y) \end{aligned}$$

With inverse functions

$$\begin{aligned} x &= h_x(u, v) \\ y &= h_y(u, v) \end{aligned}$$

Copyright Dick Startz

67

Change of variables

$$\mathbf{J} = \begin{bmatrix} \frac{\partial h_x}{\partial u} & \frac{\partial h_x}{\partial v} \\ \frac{\partial h_y}{\partial u} & \frac{\partial h_y}{\partial v} \end{bmatrix}$$

$$f_{U,V}(u, v) = f_{X,Y}(h_x(u, v), h_y(u, v)) |\mathbf{J}|$$

where $|\mathbf{J}|$ is the absolute value of the determinant.

Copyright Dick Startz

68

Sum of two independent $U(0,1)$

$$f_{X,Y} = \begin{cases} 1, & 0 \leq x \leq 1, 0 \leq y \leq 1 \\ 0, & \text{otherwise} \end{cases}$$

Transformations

$$\begin{aligned} u &= x + y \\ v &= x - y \end{aligned}$$

Copyright Dick Startz

69

The inverse functions are given by

$$\begin{aligned} x &= h_x(u, v) = \frac{u+v}{2} \\ y &= h_y(x, y) = \frac{u-v}{2} \\ \mathbf{J} &= \begin{vmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & -\frac{1}{2} \end{vmatrix} = -\frac{1}{2} \end{aligned}$$

Copyright Dick Startz

70

Joint density

$$\begin{aligned} & \begin{matrix} 0 \leq u \leq 2 \\ -1 \leq v \leq 1 \end{matrix} \\ f_{X,Y}(h_x(u, v), h_y(u, v)) &= \begin{cases} 1, & 0 \leq u \leq 2, -1 \leq v \leq 1 \\ 0, & \text{otherwise} \end{cases} \\ f_{u,v}(u, v) &= \begin{cases} 1 \times \left| -\frac{1}{2} \right|, & 0 \leq u \leq 2, -1 \leq v \leq 1 \\ 0, & \text{otherwise} \end{cases} \end{aligned}$$

Copyright Dick Startz

71

Marginal distribution of the sum

$$f_U = \int_{-\infty}^{\infty} f_{U,V}(u, v) dv$$

Suppose $0 \leq u \leq 1$, then $-u \leq v \leq u$.


$$f_U = \int_{-u}^u \frac{1}{2} dv = \frac{1}{2} [v]_{-u}^u = u$$

Suppose $1 \leq u \leq 2$, then $-(2-u) \leq v \leq (2-u)$

$$f_U = \int_{-(2-u)}^{(2-u)} \frac{1}{2} dv = \frac{1}{2} [v]_{-(2-u)}^{(2-u)} = 2 - u$$

Copyright Dick Startz

72



Copyright Dick Startz

73

Marginal distribution of the difference

$$f_V = \int_{-\infty}^{\infty} f_{U,V}(u, v) du$$

If $-1 < v < 0$, then let's look at $-2v < u < 2$

$$f_V(v) = \left[\frac{1}{2}u \right]_{-2v}^2 = 1 + v$$

If $0 < v < 1$, then let's look at $2v < u < 2$

$$f_V(v) = \left[\frac{1}{2}u \right]_{2v}^2 = 1 - v$$


Now consider

$$f_{u,v}(1,1) = \frac{1}{2} \neq f_u(1) \times f_v(1) = 1 \times 0 = 0$$

Therefore u, v are not independent.

Copyright Dick Startz

74



Copyright Dick Startz

75

Independence of sum and difference of independent uniforms

Suppose $0 \leq u \leq 1$, then $-u \leq v \leq u$.

$$f_U = \int_{-u}^u \frac{1}{2} dv = \frac{1}{2} [v]_{-u}^u = u$$

If $0 < v < 1$, then let's look at $2v < u < 2$

$$f_V(v) = \left[\frac{1}{2}u \right]_{2v}^2 = 1 - v$$

Now consider

$$f_{u,v}(1,1) = \frac{1}{2} \neq f_u(1) \times f_v(1) = 1 \times 0 = 0$$

Therefore u, v are not independent.

Copyright Dick Startz

76

Sum and difference of independent, standard normals

$$f(x, y) = (2\pi)^{-1} e^{-\frac{1}{2}x^2} e^{-\frac{1}{2}y^2}$$

Transformations

$$\begin{aligned} u &= x + y \\ v &= x - y \end{aligned}$$

Copyright Dick Startz

77

Change of variables

$$f_{U,V}(u, v) = (2\pi)^{-1} e^{-\frac{1}{2}\left(\frac{u+v}{2}\right)^2} e^{-\frac{1}{2}\left(\frac{u-v}{2}\right)^2} \times \left| -\frac{1}{2} \right|$$

$$f_{U,V}(u, v) = (2\pi)^{-1} e^{-\frac{1}{2}\left[\left(\frac{u+v}{2}\right)^2 + \left(\frac{u-v}{2}\right)^2\right]} \times \left| -\frac{1}{2} \right|$$

Copyright Dick Startz

78

Adding exponents

$$\begin{aligned} & -\frac{1}{2}\left(\frac{1}{2}\right)^2 [(u+v)^2 + (u-v)^2] \\ & = -\frac{1}{2}\left(\frac{1}{2}\right)^2 [(u^2 + v^2 + 2uv) + (u^2 + v^2 - \end{aligned}$$

Copyright Dick Startz

79

Joint density

$$\begin{aligned} f_{U,V}(u, v) &= (2\pi\sqrt{2})^{-1} e^{-\frac{1}{2}\left(\frac{u}{\sqrt{2}}\right)^2} e^{-\frac{1}{2}\left(\frac{v}{\sqrt{2}}\right)^2} \\ u, v &\sim i.i.d. N(0, \sqrt{2}) \end{aligned}$$

Quick check:

$$\begin{aligned} \text{var}(x + y) &= \text{var}(x) + \text{var}(y) + 2 \times 0 \\ 1 + 1 &= 2 \end{aligned}$$

Copyright Dick Startz

80

Functions of independent random variables

Let X and Y be independent random variables.
If

$$\begin{aligned} U &= g(X) \\ V &= h(Y) \end{aligned}$$

Then U and V are independent.

Copyright Dick Startz

81

Convolution formula

Theorem 5.2.9 If X and Y are independent continuous random variables with pdfs $f_X(x)$ and $f_Y(y)$, then the pdf of $Z = X + Y$ is

$$f_Z(z) = \int_{-\infty}^{\infty} f_X(w) f_Y(z - w) dw$$

Note: If the limits of integration may be modified if either pdf has a limited support.

Note: Because the sum is symmetric in X and Y , the convolution can also be written

$$f_Z(z) = \int_{-\infty}^{\infty} f_X(z - w) f_Y(w) dw$$

Copyright Dick Startz

82

Sum of independent $x \sim U(0,1)$ and $y \sim U(0,1)$

$$\begin{aligned} f_Z(z) &= \int_{-\infty}^{\infty} f_X(w) f_Y(z - w) dw \\ &= \int_0^1 1 \times f_Y(z - w) dw \end{aligned}$$

$$f_Z(z) = \int_0^1 f_Y(z - w) dw$$

The integrand is zero except if $0 \leq z - w \leq 1$, in which case it is 1. So if $0 \leq z \leq 1$ w can take on any value up to z

$$f_Z(z) = \int_0^z 1 dw = z$$

If z is between 1 and 2 w can be as low as $z - 1$ and as high as 1.

$$f_Z(z) = \int_{z-1}^1 1 dw = 2 - z$$


Otherwise the density equals 0

Copyright Dick Startz

83

Copyright Dick Startz

84



Copyright Dick Startz

85

Sum of independent standard normals

$$f_Z(z) = \int_{-\infty}^{\infty} \phi(w)\phi(z-w)dw$$

We can write the innards as

$$\begin{aligned} & \frac{1}{\sqrt{2\pi}} \exp\left[-\frac{w^2}{2}\right] \times \frac{1}{\sqrt{2\pi}} \exp\left[-\frac{(w-z)^2}{2}\right] \\ &= \frac{1}{2\pi} \times \exp\left(-\frac{z^2}{4}\right) \exp\left(-\left(w-\frac{z}{2}\right)^2\right) \end{aligned}$$

Copyright Dick Startz

86

Sum of independent standard normals

Putting this back into the integral gives

$$\begin{aligned} f_Z(z) &= \frac{1}{\sqrt{2\pi}} \times \exp\left(-\frac{z^2}{4}\right) \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} \times \exp\left(-\left(w-\frac{z}{2}\right)^2\right) dw \\ f_Z(z) &= \frac{1}{\sqrt{2\pi}} \times \exp\left(-\frac{z^2}{4}\right) \sqrt{5} \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} \cdot \frac{1}{\sqrt{5}} \times \exp\left(-\frac{1}{2 \cdot 5} \left(w-\frac{z}{2}\right)^2\right) dw \\ & \quad w \sim N\left(\frac{z}{2}, (\sqrt{5})^2\right) \\ f_Z(z) &= \frac{1}{\sqrt{2\pi}} \times \exp\left(-\frac{z^2}{4}\right) \sqrt{5} = \frac{1}{\sqrt{2\pi \cdot 5}} \exp\left(-\frac{1}{2 \times 5} z^2\right) \sim N(0,2) \end{aligned}$$

Copyright Dick Startz

87

Bivariate normal

If X and Y are joint normal, with means μ_x and μ_y , variances σ_x^2 and σ_y^2 , and correlation ρ , we can write the joint density as

$$\begin{aligned} f(x,y) &= \left(2\pi\sigma_x\sigma_y\sqrt{1-\rho^2}\right)^{-1} \\ & \times e^{-\frac{1}{2(1-\rho^2)}\left(\left(\frac{x-\mu_x}{\sigma_x}\right)^2 + \left(\frac{y-\mu_y}{\sigma_y}\right)^2 - 2\rho\left(\frac{x-\mu_x}{\sigma_x}\right)\left(\frac{y-\mu_y}{\sigma_y}\right)\right)} \end{aligned}$$

Copyright Dick Startz

88

Nice properties

- a) The marginal distribution of X is $N(\mu_x, \sigma_x^2)$.
- b) The marginal distribution of Y is $N(\mu_y, \sigma_y^2)$.
- c) For any constants a and b , the distribution of $ax + by$ is $N(a\mu_x + b\mu_y, a^2\sigma_x^2 + b^2\sigma_y^2 + 2ab\rho\sigma_x\sigma_y)$.

Copyright Dick Startz

89

Uncorrelated normals are independent

$$f(x, y) = \left(2\pi\sigma_x\sigma_y\sqrt{1-\rho^2}\right)^{-1} \times e^{-\frac{1}{2(1-\rho^2)}\left(\left(\frac{x-\mu_x}{\sigma_x}\right)^2 + \left(\frac{y-\mu_y}{\sigma_y}\right)^2 - 2\rho\left(\frac{x-\mu_x}{\sigma_x}\right)\left(\frac{y-\mu_y}{\sigma_y}\right)\right)}$$

Copyright Dick Startz

90

Variance-covariance matrix

Covariance

$$\sigma_{xy} = \rho\sigma_x\sigma_y$$

Variance-covariance matrix

$$\Sigma = \begin{bmatrix} \sigma_x^2 & \sigma_{xy} \\ \sigma_{xy} & \sigma_y^2 \end{bmatrix}$$

$$f(x, y) = (2\pi)^{-\frac{2}{2}}|\Sigma|^{-\frac{1}{2}}\exp\left(-\frac{1}{2}\begin{bmatrix} x & y \end{bmatrix}\Sigma^{-1}\begin{bmatrix} x \\ y \end{bmatrix}\right)$$

Copyright Dick Startz

91

Conditional normals

- $y|x \sim N, x|y \sim N$

$$E(y|x) = \mu_y + \frac{\sigma_{xy}}{\sigma_x^2}(x - \mu_x)$$

$$E(x|y) = \mu_x + \frac{\sigma_{xy}}{\sigma_y^2}(y - \mu_y)$$

$$\text{var}(y|x) = \sigma_y^2(1 - \rho^2)$$

$$\text{var}(x|y) = \sigma_x^2(1 - \rho^2)$$

$$\frac{\text{var } E(y|x)}{\text{var}(y)} = \rho^2$$

Copyright Dick Startz

92

Conditional normals

- Theory

$$E(y|x) = \mu_y + \frac{\sigma_{xy}}{\sigma_x^2} (x - \mu_x)$$

- Regression

$$\hat{\beta} = \frac{\sum (x - \bar{x})(y - \bar{y})}{\sum (x - \bar{x})^2}$$

Copyright Dick Startz

93

$$\frac{\text{var } E(y|x)}{\text{var}(y)} = \rho^2 \text{ regression analogy}$$

$$y = \beta x + \varepsilon$$

The forecast is

$$E(y|x) = \beta x$$

The explained variation is

$$\text{var}(E(y|x))$$

So the R^2 is

$$R^2 = \frac{\text{var}(E(y|x))}{\text{var}(y)}$$

Copyright Dick Startz

94

Question for today

Suppose that employers observe a signal of productivity, mp_i , with error.

$$s_i = mp_i + \varepsilon_i$$

where marginal product and the error are joint normal, and uncorrelated, and the mean error is zero. So the error is pure noise. If employers observe the signal for individuals and pay expected marginal product, what does the wage schedule look like, and in particular what happens to an individual's wage if they increase their productivity by 1.0?

Copyright Dick Startz

95

Simple lemons

Suppose used cars have a value distribution $f(v)$. You know the value of your car, v_i , but no one else does. For a cost c you can have the value of your car creditably certified by a mechanic. If you do, the car will sell for v . If not the car will sell for the average value of uncertified cars.

Copyright Dick Startz

99

Marginal lemon

- Suppose we call the value at which certification is marginal v^* . Then uncertified cars sell for

$$E(v|v < v^*)$$

Or

$$\int_{-\infty}^{v^*} v f(v|v \leq v^*) dv = \int_{-\infty}^{v^*} \frac{v f(v)}{\int_{-\infty}^{v^*} f(v) dv} dv$$

Copyright Dick Startz

100

Gain from certification

$$v_i - \int_{-\infty}^{v^*} \frac{v f(v)}{\int_{-\infty}^{v^*} f(v) dv} dv - c$$

$$0 = v^* - \int_{-\infty}^{v^*} \frac{v f(v)}{\int_{-\infty}^{v^*} f(v) dv} dv - c$$

Copyright Dick Startz

101

Uniform example

$$v \sim U[0, b]$$

$$f(v) = \frac{1}{b}$$

$$\int_{-\infty}^{v^*} f(v) dv = F(v^*) = \frac{1}{b} [v]_0^{v^*} = \frac{v^*}{b}$$

So conditional expectation is:

$$\frac{1}{(v^*/b)} \int_{-\infty}^{v^*} v f(v) dv = \frac{1}{(v^*/b)} \frac{1}{2b} [v^2]_0^{v^*} = \frac{v^{*2}}{2v^*} = \frac{v^*}{2}$$

Copyright Dick Startz

102

Uniform example

$$0 = v^* - \frac{v^*}{2} - c$$

$$v^* = 2c$$

Copyright Dick Startz

103

Multivariate normal

$$\begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \sim N \left(\begin{bmatrix} \mu_1 \\ \mu_2 \\ \vdots \\ \mu_n \end{bmatrix}, \Sigma = \begin{bmatrix} \sigma_1^2 & \sigma_{12} & \cdots & \sigma_{1n} \\ \sigma_{12} & \sigma_2^2 & \cdots & \\ \vdots & \vdots & \ddots & \\ \sigma_{1n} & & & \sigma_n^2 \end{bmatrix} \right)$$

$$f(x) = (2\pi)^{-\frac{n}{2}} |\Sigma|^{-\frac{1}{2}} \exp \left(-\frac{1}{2} (x - \mu)' \Sigma^{-1} (x - \mu) \right)$$

Copyright Dick Startz

104

Marginal and conditional distributions

$$\vec{x}_1 \sim N(\vec{\mu}_1, \Sigma_1)$$

$$E(\vec{x}_1 | \vec{x}_2) = \vec{\mu}_1 + \Sigma_{12} \Sigma_2^{-1} (\vec{x}_2 - \vec{\mu}_2)$$

Copyright Dick Startz

105

Conditional distribution and regression

• Theory

$$E(\vec{x}_1 | \vec{x}_2) = \vec{\mu}_1 + \Sigma_{12} \Sigma_2^{-1} (\vec{x}_2 - \vec{\mu}_2)$$

• Regression

$$\hat{\beta} \leftarrow (X'X)^{-1} X'y$$

Copyright Dick Startz

106

Linear combinations of joint normals

If L is an $m \times k$ matrix of constants then

$$Lx \sim N(L\mu, L\Sigma L')$$

Copyright Dick Startz

107

Useful application of linear combination of normals

$$y = X\beta + \varepsilon, \varepsilon \sim N(0, \sigma^2 I)$$

where X is an $n \times k$ fixed matrix.

$$\begin{aligned}\hat{\beta} &\equiv (X'X)^{-1}X'y \\ &= (X'X)^{-1}X'X\beta + (X'X)^{-1}X'\varepsilon \\ &= \beta + (X'X)^{-1}X'\varepsilon\end{aligned}$$

$$\begin{aligned}\hat{\beta} &\sim N(\beta, (X'X)^{-1}X'\sigma^2 I X(X'X)^{-1}) \\ &= N(\beta, \sigma^2 (X'X)^{-1})\end{aligned}$$

Copyright Dick Startz

108

Vector bivariate normal

If

$$\begin{bmatrix} x \\ y \end{bmatrix} \sim N \left(\begin{bmatrix} \mu_x \\ \mu_y \end{bmatrix}, \begin{bmatrix} \Sigma_{xx} & \Sigma_{xy} \\ \Sigma_{xy} & \Sigma_{yy} \end{bmatrix} \right)$$

Then x and y are independent iff $\Sigma_{xy} = 0$.

Linear combinations of x and y can be written

$$\begin{bmatrix} L_x & 0 \\ 0 & L_y \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} \sim N \left(\begin{bmatrix} L_x \mu_x \\ L_y \mu_y \end{bmatrix}, \begin{bmatrix} L_x \Sigma_{xx} L_x' & L_x \Sigma_{xy} L_y' \\ L_x \Sigma_{xy} L_y' & L_y \Sigma_{yy} L_y' \end{bmatrix} \right)$$

$L_x x$ and $L_y y$ are independent iff their covariance, $L_x \Sigma_{xy} L_y'$, equals zero.

Copyright Dick Startz

109

Scalar Cholesky factorization

Cholesky factorization of σ^2 is σ . If

$$\varepsilon \sim N(0, 1)$$

then

$$\begin{aligned}x &= \sigma \varepsilon \sim N(0, \sigma^2) \\ x \times \sigma^{-1} &\sim N(0, 1)\end{aligned}$$

Copyright Dick Startz

110

Cholesky factorization

$\exists C$ s.t. C is lower triangular, nonsingular.

$$CC' = \Sigma$$

$$C = \begin{bmatrix} \sigma_1 & 0 \\ \frac{\sigma_{12}}{\sigma_1} & \sigma_2^2 - \left(\frac{\sigma_{12}}{\sigma_1}\right)^2 \end{bmatrix}$$

n.b. Matlab does upper triangular, $C'C = \Sigma$

Copyright Dick Startz

111

If ε is distributed i.i.d. standard normal,
 $\varepsilon \sim N(0, I)$,
 $C\varepsilon \sim N(C \times 0, CIC') = N(0, \Sigma)$

If $x \sim N(0, \Sigma)$,
 $C^{-1}x \sim N(0, C^{-1}\Sigma C'^{-1}) = N(0, C^{-1}[CC']C'^{-1})$
 $= N(0, [C^{-1}C][C'C'^{-1}]) = N(0, I)$

Copyright Dick Startz

112

Example

If $\varepsilon_1, \varepsilon_2$ are i.i.d standard normal, and

$$x_1 = \sigma_1 \varepsilon_1$$

$$x_2 = \frac{\sigma_{12}}{\sigma_1} \varepsilon_1 + \left(\sigma_2^2 - \left(\frac{\sigma_{12}}{\sigma_1} \right)^2 \right) \varepsilon_2$$

then

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \sim N\left(0, \begin{bmatrix} \sigma_1^2 & \sigma_{12} \\ \sigma_{12} & \sigma_2^2 \end{bmatrix}\right)$$

Copyright Dick Startz

113

$$\chi^2$$

Theorem: The sum of the squares of n independent standard normals is distributed χ_n^2 .

Examples:

If $\varepsilon \sim iid N(0,1)$, then $\sum \varepsilon^2 \sim \chi_n^2$.

If $x_i \sim ind N(\mu_i, \sigma_i^2)$, then $\sum \left(\frac{x_i - \mu_i}{\sigma_i} \right)^2 \sim \chi_n^2$.

Copyright Dick Startz

114

$$\chi_d^2 \text{ or } \chi^2(d)$$

$$f(x) = \frac{1}{2^{\frac{d}{2}} \Gamma\left(\frac{d}{2}\right)} x^{\frac{d}{2}-1} e^{-\frac{x}{2}}$$

$$E(x) = d$$

$$\text{var}(x) = 2d$$

- If $\varepsilon \sim N(0,1)$, $E(\varepsilon^2) = 1$

Copyright Dick Startz

115

Class assignment

Prove that if $x \sim \chi_1^2$, \sqrt{x} is distributed standard half-normal, that is, like a standard normal distribution defined only on a non-negative support.

Copyright Dick Startz

116

Sum of independent χ^2 is χ^2

If $x_i \sim \text{ind} \chi_{k_i}^2$, then

$$x_1 + \dots + x_k \sim \chi_{\sum k_i}^2$$

Copyright Dick Startz

117

Example

$$y_i, i = 1, \dots, n \sim N(\mu_y, \sigma^2)$$

μ_y is known

$$H_0: \sigma^2 = \sigma_0^2$$

$$\sum \left(\frac{y_i - \mu_y}{\sigma_0} \right)^2 \sim \chi_n^2$$

Copyright Dick Startz

118

t-distribution


If $x \sim N(0,1)$ and $S \sim \chi_d^2$ and x, S are independent

$$\frac{x}{\sqrt{S/d}} \sim t_d$$

$$\lim_{n \rightarrow \infty} t_n = N(0,1)$$

Copyright Dick Startz

119



Copyright Dick Startz

120

F-distribution

If $s_1 \sim \chi^2_{d_1}$ and $s_2 \sim \chi^2_{d_2}$, s_1, s_2 independent

$$\frac{s_1/d_1}{s_2/d_2} \sim F(d_1, d_2)$$

$$t_d^2 = F(1, d)$$

$$\lim_{d_2 \rightarrow \infty} F(d_1, d_2) = \chi^2_{d_1}/d_1$$

Copyright Dick Startz

121