

Conditional probability

Using Densities

If A and B are events in S , and $P(B) > 0$, then the *conditional probability* of A given B , written $P(A|B)$, is

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

Copyright Dick Startz

1

Copyright Dick Startz

2

Conditional probability example

$$\begin{aligned} S &= \{1, 2, 3, 4, 5, 6\} \\ A &= \{2\}, P(A) = \frac{1}{6} \\ B &= \{2\} \cup \{4\} \cup \{6\}, P(B) = \frac{1}{2} \\ A \cap B &= \{2\}, P(A \cap B) = \frac{1}{6} \\ P(A|B) &= \frac{P(A \cap B)}{P(B)} = \frac{1/6}{1/2} = \frac{1}{3} \end{aligned}$$

Copyright Dick Startz

3

Another example

Type	probability
Productive, talks	.4
Productive, quiet	.2
Useless, talks	.1
Useless, quiet	.3

Now suppose that the value of a productive RA is 1 and the value of an unproductive RA is zero and that the cost of an RA is 0.7 and that you want to maximize

$$V = P(\text{Prod}) \times (1 - .7) + P(\text{Unprod}) \times (0 - .7)$$

Copyright Dick Startz

4

Insurance market

Consider a health insurance plan. If a person is sick, the plan will have to spend C . A healthy person costs nothing. The probability that a young person will be sick is p_y ; for an old person p_o , where $p_o > p_y$. A fraction f of the population is young. People voluntarily buy insurance if the premium is equal to or below their expected cost of illness. The insurance company prices the plan so as to break even.

- A. Suppose participation is mandatory. What will be the premium?
- B. Suppose participation is voluntary. What will be the premium?

Copyright Dick Startz

7

Bayes rule

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

$$P(A \cap B) = P(A|B)P(B)$$

$$P(A \cap B) = P(B \cap A) = P(B|A)P(A)$$

Equating these gives

$$P(A|B) = P(B|A) \frac{P(A)}{P(B)}$$

Copyright Dick Startz

9

Bayes rule

Let A_1, A_2, \dots be a partition of the sample space, and let B be any set. Then $\forall i$

$$P(A_i|B) = \frac{P(B|A_i)P(A_i)}{\sum_{i=1}^{\infty} P(B|A_i)P(A_i)}$$

Copyright Dick Startz

10

Somewhat true facts about tests for tuberculosis

- Tuberculosis is a serious disease that is on the increase.
- Incidence is about 2 people per 10,000 (varies across countries and time)
- Inexpensive tests are pretty accurate.
 - Probability of a false positive is 2 out of 100
 - Probability of a false negative is 1 out of 100

Copyright Dick Startz

11

Example of tuberculosis test

- B is outcome of tuberculosis test
 - $B \in \{yes, no\}$
- A is whether patient has tuberculosis
 - $A \in \{yes, no\}$
 - $p(A = yes) = 2/10,000$
- $p(B = yes|A = yes) = .99$
- $p(B = yes|A = no) = 0.02$

What is $p(A = yes|B = yes)$, i.e. tuberculosis if test is positive?

Copyright Dick Startz

12

Example of tuberculosis test, cont.

$$P(\text{tuber}|\text{pos}) = \frac{P(\text{pos}|\text{tuber}) \times P(\text{tuber})}{P(\text{pos}|\text{tuber})P(\text{tuber}) + P(\text{pos}|\sim\text{tuber})P(\sim\text{tuber})}$$

$$P(\text{tuber}|\text{pos}) = \frac{.99 \times .0002}{.99 \times .0002 + .02 \times .9998} = 0.0098$$

Copyright Dick Startz

13

Independence

Two events, A and B are *statistically independent* if

$$P(A \cap B) = P(A) \times P(B)$$

Intuition:

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

$$P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{P(A)P(B)}{P(B)} = P(A)$$

Copyright Dick Startz

14

Example of not independent

Type	probability
Productive, talks	.4
Productive, quiet	.2
Useless, talks	.1
Useless, quiet	.3

$$P(\text{Prod} \cap \text{Talks}) = .4$$

$$P(\text{Prod}) = .6$$

$$P(\text{Talks}) = .5$$

$$.4 \neq .6 \times .5$$

Copyright Dick Startz

15

Further implications of independence

If A and B are independent events, then the following pairs are also independent:

- a) A and B^c
- b) A^c and B
- c) A^c and B^c

Copyright Dick Startz

16

Random variable

A *random variable* is a function from a sample space S into the real numbers.

The *cumulative distribution function* or *cdf* of a random variable X , denoted by $F_X(x)$, is defined by

$$F_X(x) = P_X(X \leq x), \forall x$$

Copyright Dick Startz

17

Sum of Two Dice CDF

$$\begin{aligned} F(1) &= 0 \\ F(2) &= \frac{1}{36} \\ F(3) &= \frac{1}{36} + \frac{2}{36} \\ &\vdots \\ F(12) &= 1 \end{aligned}$$

Copyright Dick Startz

18


CDF properties

The function $F(x)$ is a cdf iff the following three conditions hold:

- a) $\lim_{x \rightarrow -\infty} F(x) = 0$ and $\lim_{x \rightarrow \infty} F(x) = 1$.
- b) $F(x)$ is a nondecreasing function of x .
- c) $F(x)$ is right-continuous: i.e.,
 $\forall x_0, \lim_{x \downarrow x_0} F(x) = F(x_0)$.

Copyright Dick Startz

19



Copyright Dick Startz

20

Logistic cdf

$$F_X(x) = \frac{1}{1 + e^{-\frac{(x-\mu)}{s}}}, s > 0$$

$$a) \lim_{x \rightarrow -\infty} F(x) = 0 \text{ since } \lim_{x \rightarrow -\infty} e^{-x} = \infty \text{ and } \lim_{x \rightarrow \infty} F(x) = 1 \text{ since } \lim_{x \rightarrow \infty} e^{-x} = 0.$$


b) $F(x)$ is a nondecreasing function of x since

$$\frac{d}{dx} F_X(x) = \frac{e^{-\frac{(x-\mu)}{s}}}{s(1 + e^{-\frac{(x-\mu)}{s}})^2} > 0$$

c) Is continuous and therefore right-continuous.

Copyright Dick Startz

21



Copyright Dick Startz

22

Identically distributed random variables

The random variables X and Y are identically distributed if $F_X(x) = F_Y(x) \forall x$

Copyright Dick Startz

23

Density functions

The *probability mass function* for a discrete distribution is

$$f_X(x) = P(X = x)$$

$$F_X(x) = \sum_{k=-\infty}^x f_X(x)$$

The *probability density function* a continuous distribution is

$$f_X(x) = \frac{d}{dx} F_X(x)$$

$$F_X(x) = \int_{-\infty}^x f_X(x) dx$$

Copyright Dick Startz

24

Logistic density, $\mu = 0, s = 1$




Figure 1.6.1. Area under logistic curve

Copyright Dick Startz

25

Expected value

$$E(x) = \sum_{-\infty}^x x \cdot f(x)$$

$$E(x) = \int_{-\infty}^x x \cdot f(x) dx$$

$$E(bet) = -1 \times p + 1 \times (1 - p)$$

$$= 1 \times (1 - 2p)$$

Copyright Dick Startz

26

Variance

$$\text{var}(x) = \sum (x - E(x))^2 \cdot f(x)$$

$$\text{var}(x) = \int_{-\infty}^x (x - E(x))^2 \cdot f(x) dx$$

$$\text{standard deviation} \equiv \sqrt{\text{var}}$$

$$E(bet) = B(1 - 2p)$$

$$\text{var}(bet) = (-B - B(1 - 2p))^2 p + (B - B(1 - 2p))^2 (1 - p)$$

$$\text{If } p = \frac{1}{2}, \text{var}(bet) = B^2, \text{std dev}(bet) = B$$

Copyright Dick Startz

27

Bernoulli

$$f(x) = \begin{cases} p, & x = 1 \\ 1 - p, & x = 0 \\ 0, & \text{otherwise} \end{cases}$$

$$f(x|p) = p^x(1-p)^{1-x}, x \in \{0,1\}$$

$$E(x) = 1 \times p + 0 \times (1-p) = p$$

$$\text{var}(x) = (1-p)^2 \times p + (0-p)^2 \times (1-p)$$

$$= p(1-p)$$

Copyright Dick Startz

28

Support

The **support** of a distribution is the set of values for which the probability is positive.

The support of the Bernoulli is $\{0,1\}$.

Copyright Dick Startz

29

Binomial distribution

$$f(x|n, p) = \binom{n}{x} p^x (1-p)^{n-x}$$

$$E(x) = np$$

$$\text{var}(x) = np(1-p)$$

Copyright Dick Startz

30

Random outcome of a test statistic

Consider a test statistic that rejects a true null hypothesis a fraction $p = 0.05$ of the time. Suppose we have $n = 20$ independent applications of this test. Draw a plot of the pdf of the number of rejections (an application of the binomial distribution). What is the probability of more than 1 rejection?

Copyright Dick Startz

31