

You Perturb It, You Pay According to $\mathcal{K}(A)$: Norms and Error Analysis

Nate DeMaagd, Kurt O'Hearn

MTH 499-02

March 25, 2013

- Norms: properties, types, examples

- Norms: properties, types, examples
- Theorems, results on norms

- Norms: properties, types, examples
- Theorems, results on norms
- Relative/absolute error in perturbations, conditioning number

Definition

Let V be a vector space. A norm is a function $\|\cdot\| : V \rightarrow \mathbb{R}^+$ that has the following properties:

- $\|x\| > 0$ if $x \neq 0$, $x \in V$
- $\|\lambda x\| = |\lambda| \|x\|$ if $\lambda \in \mathbb{R}$, $x \in V$
- $\|x + y\| \leq \|x\| + \|y\|$ for all $x, y \in V$ (triangle inequality)

Definition

Let V be a vector space. A norm is a function $\|\cdot\| : V \rightarrow \mathbb{R}^+$ that has the following properties:

- $\|x\| > 0$ if $x \neq 0$, $x \in V$
- $\|\lambda x\| = |\lambda| \|x\|$ if $\lambda \in \mathbb{R}$, $x \in V$
- $\|x + y\| \leq \|x\| + \|y\|$ for all $x, y \in V$ (triangle inequality)

Other comments:

- Norm can be thought of as generalization of absolute value
- Interpretation: vector length in \mathbb{R}^2 , \mathbb{R}^3

Vector Norms:

Vector Norms:

- l_k norm

$$l_k = \|x\|_k = \left(\sum_{i=1}^n |x_i|^k \right)^{1/k} \quad \text{where } k \in \{1, 2, \dots\} \text{ and } x = (x_1, \dots, x_n)^T$$

Vector Norms:

- l_k norm

$$l_k = \|x\|_k = \left(\sum_{i=1}^n |x_i|^k \right)^{1/k} \quad \text{where } k \in \{1, 2, \dots\} \text{ and } x = (x_1, \dots, x_n)^T$$

- Some common instances of the l_k norm:

k	l_k
1	$\ x\ _1 = \sum_{i=1}^n x_i $
2	$\ x\ _2 = \left(\sum_{i=1}^n x_i^2 \right)^{1/2}$
∞	$\ x\ _\infty = \max_{1 \leq i \leq n} x_i $

- Consider vectors in \mathbb{R}^3

$$x = (2, -1, 3)^T \quad y = (7, 2, 5)^T \quad z = (-3, 1, -9)^T$$

	$\ \cdot\ _1$	$\ \cdot\ _2$	$\ \cdot\ _\infty$
x			
y			
z			

- Consider vectors in \mathbb{R}^3

$$x = (2, -1, 3)^T \quad y = (7, 2, 5)^T \quad z = (-3, 1, -9)^T$$

	$\ \cdot\ _1$	$\ \cdot\ _2$	$\ \cdot\ _\infty$
x	6		
y			
z			

$$\|x\|_1 = 2 + 1 + 3 = 6$$

- Consider vectors in \mathbb{R}^3

$$x = (2, -1, 3)^T \quad y = (7, 2, 5)^T \quad z = (-3, 1, -9)^T$$

	$\ \cdot\ _1$	$\ \cdot\ _2$	$\ \cdot\ _\infty$
x	6	3.74	
y			
z			

$$\|x\|_2 = \sqrt{2^2 + 1^2 + 3^2} = 3.74$$

- Consider vectors in \mathbb{R}^3

$$x = (2, -1, 3)^T \quad y = (7, 2, 5)^T \quad z = (-3, 1, -9)^T$$

	$\ \cdot\ _1$	$\ \cdot\ _2$	$\ \cdot\ _\infty$
x	6	3.74	3
y			
z			

$$\|x\|_\infty = 3$$

- Consider vectors in \mathbb{R}^3

$$x = (2, -1, 3)^T \quad y = (7, 2, 5)^T \quad z = (-3, 1, -9)^T$$

	$\ \cdot\ _1$	$\ \cdot\ _2$	$\ \cdot\ _\infty$
x	6	3.74	3
y	14	8.83	7
z	13	9.54	9

Definition

A matrix norm subordinate to a vector norm $\|\cdot\|$ is defined as

$$\|A\| = \sup \{ \|Au\| : u \in \mathbb{R}^n, \|u\| = 1 \},$$

where A is an $n \times n$ matrix.

Definition

A matrix norm subordinate to a vector norm $\|\cdot\|$ is defined as

$$\|A\| = \sup \{ \|Au\| : u \in \mathbb{R}^n, \|u\| = 1 \},$$

where A is an $n \times n$ matrix.

- Two common subordinate matrix norms

Definition

A matrix norm subordinate to a vector norm $\|\cdot\|$ is defined as

$$\|A\| = \sup \{ \|Au\| : u \in \mathbb{R}^n, \|u\| = 1 \},$$

where A is an $n \times n$ matrix.

- Two common subordinate matrix norms
 - Subordinate to vector norm $\|\cdot\|_\infty$ is

$$\|A\|_\infty = \max_{1 \leq i \leq n} \sum_{j=1}^n |a_{ij}|,$$

which is the maximum absolute row sum of A .

Definition

A matrix norm subordinate to a vector norm $\|\cdot\|$ is defined as

$$\|A\| = \sup \{ \|Au\| : u \in \mathbb{R}^n, \|u\| = 1 \},$$

where A is an $n \times n$ matrix.

- Two common subordinate matrix norms

- Subordinate to vector norm $\|\cdot\|_\infty$ is

$$\|A\|_\infty = \max_{1 \leq i \leq n} \sum_{j=1}^n |a_{ij}|,$$

which is the maximum absolute row sum of A .

- Subordinate to vector norm $\|\cdot\|_2$ is

$$\begin{aligned} \|A\|_2 &= \sup_{\|x\|_2=1} \{\|Ax\|_2\} \\ &= \sqrt{\rho(A^T A)}, \end{aligned}$$

where $\rho(A^T A)$ is spectral radius of $A^T A$, largest eigenvalue of $A^T A$.

- Consider matrix $A = \begin{bmatrix} 1 & 1 \\ 2 & 1 \end{bmatrix}$

- Consider matrix $A = \begin{bmatrix} 1 & 1 \\ 2 & 1 \end{bmatrix}$
- $(A^T A) = \begin{bmatrix} 5 & 3 \\ 3 & 2 \end{bmatrix}$

- Consider matrix $A = \begin{bmatrix} 1 & 1 \\ 2 & 1 \end{bmatrix}$
- $(A^T A) = \begin{bmatrix} 5 & 3 \\ 3 & 2 \end{bmatrix}$
- For $A^T A$, eigenvalues are $\sigma_1^2 = 6.8541$ and $\sigma_2^2 = 0.1459$.

- Consider matrix $A = \begin{bmatrix} 1 & 1 \\ 2 & 1 \end{bmatrix}$
- $(A^T A) = \begin{bmatrix} 5 & 3 \\ 3 & 2 \end{bmatrix}$
- For $A^T A$, eigenvalues are $\sigma_1^2 = 6.8541$ and $\sigma_2^2 = 0.1459$.
- So, $\|A\|_2 = \sqrt{6.8541} \approx 2.618$.

- Consider matrix $B = \begin{bmatrix} 3 & 2 & 4 \\ 2 & 0 & 2 \\ 4 & 2 & 3 \end{bmatrix}$

- Consider matrix $B = \begin{bmatrix} 3 & 2 & 4 \\ 2 & 0 & 2 \\ 4 & 2 & 3 \end{bmatrix}$
- $\|B\|_2$ is simply the maximum absolute row sum, which is 9.

Theorem (Subordinate Matrix Norm)

If $\|\cdot\|$ is any norm in \mathbb{R}^n , then the equation

$$\|A\| = \sup_{\|u\|=1} \{\|Au\| : u \in \mathbb{R}^n\}$$

defines a norm on the linear space of all $n \times n$ matrices.

Theorem (Subordinate Matrix Norm)

If $\|\cdot\|$ is any norm in \mathbb{R}^n , then the equation

$$\|A\| = \sup_{\|u\|=1} \{\|Au\| : u \in \mathbb{R}^n\}$$

defines a norm on the linear space of all $n \times n$ matrices.

Proof.

TODO □

Theorem (Infinity Matrix Norm)

If the vectorm norm $\|\cdot\|_\infty$ is defined by

$$\|x\|_\infty = \max_{1 \leq i \leq n} |x_i|$$

then its subordinate matrix norm is given by

$$\|A\|_\infty = \max_{1 \leq i \leq n} \sum_{j=1}^n |a_{ij}|.$$

Theorem (Infinity Matrix Norm)

If the vectorm norm $\|\cdot\|_\infty$ is defined by

$$\|x\|_\infty = \max_{1 \leq i \leq n} |x_i|$$

then its subordinate matrix norm is given by

$$\|A\|_\infty = \max_{1 \leq i \leq n} \sum_{j=1}^n |a_{ij}|.$$

Proof.

TODO □

TODO

The Cholesky Factorization

The Cholesky Factorization

- Given A , we can write A as $A = LL^T$ with L lower triangular

The Cholesky Factorization

- Given A , we can write A as $A = LL^T$ with L lower triangular
- Necessary conditions on A :

The Cholesky Factorization

- Given A , we can write A as $A = LL^T$ with L lower triangular
- Necessary conditions on A :
 - Real-valued entries ($A \in \mathbb{R}^{n \times n}$)
 - Symmetric ($A = A^T$)
 - Positive definite ($x^T Ax > 0$ for all $x \in \mathbb{R}^n$, $x \neq 0_n$)

The Cholesky Factorization

- Given A , we can write A as $A = LL^T$ with L lower triangular
- Necessary conditions on A :
 - Real-valued entries ($A \in \mathbb{R}^{n \times n}$)
 - Symmetric ($A = A^T$)
 - Positive definite ($x^T Ax > 0$ for all $x \in \mathbb{R}^n$, $x \neq 0_n$)
- Cholesky: special case of LU -factorization

The Cholesky Factorization

- Given A , we can write A as $A = LL^T$ with L lower triangular
- Necessary conditions on A :
 - Real-valued entries ($A \in \mathbb{R}^{n \times n}$)
 - Symmetric ($A = A^T$)
 - Positive definite ($x^T Ax > 0$ for all $x \in \mathbb{R}^n$, $x \neq 0_n$)
- Cholesky: special case of LU -factorization
- Why are factorizations important? Connections later in pivoting

The Cholesky Factorization

- Given A , we can write A as $A = LL^T$ with L lower triangular
- Necessary conditions on A :
 - Real-valued entries ($A \in \mathbb{R}^{n \times n}$)
 - Symmetric ($A = A^T$)
 - Positive definite ($x^T Ax > 0$ for all $x \in \mathbb{R}^n$, $x \neq 0_n$)
- Cholesky: special case of LU -factorization
- Why are factorizations important? Connections later in pivoting

Theorem

If A is a real, symmetric, and positive definite matrix, then $A = LL^T$ for a unique upper triangular L .

Theorem

If A is a real, symmetric, and positive definite matrix, then $A = LL^T$ for a unique upper triangular L .

Proof.

Existence:

Theorem

If A is a real, symmetric, and positive definite matrix, then $A = LL^T$ for a unique upper triangular L .

Proof.

Existence:

- A positive definite $\Rightarrow A$ nonsingular

Theorem

If A is a real, symmetric, and positive definite matrix, then $A = LL^T$ for a unique upper triangular L .

Proof.

Existence:

- A positive definite $\Rightarrow A$ nonsingular
- Leading principle minors of A positive definite
 \Rightarrow principle minors nonsingular $\Rightarrow A = LU$ (LU factorization exists)

Theorem

If A is a real, symmetric, and positive definite matrix, then $A = LL^T$ for a unique upper triangular L .

Proof.

Existence:

- A positive definite $\Rightarrow A$ nonsingular
- Leading principle minors of A positive definite
 \Rightarrow principle minors nonsingular $\Rightarrow A = LU$ (LU factorization exists)
- $LU = A = A^T = U^T L^T \Leftrightarrow U(L^T)^{-1} = L^{-1}U^T$

Theorem

If A is a real, symmetric, and positive definite matrix, then $A = LL^T$ for a unique upper triangular L .

Proof.

Existence:

- A positive definite $\Rightarrow A$ nonsingular
- Leading principle minors of A positive definite
 \Rightarrow principle minors nonsingular $\Rightarrow A = LU$ (LU factorization exists)
- $LU = A = A^T = U^T L^T \Leftrightarrow U(L^T)^{-1} = L^{-1}U^T$
- $U(L^T)^{-1}$ upper triangular, $L^{-1}U^T$ lower triangular
 \Rightarrow both sides diagonal (D)

Theorem

If A is a real, symmetric, and positive definite matrix, then $A = LL^T$ for a unique upper triangular L .

Proof.

Existence:

- A positive definite $\Rightarrow A$ nonsingular
- Leading principle minors of A positive definite
 \Rightarrow principle minors nonsingular $\Rightarrow A = LU$ (LU factorization exists)
- $LU = A = A^T = U^T L^T \Leftrightarrow U(L^T)^{-1} = L^{-1}U^T$
- $U(L^T)^{-1}$ upper triangular, $L^{-1}U^T$ lower triangular
 \Rightarrow both sides diagonal (D)
- $U = DL^T$

Theorem

If A is a real, symmetric, and positive definite matrix, then $A = LL^T$ for a unique upper triangular L .

Proof.

Existence:

- A positive definite $\Rightarrow A$ nonsingular
- Leading principle minors of A positive definite
 \Rightarrow principle minors nonsingular $\Rightarrow A = LU$ (LU factorization exists)
- $LU = A = A^T = U^T L^T \Leftrightarrow U(L^T)^{-1} = L^{-1}U^T$
- $U(L^T)^{-1}$ upper triangular, $L^{-1}U^T$ lower triangular
 \Rightarrow both sides diagonal (D)
- $U = DL^T$
- D has positive diagonal entries $\Rightarrow D = D^{1/2}D^{1/2}$

Theorem

If A is a real, symmetric, and positive definite matrix, then $A = LL^T$ for a unique upper triangular L .

Proof.

Existence:

- A positive definite $\Rightarrow A$ nonsingular
- Leading principle minors of A positive definite
 \Rightarrow principle minors nonsingular $\Rightarrow A = LU$ (LU factorization exists)
- $LU = A = A^T = U^T L^T \Leftrightarrow U(L^T)^{-1} = L^{-1}U^T$
- $U(L^T)^{-1}$ upper triangular, $L^{-1}U^T$ lower triangular
 \Rightarrow both sides diagonal (D)
- $U = DL^T$
- D has positive diagonal entries $\Rightarrow D = D^{1/2}D^{1/2}$
- $A = L^*(L^*)^T$, where $L^* = LD^{1/2}$

Theorem

If A is a real, symmetric, and positive definite matrix, then $A = LL^T$ for a unique upper triangular L .

Proof.

Existence:

- A positive definite $\Rightarrow A$ nonsingular
- Leading principle minors of A positive definite
 \Rightarrow principle minors nonsingular $\Rightarrow A = LU$ (LU factorization exists)
- $LU = A = A^T = U^T L^T \Leftrightarrow U(L^T)^{-1} = L^{-1}U^T$
- $U(L^T)^{-1}$ upper triangular, $L^{-1}U^T$ lower triangular
 \Rightarrow both sides diagonal (D)
- $U = DL^T$
- D has positive diagonal entries $\Rightarrow D = D^{1/2}D^{1/2}$
- $A = L^*(L^*)^T$, where $L^* = LD^{1/2}$

Unicity: straightforward (assume two, show same)




```
1: function CHOLESKY( $n, (a_{ij})$ )
2:   for  $k = 1$  to  $n$  do
3:      $l_{kk} \leftarrow \left( a_{kk} - \sum_{s=1}^{k-1} l_{ks}^2 \right)^{1/2}$ 
4:     for  $i = k + 1$  to  $n$  do
5:        $l_{ik} \leftarrow \left( a_{ik} - \sum_{s=1}^{k-1} l_{is} l_{ks} \right) / l_{kk}$ 
6:   return  $(l_{ij})$ 
```

where

- $A = (a_{ij})$: factored matrix
- n : order of A
- $L = (l_{ij})$: resulting lower triangular matrix

$$A = \begin{bmatrix} 25 & 15 & -5 \\ 15 & 18 & 0 \\ -5 & 0 & 11 \end{bmatrix}$$

$$L = \begin{bmatrix} & 0 & 0 \\ & & 0 \\ & & \end{bmatrix}$$

```
1: function CHOLESKY( $n, (a_{ij})$ )
2:   for  $k = 1$  to  $n$  do
3:      $l_{kk} \leftarrow \left( a_{kk} - \sum_{s=1}^{k-1} l_{ks}^2 \right)^{1/2}$ 
4:     for  $i = k + 1$  to  $n$  do
5:        $l_{ik} \leftarrow \left( a_{ik} - \sum_{s=1}^{k-1} l_{is} l_{ks} \right) / l_{kk}$ 
6:   return ( $l_{ij}$ )
```

$$A = \begin{bmatrix} 25 & 15 & -5 \\ 15 & 18 & 0 \\ -5 & 0 & 11 \end{bmatrix}$$

$$L = \begin{bmatrix} 5 & 0 & 0 \\ & & 0 \\ & & \end{bmatrix}$$

```

1: function CHOLESKY( $n, (a_{ij})$ )
2:   for  $k = 1$  to  $n$  do
3:      $l_{kk} \leftarrow \left( a_{kk} - \sum_{s=1}^{k-1} l_{ks}^2 \right)^{1/2}$ 
4:     for  $i = k + 1$  to  $n$  do
5:        $l_{ik} \leftarrow \left( a_{ik} - \sum_{s=1}^{k-1} l_{is} l_{ks} \right) / l_{kk}$ 
6:   return ( $l_{ij}$ )
    
```

$k = 1 :$

$$l_{11} = (25 - 0)^{1/2} = 5$$

$$A = \begin{bmatrix} 25 & 15 & -5 \\ 15 & 18 & 0 \\ -5 & 0 & 11 \end{bmatrix}$$

$$L = \begin{bmatrix} 5 & 0 & 0 \\ 3 & & \\ & & \end{bmatrix}$$

```
1: function CHOLESKY( $n, (a_{ij})$ )
2:   for  $k = 1$  to  $n$  do
3:      $l_{kk} \leftarrow \left( a_{kk} - \sum_{s=1}^{k-1} l_{ks}^2 \right)^{1/2}$ 
4:     for  $i = k + 1$  to  $n$  do
5:        $l_{ik} \leftarrow \left( a_{ik} - \sum_{s=1}^{k-1} l_{is} l_{ks} \right) / l_{kk}$ 
6:   return ( $l_{ij}$ )
```

$$k = 1, \quad i = 2 :$$

$$l_{21} = (15 - 0)/5 = 3$$

$$A = \begin{bmatrix} 25 & 15 & -5 \\ 15 & 18 & 0 \\ -5 & 0 & 11 \end{bmatrix}$$

$$L = \begin{bmatrix} 5 & 0 & 0 \\ 3 & & 0 \\ -1 & & \end{bmatrix}$$

```

1: function CHOLESKY( $n, (a_{ij})$ )
2:   for  $k = 1$  to  $n$  do
3:      $l_{kk} \leftarrow \left( a_{kk} - \sum_{s=1}^{k-1} l_{ks}^2 \right)^{1/2}$ 
4:     for  $i = k + 1$  to  $n$  do
5:        $l_{ik} \leftarrow \left( a_{ik} - \sum_{s=1}^{k-1} l_{is} l_{ks} \right) / l_{kk}$ 
6:   return ( $l_{ij}$ )
    
```

$k = 1, \quad i = 3 :$

$$l_{31} = (-5 - 0)/5 = -1$$

$$A = \begin{bmatrix} 25 & 15 & -5 \\ 15 & 18 & 0 \\ -5 & 0 & 11 \end{bmatrix}$$

$$L = \begin{bmatrix} 5 & 0 & 0 \\ 3 & 3 & 0 \\ -1 & & \end{bmatrix}$$

```

1: function CHOLESKY( $n, (a_{ij})$ )
2:   for  $k = 1$  to  $n$  do
3:      $l_{kk} \leftarrow \left( a_{kk} - \sum_{s=1}^{k-1} l_{ks}^2 \right)^{1/2}$ 
4:     for  $i = k + 1$  to  $n$  do
5:        $l_{ik} \leftarrow \left( a_{ik} - \sum_{s=1}^{k-1} l_{is} l_{ks} \right) / l_{kk}$ 
6:   return  $(l_{ij})$ 

```

$k = 2 :$

$$l_{22} = (18 - 3^2)^{1/2} = 3$$

$$A = \begin{bmatrix} 25 & 15 & -5 \\ 15 & 18 & 0 \\ -5 & 0 & 11 \end{bmatrix}$$

$$L = \begin{bmatrix} 5 & 0 & 0 \\ 3 & 3 & 0 \\ -1 & 1 & \end{bmatrix}$$

```

1: function CHOLESKY( $n, (a_{ij})$ )
2:   for  $k = 1$  to  $n$  do
3:      $l_{kk} \leftarrow \left( a_{kk} - \sum_{s=1}^{k-1} l_{ks}^2 \right)^{1/2}$ 
4:     for  $i = k + 1$  to  $n$  do
5:        $l_{ik} \leftarrow \left( a_{ik} - \sum_{s=1}^{k-1} l_{is} l_{ks} \right) / l_{kk}$ 
6:   return ( $l_{ij}$ )

```

$k = 2, \quad i = 3 :$

$$l_{32} = (0 - (-1)(3)) / 3 = 1$$

$$A = \begin{bmatrix} 25 & 15 & -5 \\ 15 & 18 & 0 \\ -5 & 0 & 11 \end{bmatrix}$$

$$L = \begin{bmatrix} 5 & 0 & 0 \\ 3 & 3 & 0 \\ -1 & 1 & 3 \end{bmatrix}$$

```

1: function CHOLESKY( $n, (a_{ij})$ )
2:   for  $k = 1$  to  $n$  do
3:      $l_{kk} \leftarrow \left( a_{kk} - \sum_{s=1}^{k-1} l_{ks}^2 \right)^{1/2}$ 
4:     for  $i = k + 1$  to  $n$  do
5:        $l_{ik} \leftarrow \left( a_{ik} - \sum_{s=1}^{k-1} l_{is} l_{ks} \right) / l_{kk}$ 
6:   return  $(l_{ij})$ 

```

$k = 3 :$

$$l_{33} = (11 - (-1)^2 - (-1)^2) = 3$$

- Verify correctness:

$$L = \begin{bmatrix} 5 & 0 & 0 \\ 3 & 3 & 0 \\ -1 & 1 & 3 \end{bmatrix}$$

- Verify correctness:

$$L = \begin{bmatrix} 5 & 0 & 0 \\ 3 & 3 & 0 \\ -1 & 1 & 3 \end{bmatrix} \implies L^T = \begin{bmatrix} 5 & 3 & -1 \\ 0 & 3 & 1 \\ 0 & 0 & 3 \end{bmatrix}$$

- Verify correctness:

$$L = \begin{bmatrix} 5 & 0 & 0 \\ 3 & 3 & 0 \\ -1 & 1 & 3 \end{bmatrix} \implies L^T = \begin{bmatrix} 5 & 3 & -1 \\ 0 & 3 & 1 \\ 0 & 0 & 3 \end{bmatrix}$$

$$LL^T = \begin{bmatrix} 5 & 0 & 0 \\ 3 & 3 & 0 \\ -1 & 1 & 3 \end{bmatrix} \begin{bmatrix} 5 & 3 & -1 \\ 0 & 3 & 1 \\ 0 & 0 & 3 \end{bmatrix} = \begin{bmatrix} 25 & 15 & -5 \\ 15 & 18 & 0 \\ -5 & 0 & 11 \end{bmatrix} = A$$

Gaussian Elimination

Gaussian Elimination

- Process which attempts to solve a linear system by reducing the coefficient matrix to triangular form using elementary row operations (the result can then be easily solved using forward/backward substitution)

Gaussian Elimination

- Process which attempts to solve a linear system by reducing the coefficient matrix to triangular form using elementary row operations (the result can then be easily solved using forward/backward substitution)
- Possible row operations:
 - Interchange
 - Scaling
 - Addition

Gaussian Elimination

- Process which attempts to solve a linear system by reducing the coefficient matrix to triangular form using elementary row operations (the result can then be easily solved using forward/backward substitution)
- Possible row operations:
 - Interchange
 - Scaling
 - Addition
- Gaussian elimination only utilizes scaling and addition!

```
1: function GE( $n, (a_{ij})$ )
2:   for  $k = 1$  to  $n - 1$  do
3:     for  $i = k + 1$  to  $n$  do
4:        $z \leftarrow a_{ik} / a_{kk}$ 
5:        $a_{ik} \leftarrow 0$ 
6:       for  $j = k + 1$  to  $n$  do
7:          $a_{ij} \leftarrow a_{ij} - za_{kj}$ 
8:   return  $(a_{ij})$ 
```


- Recall: row operations are equivalent to left multiplication of coefficient matrix by elementary matrices

- Recall: row operations are equivalent to left multiplication of coefficient matrix by elementary matrices
 - Example: scale row 2 of $A \in \mathbb{R}^{3 \times 3}$ by 4

$$EA, \quad E = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

- Recall: row operations are equivalent to left multiplication of coefficient matrix by elementary matrices
 - Example: scale row 2 of $A \in \mathbb{R}^{3 \times 3}$ by 4

$$EA, \quad E = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

- Define G.E. recursively in terms of multiplication by these matrices:

$$A_{k+1} = E_k A_k$$

- Recall: row operations are equivalent to left multiplication of coefficient matrix by elementary matrices
 - Example: scale row 2 of $A \in \mathbb{R}^{3 \times 3}$ by 4

$$EA, \quad E = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

- Define G.E. recursively in terms of multiplication by these matrices:

$$A_{k+1} = E_k A_k$$

- Note that algorithm terminates after $s = T_n$ operations with $A_{n \times n}$

- Recall: row operations are equivalent to left multiplication of coefficient matrix by elementary matrices
 - Example: scale row 2 of $A \in \mathbb{R}^{3 \times 3}$ by 4

$$EA, \quad E = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

- Define G.E. recursively in terms of multiplication by these matrices:

$$A_{k+1} = E_k A_k$$

- Note that algorithm terminates after $s = T_n$ operations with $A_{n \times n}$
- Define $U = A_{s+1}$ and observe that

$$U = E_s \cdots E_1 A \quad \text{or} \quad A = E_1^{-1} \cdots E_s^{-1} U = LU$$

- Recall: row operations are equivalent to left multiplication of coefficient matrix by elementary matrices
 - Example: scale row 2 of $A \in \mathbb{R}^{3 \times 3}$ by 4

$$EA, \quad E = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

- Define G.E. recursively in terms of multiplication by these matrices:

$$A_{k+1} = E_k A_k$$

- Note that algorithm terminates after $s = T_n$ operations with $A_{n \times n}$
- Define $U = A_{s+1}$ and observe that

$$U = E_s \cdots E_1 A \quad \text{or} \quad A = E_1^{-1} \cdots E_s^{-1} U = LU$$

- So, Gaussian elimination constructs an LU factorization (and hence is functionally equivalent to the LU factorization algorithm!)

Pivoting Methods

Pivoting Methods

- Problem: Gaussian elimination breaks down (spectacular examples pg. 167–169)

Pivoting Methods

- Problem: Gaussian elimination breaks down (spectacular examples pg. 167–169)
- Solution: incorporate row interchange operations via permutation matrix (we call this pivoting)

Pivoting Methods

- Problem: Gaussian elimination breaks down (spectacular examples pg. 167–169)
- Solution: incorporate row interchange operations via permutation matrix (we call this pivoting)
- Types of pivoting:
 - Partial/complete: pertains to the amount of the matrix examined to find where to pivot

Pivoting Methods

- Problem: Gaussian elimination breaks down (spectacular examples pg. 167–169)
- Solution: incorporate row interchange operations via permutation matrix (we call this pivoting)
- Types of pivoting:
 - Partial/complete: pertains to the amount of the matrix examined to find where to pivot
 - Partial: analyze one row
 - Complete: analyze entire submatrix

Pivoting Methods

- Problem: Gaussian elimination breaks down (spectacular examples pg. 167–169)
- Solution: incorporate row interchange operations via permutation matrix (we call this pivoting)
- Types of pivoting:
 - Partial/complete: pertains to the amount of the matrix examined to find where to pivot
 - Partial: analyze one row
 - Complete: analyze entire submatrix
 - Unscaled/scaled: pertains to how the potential pivot locations are compared

Pivoting Methods

- Problem: Gaussian elimination breaks down (spectacular examples pg. 167–169)
- Solution: incorporate row interchange operations via permutation matrix (we call this pivoting)
- Types of pivoting:
 - Partial/complete: pertains to the amount of the matrix examined to find where to pivot
 - Partial: analyze one row
 - Complete: analyze entire submatrix
 - Unscaled/scaled: pertains to how the potential pivot locations are compared
 - Unscaled: compare absolute value of entries directly
 - Scaled: compare ratios of pivot entries to maximum entries in magnitude in each row

```
1: function GE_PP( $n, (a_{ij}), (p_i)$ )
2:   for  $k = 1$  to  $n - 1$  do
3:     for  $i = k + 1$  to  $n$  do
4:        $z \leftarrow a_{p_i k} / a_{p_k k}$ 
5:        $a_{p_i k} \leftarrow 0$ 
6:       for  $j = k + 1$  to  $n$  do
7:          $a_{p_i j} \leftarrow a_{p_i j} - z a_{p_k j}$ 
8:   return  $(a_{ij})$ 
```

$$A_1 = \begin{bmatrix} 1 & 3 & 6 \\ 2 & 1 & 1 \\ 1 & 3 & 3 \end{bmatrix}$$

$$p = [1, \quad 2, \quad 3]$$

```

1: function GE_PP( $n, (a_{ij}), (p_i)$ )
2:   for  $k = 1$  to  $n - 1$  do
3:     for  $i = k + 1$  to  $n$  do
4:        $z \leftarrow a_{p_i k} / a_{p_k k}$ 
5:        $a_{p_i k} \leftarrow 0$ 
6:       for  $j = k + 1$  to  $n$  do
7:          $a_{p_i j} \leftarrow a_{p_i j} - z a_{p_k j}$ 
8:   return ( $a_{ij}$ )
    
```

$$\max(|1|, |2|, |1|) = 2$$

$$A_1 = \begin{bmatrix} 1 & 3 & 6 \\ 2 & 1 & 1 \\ 1 & 3 & 3 \end{bmatrix}$$

$$p = [2, \quad 1, \quad 3]$$

```

1: function GE_PP( $n, (a_{ij}), (p_i)$ )
2:   for  $k = 1$  to  $n - 1$  do
3:     for  $i = k + 1$  to  $n$  do
4:        $z \leftarrow a_{p_i k} / a_{p_k k}$ 
5:        $a_{p_i k} \leftarrow 0$ 
6:       for  $j = k + 1$  to  $n$  do
7:          $a_{p_i j} \leftarrow a_{p_i j} - z a_{p_k j}$ 
8:   return ( $a_{ij}$ )
    
```

$$\max(|1|, |2|, |1|) = 2$$

$$R_1 \leftarrow R_1 - \frac{1}{2} R_2$$

$$R_3 \leftarrow R_3 - \frac{1}{2} R_2$$

$$R_2 \leftarrow \frac{1}{2} R_2$$

$$A_2 = \begin{bmatrix} 0 & 2.5 & 5.5 \\ 1 & 0.5 & 0.5 \\ 0 & 2.5 & 2.5 \end{bmatrix}$$

$$p = [2, \quad 1, \quad 3]$$

```

1: function GE_PP( $n, (a_{ij}), (p_i)$ )
2:   for  $k = 1$  to  $n - 1$  do
3:     for  $i = k + 1$  to  $n$  do
4:        $z \leftarrow a_{p_i k} / a_{p_k k}$ 
5:        $a_{p_i k} \leftarrow 0$ 
6:       for  $j = k + 1$  to  $n$  do
7:          $a_{p_i j} \leftarrow a_{p_i j} - z a_{p_k j}$ 
8:   return ( $a_{ij}$ )
    
```

$$\max(|1|, |2|, |1|) = 2$$

$$R_1 \leftarrow R_1 - \frac{1}{2} R_2$$

$$R_3 \leftarrow R_3 - \frac{1}{2} R_2$$

$$R_2 \leftarrow \frac{1}{2} R_2$$

$$A_2 = \begin{bmatrix} 0 & 2.5 & 5.5 \\ 1 & 0.5 & 0.5 \\ 0 & 2.5 & 2.5 \end{bmatrix}$$

$$p = [2, \quad 1, \quad 3]$$

```

1: function GE_PP( $n, (a_{ij}), (p_i)$ )
2:   for  $k = 1$  to  $n - 1$  do
3:     for  $i = k + 1$  to  $n$  do
4:        $z \leftarrow a_{p_i k} / a_{p_k k}$ 
5:        $a_{p_i k} \leftarrow 0$ 
6:       for  $j = k + 1$  to  $n$  do
7:          $a_{p_i j} \leftarrow a_{p_i j} - z a_{p_k j}$ 
8:   return ( $a_{ij}$ )

```

$$\max(|2.5|, |2.5|) = 2.5$$

$$A_2 = \begin{bmatrix} 0 & 2.5 & 5.5 \\ 1 & 0.5 & 0.5 \\ 0 & 2.5 & 2.5 \end{bmatrix}$$

$$p = [2, \quad 1, \quad 3]$$

```

1: function GE_PP( $n, (a_{ij}), (p_i)$ )
2:   for  $k = 1$  to  $n - 1$  do
3:     for  $i = k + 1$  to  $n$  do
4:        $z \leftarrow a_{p_i k} / a_{p_k k}$ 
5:        $a_{p_i k} \leftarrow 0$ 
6:       for  $j = k + 1$  to  $n$  do
7:          $a_{p_i j} \leftarrow a_{p_i j} - z a_{p_k j}$ 
8:   return ( $a_{ij}$ )
    
```

$$\max(|2.5|, |2.5|) = 2.5$$

$$R_3 \leftarrow R_3 - R_2$$

$$R_1 \leftarrow \frac{2}{5} R_1$$

$$A_3 = \begin{bmatrix} 0 & 1 & 2.2 \\ 1 & 0.5 & -0.6 \\ 0 & 0 & -3 \end{bmatrix}$$

$$p = [2, \quad 1, \quad 3]$$

```

1: function GE_PP( $n, (a_{ij}), (p_i)$ )
2:   for  $k = 1$  to  $n - 1$  do
3:     for  $i = k + 1$  to  $n$  do
4:        $z \leftarrow a_{p_i k} / a_{p_k k}$ 
5:        $a_{p_i k} \leftarrow 0$ 
6:       for  $j = k + 1$  to  $n$  do
7:          $a_{p_i j} \leftarrow a_{p_i j} - z a_{p_k j}$ 
8:   return ( $a_{ij}$ )

```

$$\max(|2.5|, |2.5|) = 2.5$$

$$R_3 \leftarrow R_3 - R_2$$

$$R_1 \leftarrow \frac{2}{5} R_1$$

$$A_3 = \begin{bmatrix} 0 & 1 & 2.2 \\ 1 & 0.5 & -0.6 \\ 0 & 0 & -3 \end{bmatrix}$$

$$p = [2, \quad 1, \quad 3]$$

```

1: function GE_PP( $n, (a_{ij}), (p_i)$ )
2:   for  $k = 1$  to  $n - 1$  do
3:     for  $i = k + 1$  to  $n$  do
4:        $z \leftarrow a_{p_i k} / a_{p_k k}$ 
5:        $a_{p_i k} \leftarrow 0$ 
6:       for  $j = k + 1$  to  $n$  do
7:          $a_{p_i j} \leftarrow a_{p_i j} - z a_{p_k j}$ 
8:   return ( $a_{ij}$ )

```

$$R_3 \leftarrow -\frac{1}{3}R_3$$

$$A_4 = \begin{bmatrix} 0 & 1 & 2.2 \\ 1 & 0.5 & -0.6 \\ 0 & 0 & 1 \end{bmatrix}$$

$$p = [2, \quad 1, \quad 3]$$

```

1: function GE_PP( $n, (a_{ij}), (p_i)$ )
2:   for  $k = 1$  to  $n - 1$  do
3:     for  $i = k + 1$  to  $n$  do
4:        $z \leftarrow a_{p_i k} / a_{p_k k}$ 
5:        $a_{p_i k} \leftarrow 0$ 
6:       for  $j = k + 1$  to  $n$  do
7:          $a_{p_i j} \leftarrow a_{p_i j} - z a_{p_k j}$ 
8:   return ( $a_{ij}$ )

```

$$R_3 \leftarrow -\frac{1}{3}R_3$$