

# A Summary of Methods of Solution for Linear Systems

Nate DeMaagd, Kurt O'Hearn

MTH 499-02

April 23, 2013



- Types of methods

- Types of methods
- Direct methods and examples

- Types of methods
- Direct methods and examples
- Indirect methods and examples



Recall: For the linear system  $Ax = b$  with  $A_{m \times n}$ :

Recall: For the linear system  $Ax = b$  with  $A_{m \times n}$ :

System Type	Possible Number of Solutions
Square ( $m = n$ )	None, Unique
Overdetermined ( $m > n$ )	None, Unique
Underdetermined ( $m < n$ )	None, Infinite



Recall: For the linear system  $Ax = b$  with  $A_{m \times n}$ :

System Type	Possible Number of Solutions
Square ( $m = n$ )	None, Unique
Overdetermined ( $m > n$ )	None, Unique
Underdetermined ( $m < n$ )	None, Infinite

Types of Methods

- Direct methods

Recall: For the linear system  $Ax = b$  with  $A_{m \times n}$ :

System Type	Possible Number of Solutions
Square ( $m = n$ )	None, Unique
Overdetermined ( $m > n$ )	None, Unique
Underdetermined ( $m < n$ )	None, Infinite

## Types of Methods

- Direct methods
  - Execute a predetermined number of computations to produce a result

Recall: For the linear system  $Ax = b$  with  $A_{m \times n}$ :

System Type	Possible Number of Solutions
Square ( $m = n$ )	None, Unique
Overdetermined ( $m > n$ )	None, Unique
Underdetermined ( $m < n$ )	None, Infinite

## Types of Methods

- Direct methods
  - Execute a predetermined number of computations to produce a result
  - Methods: compute  $A^{-1}$ , transform  $A$  using factorizations/pivoting

Recall: For the linear system  $Ax = b$  with  $A_{m \times n}$ :

System Type	Possible Number of Solutions
Square ( $m = n$ )	None, Unique
Overdetermined ( $m > n$ )	None, Unique
Underdetermined ( $m < n$ )	None, Infinite

## Types of Methods

- Direct methods
  - Execute a predetermined number of computations to produce a result
  - Methods: compute  $A^{-1}$ , transform  $A$  using factorizations/pivoting
- Indirect methods

Recall: For the linear system  $Ax = b$  with  $A_{m \times n}$ :

System Type	Possible Number of Solutions
Square ( $m = n$ )	None, Unique
Overdetermined ( $m > n$ )	None, Unique
Underdetermined ( $m < n$ )	None, Infinite

## Types of Methods

- Direct methods
  - Execute a predetermined number of computations to produce a result
  - Methods: compute  $A^{-1}$ , transform  $A$  using factorizations/pivoting
- Indirect methods
  - Generate a sequence of intermediate results which (hopefully) produce the desired final result

Recall: For the linear system  $Ax = b$  with  $A_{m \times n}$ :

System Type	Possible Number of Solutions
Square ( $m = n$ )	None, Unique
Overdetermined ( $m > n$ )	None, Unique
Underdetermined ( $m < n$ )	None, Infinite

## Types of Methods

- Direct methods
  - Execute a predetermined number of computations to produce a result
  - Methods: compute  $A^{-1}$ , transform  $A$  using factorizations/pivoting
- Indirect methods
  - Generate a sequence of intermediate results which (hopefully) produce the desired final result
  - Methods:

Recall: For the linear system  $Ax = b$  with  $A_{m \times n}$ :

System Type	Possible Number of Solutions
Square ( $m = n$ )	None, Unique
Overdetermined ( $m > n$ )	None, Unique
Underdetermined ( $m < n$ )	None, Infinite

## Types of Methods

- Direct methods
  - Execute a predetermined number of computations to produce a result
  - Methods: compute  $A^{-1}$ , transform  $A$  using factorizations/pivoting
- Indirect methods
  - Generate a sequence of intermediate results which (hopefully) produce the desired final result
  - Methods:
    - Richardson, Jacobi, Gauss-Seidel, SOR, steepest descent, conjugate gradient

Recall: For the linear system  $Ax = b$  with  $A_{m \times n}$ :

System Type	Possible Number of Solutions
Square ( $m = n$ )	None, Unique
Overdetermined ( $m > n$ )	None, Unique
Underdetermined ( $m < n$ )	None, Infinite

## Types of Methods

- Direct methods
  - Execute a predetermined number of computations to produce a result
  - Methods: compute  $A^{-1}$ , transform  $A$  using factorizations/pivoting
- Indirect methods
  - Generate a sequence of intermediate results which (hopefully) produce the desired final result
  - Methods:
    - Richardson, Jacobi, Gauss-Seidel, SOR, steepest descent, conjugate gradient

Note: methods can be general or exploit certain matrix characteristics





Compute  $A^{-1}$

Compute  $A^{-1}$

- Most likely too difficult but not always (e.g., diagonal)

Compute  $A^{-1}$

- Most likely too difficult but not always (e.g., diagonal)

Gaussian Elimination

Compute  $A^{-1}$

- Most likely too difficult but not always (e.g., diagonal)

Gaussian Elimination

- Solve by reducing the coefficient matrix to triangular form using elementary row operations

Compute  $A^{-1}$

- Most likely too difficult but not always (e.g., diagonal)

Gaussian Elimination

- Solve by reducing the coefficient matrix to triangular form using elementary row operations
- Result can then be easily solved using forward/backward substitution

Compute  $A^{-1}$

- Most likely too difficult but not always (e.g., diagonal)

Gaussian Elimination

- Solve by reducing the coefficient matrix to triangular form using elementary row operations
- Result can then be easily solved using forward/backward substitution
- Possible row operations:
  - Interchange
  - Scaling
  - Addition

Compute  $A^{-1}$

- Most likely too difficult but not always (e.g., diagonal)

Gaussian Elimination

- Solve by reducing the coefficient matrix to triangular form using elementary row operations
- Result can then be easily solved using forward/backward substitution
- Possible row operations:
  - Interchange
  - Scaling
  - Addition
- Recall: GE is equivalent to computing  $LU$  factorization



## Pivoting Methods

## Pivoting Methods

- Types of pivoting:
  - Partial/complete: pertains to the amount of the matrix examined to find where to pivot

## Pivoting Methods

- Types of pivoting:
  - Partial/complete: pertains to the amount of the matrix examined to find where to pivot
    - Partial: analyze one row
    - Complete: analyze entire submatrix

## Pivoting Methods

- Types of pivoting:
  - Partial/complete: pertains to the amount of the matrix examined to find where to pivot
    - Partial: analyze one row
    - Complete: analyze entire submatrix
  - Unscaled/scaled: pertains to how the potential pivot locations are compared

## Pivoting Methods

- Types of pivoting:
  - Partial/complete: pertains to the amount of the matrix examined to find where to pivot
    - Partial: analyze one row
    - Complete: analyze entire submatrix
  - Unscaled/scaled: pertains to how the potential pivot locations are compared
    - Unscaled: compare absolute value of entries directly
    - Scaled: compare ratios of pivot entries to maximum entries in magnitude in each row

$$A_1 = \begin{bmatrix} 1 & 3 & 6 \\ 2 & 1 & 1 \\ 1 & 3 & 3 \end{bmatrix}$$

$$p = [1, \quad 2, \quad 3]$$

```

1: function GE_SPP( $n, (a_{ij}), (p_i)$ )
2:   for  $k = 1$  to  $n - 1$  do
3:     for  $i = k + 1$  to  $n$  do
4:        $z \leftarrow a_{p_i k} / a_{p_k k}$ 
5:        $a_{p_i k} \leftarrow 0$ 
6:       for  $j = k + 1$  to  $n$  do
7:          $a_{p_i j} \leftarrow a_{p_i j} - z a_{p_k j}$ 
8:   return  $(a_{ij})$ 

```

$$\max(|1|, |2|, |1|) = 2$$

$$A_1 = \begin{bmatrix} 1 & 3 & 6 \\ 2 & 1 & 1 \\ 1 & 3 & 3 \end{bmatrix}$$

$$p = [2, \quad 1, \quad 3]$$

```

1: function GE_SPP( $n, (a_{ij}), (p_i)$ )
2:   for  $k = 1$  to  $n - 1$  do
3:     for  $i = k + 1$  to  $n$  do
4:        $z \leftarrow a_{p_i k} / a_{p_k k}$ 
5:        $a_{p_i k} \leftarrow 0$ 
6:       for  $j = k + 1$  to  $n$  do
7:          $a_{p_i j} \leftarrow a_{p_i j} - z a_{p_k j}$ 
8:   return ( $a_{ij}$ )

```

$$\max(|1|, |2|, |1|) = 2$$

$$R_1 \leftarrow R_1 - \frac{1}{2}R_2$$

$$R_3 \leftarrow R_3 - \frac{1}{2}R_2$$

$$R_2 \leftarrow \frac{1}{2}R_2$$

$$A_2 = \begin{bmatrix} 0 & 2.5 & 5.5 \\ 1 & 0.5 & 0.5 \\ 0 & 2.5 & 2.5 \end{bmatrix}$$

$$p = [2, \quad 1, \quad 3]$$

```

1: function GE_SPP( $n, (a_{ij}), (p_i)$ )
2:   for  $k = 1$  to  $n - 1$  do
3:     for  $i = k + 1$  to  $n$  do
4:        $z \leftarrow a_{p_i k} / a_{p_k k}$ 
5:        $a_{p_i k} \leftarrow 0$ 
6:       for  $j = k + 1$  to  $n$  do
7:          $a_{p_i j} \leftarrow a_{p_i j} - z a_{p_k j}$ 
8:   return ( $a_{ij}$ )

```

$$\max(|1|, |2|, |1|) = 2$$

$$R_1 \leftarrow R_1 - \frac{1}{2} R_2$$

$$R_3 \leftarrow R_3 - \frac{1}{2} R_2$$

$$R_2 \leftarrow \frac{1}{2} R_2$$



$$A_2 = \begin{bmatrix} 0 & 2.5 & 5.5 \\ 1 & 0.5 & 0.5 \\ 0 & 2.5 & 2.5 \end{bmatrix}$$

$$p = [2, \quad 1, \quad 3]$$

```

1: function GE_SPP( $n, (a_{ij}), (p_i)$ )
2:   for  $k = 1$  to  $n - 1$  do
3:     for  $i = k + 1$  to  $n$  do
4:        $z \leftarrow a_{p_i k} / a_{p_k k}$ 
5:        $a_{p_i k} \leftarrow 0$ 
6:       for  $j = k + 1$  to  $n$  do
7:          $a_{p_i j} \leftarrow a_{p_i j} - z a_{p_k j}$ 
8:   return  $(a_{ij})$ 

```

$$\max(|2.5|, |2.5|) = 2.5$$

$$A_2 = \begin{bmatrix} 0 & 2.5 & 5.5 \\ 1 & 0.5 & 0.5 \\ 0 & 2.5 & 2.5 \end{bmatrix}$$

$$p = [2, \quad 1, \quad 3]$$

```

1: function GE_SPP( $n, (a_{ij}), (p_i)$ )
2:   for  $k = 1$  to  $n - 1$  do
3:     for  $i = k + 1$  to  $n$  do
4:        $z \leftarrow a_{p_i k} / a_{p_k k}$ 
5:        $a_{p_i k} \leftarrow 0$ 
6:       for  $j = k + 1$  to  $n$  do
7:          $a_{p_i j} \leftarrow a_{p_i j} - z a_{p_k j}$ 
8:   return  $(a_{ij})$ 

```

$$\max(|2.5|, |2.5|) = 2.5$$

$$R_3 \leftarrow R_3 - R_2$$

$$R_1 \leftarrow \frac{2}{5} R_1$$

$$A_3 = \begin{bmatrix} 0 & 1 & 2.2 \\ 1 & 0.5 & -0.6 \\ 0 & 0 & -3 \end{bmatrix}$$

$$p = [2, \quad 1, \quad 3]$$

```

1: function GE_SPP( $n, (a_{ij}), (p_i)$ )
2:   for  $k = 1$  to  $n - 1$  do
3:     for  $i = k + 1$  to  $n$  do
4:        $z \leftarrow a_{p_i k} / a_{p_k k}$ 
5:        $a_{p_i k} \leftarrow 0$ 
6:       for  $j = k + 1$  to  $n$  do
7:          $a_{p_i j} \leftarrow a_{p_i j} - z a_{p_k j}$ 
8:   return ( $a_{ij}$ )

```

$$\max(|2.5|, |2.5|) = 2.5$$

$$R_3 \leftarrow R_3 - R_2$$

$$R_1 \leftarrow \frac{2}{5} R_1$$

$$A_3 = \begin{bmatrix} 0 & 1 & 2.2 \\ 1 & 0.5 & -0.6 \\ 0 & 0 & -3 \end{bmatrix}$$

$$p = [2, \quad 1, \quad 3]$$

```

1: function GE_SPP( $n, (a_{ij}), (p_i)$ )
2:   for  $k = 1$  to  $n - 1$  do
3:     for  $i = k + 1$  to  $n$  do
4:        $z \leftarrow a_{p_i k} / a_{p_k k}$ 
5:        $a_{p_i k} \leftarrow 0$ 
6:       for  $j = k + 1$  to  $n$  do
7:          $a_{p_i j} \leftarrow a_{p_i j} - z a_{p_k j}$ 
8:   return ( $a_{ij}$ )
    
```

$$R_3 \leftarrow -\frac{1}{3}R_3$$

$$A_4 = \begin{bmatrix} 0 & 1 & 2.2 \\ 1 & 0.5 & -0.6 \\ 0 & 0 & 1 \end{bmatrix}$$

$$p = [2, \quad 1, \quad 3]$$

```

1: function GE_SPP( $n, (a_{ij}), (p_i)$ )
2:   for  $k = 1$  to  $n - 1$  do
3:     for  $i = k + 1$  to  $n$  do
4:        $z \leftarrow a_{p_i k} / a_{p_k k}$ 
5:        $a_{p_i k} \leftarrow 0$ 
6:       for  $j = k + 1$  to  $n$  do
7:          $a_{p_i j} \leftarrow a_{p_i j} - z a_{p_k j}$ 
8:   return ( $a_{ij}$ )

```

$$R_3 \leftarrow -\frac{1}{3}R_3$$



Recall: General Conditions for Iterative Method Convergence

## Recall: General Conditions for Iterative Method Convergence

## Theorem

*For the linear system  $Ax = b$  with  $A$  invertible, define the iteration formula*

$$x^{(k)} = Gx^{(k-1)} + c.$$

*The sequence  $\{x^{(k)}\}$  will converge to  $(I - G)^{-1}c$  provided that  $\rho(G) < 1$ .*



## Recall: General Conditions for Iterative Method Convergence

## Theorem

*For the linear system  $Ax = b$  with  $A$  invertible, define the iteration formula*

$$x^{(k)} = Gx^{(k-1)} + c.$$

*The sequence  $\{x^{(k)}\}$  will converge to  $(I - G)^{-1}c$  provided that  $\rho(G) < 1$ .*

## Methods

- Richardson
- Jacobi
- Gauss-Seidel
- SOR
- Steepest Descent
- Conjugate Gradient

## Method

- Richardson

## System Stipulations

- $A$  invertible

## Iteration Formula

- $x^{(k)} = (I - A)x^{(k-1)} + b = x^{(k-1)} + r^{(k-1)}$

## Convergence Criteria

- $\rho(I - A) < 1$



- Consider matrix  $A = \begin{bmatrix} 6 & 1 & 1 \\ 2 & 4 & 0 \\ 1 & 2 & 6 \end{bmatrix}$  with solution  $b = \begin{bmatrix} 12 \\ 0 \\ 6 \end{bmatrix}$ .

- Consider matrix  $A = \begin{bmatrix} 6 & 1 & 1 \\ 2 & 4 & 0 \\ 1 & 2 & 6 \end{bmatrix}$  with solution  $b = \begin{bmatrix} 12 \\ 0 \\ 6 \end{bmatrix}$ .
- Algorithm is  $x^{(k+1)} = (I - \omega A)x^{(k)} + \omega b^{(k)}$  for some scalar  $\omega \neq 0$  such that  $|r| < 1$ .

- Consider matrix  $A = \begin{bmatrix} 6 & 1 & 1 \\ 2 & 4 & 0 \\ 1 & 2 & 6 \end{bmatrix}$  with solution  $b = \begin{bmatrix} 12 \\ 0 \\ 6 \end{bmatrix}$ .
- Algorithm is  $x^{(k+1)} = (I - \omega A)x^{(k)} + \omega b^{(k)}$  for some scalar  $\omega \neq 0$  such that  $|r| < 1$ .
- Let  $x^{(0)} = [2 \quad 2 \quad 2]^T$  and  $\omega = \frac{1}{6}$ .

- Consider matrix  $A = \begin{bmatrix} 6 & 1 & 1 \\ 2 & 4 & 0 \\ 1 & 2 & 6 \end{bmatrix}$  with solution  $b = \begin{bmatrix} 12 \\ 0 \\ 6 \end{bmatrix}$ .
- Algorithm is  $x^{(k+1)} = (I - \omega A)x^{(k)} + \omega b^{(k)}$  for some scalar  $\omega \neq 0$  such that  $|r| < 1$ .
- Let  $x^{(0)} = [2 \quad 2 \quad 2]^T$  and  $\omega = \frac{1}{6}$ .
- 

$$\begin{aligned} x^{(1)} &= \begin{bmatrix} 0 & -1/6 & -1/6 \\ -1/3 & 1/3 & 0 \\ -1/6 & -1/3 & 0 \end{bmatrix} \begin{bmatrix} 2 \\ 2 \\ 2 \end{bmatrix} + \frac{1}{6} \begin{bmatrix} 12 \\ 0 \\ 6 \end{bmatrix} \\ &= [4/3 \quad 0 \quad 0]^T \end{aligned}$$

- Consider matrix  $A = \begin{bmatrix} 6 & 1 & 1 \\ 2 & 4 & 0 \\ 1 & 2 & 6 \end{bmatrix}$  with solution  $b = \begin{bmatrix} 12 \\ 0 \\ 6 \end{bmatrix}$ .
- Algorithm is  $x^{(k+1)} = (I - \omega A)x^{(k)} + \omega b^{(k)}$  for some scalar  $\omega \neq 0$  such that  $|r| < 1$ .
- Let  $x^{(0)} = [2 \quad 2 \quad 2]^T$  and  $\omega = \frac{1}{6}$ .
- 

$$\begin{aligned} x^{(1)} &= \begin{bmatrix} 0 & -1/6 & -1/6 \\ -1/3 & 1/3 & 0 \\ -1/6 & -1/3 & 0 \end{bmatrix} \begin{bmatrix} 2 \\ 2 \\ 2 \end{bmatrix} + \frac{1}{6} \begin{bmatrix} 12 \\ 0 \\ 6 \end{bmatrix} \\ &= [4/3 \quad 0 \quad 0]^T \end{aligned}$$

- After 12 iterations,  $x^{(12)} \approx [2 \quad -1 \quad 1]^T$ .



## Method

- Jacobi

## System Stipulations

- $A$  invertible
- $A$  diagonally dominant

## Iteration Formula

- $x^{(k)} = (I - D^{-1}A)x^{(k-1)} + D^{-1}b$
- $D = a_{ii}$

## Convergence Criteria

- $\rho(I - D^{-1}A) < 1$

## Method

- Gauss-Seidel

## System Stipulations

- $A$  invertible
- $A$  diagonally dominant

## Iteration Formula

- $x^{(k)} = (I - L^{-1}A)x^{(k-1)} + L^{-1}b$
- $L = a_{ij}, i \geq j$

## Convergence Criteria

- $\rho(I - L^{-1}A) < 1$

## Method

- SOR

## System Stipulations

- Complex system
- $A$ : positive definite, Hermitian

## Iteration Formula

- $x^{(k+1)} = (I - (\alpha D - C)^{-1}A)x^{(k)} + (\alpha D - C)^{-1}b$
- $D$ : positive definite, Hermitian
- $C : C + C^* = D - A$
- $\alpha \in \mathbb{R} \ni \alpha > \frac{1}{2}$

## Convergence Criteria

- $\rho(I - (\alpha D - C)^{-1}A) < 1$

## Typical values

- $D = a_{ii}, C = -a_{ij} \ni i \geq j$

## SOR Example

## Method

- Steepest Descent

## System Stipulations

- $A$  invertible
- $A$  symmetric, positive definite

## Iteration Formula

- $x^{(k)} = x^{(k-1)} + t_{k-1}v^{(k-1)}$
- $t_{k-1} = \frac{\langle v^{(k-1)}, b - Ax^{(k-1)} \rangle}{\langle v^{(k-1)}, Av^{(k-1)} \rangle}$
- $v^{(k-1)} = -\nabla x^{(k-1)} = -\frac{d}{dx}q(x^{(k-1)})$
- $q(x) = \langle x, Ax \rangle - 2\langle x, b \rangle$

## Method

- Conjugate Gradient

## System Stipulations

- $A$  invertible
- $A$  symmetric, positive definite

## Iteration Formula

- $x^{(k)} = x^{(k-1)} + t_{k-1}v^{(k-1)}$
- $t_{k-1} = \frac{\langle v^{(k-1)}, b - Ax^{(k-1)} \rangle}{\langle v^{(k-1)}, Av^{(k-1)} \rangle}$
- $\{v^{(0)}, \dots, v^{(n)}\}$  are  $A$ -orthogonal
- $v^{(k-1)} = -\nabla x^{(k-1)} = -\frac{d}{dx}q(x^{(k-1)})$
- $q(x) = \langle x, Ax \rangle - 2\langle x, b \rangle$