

Horner's Method and Finding Roots of Polynomials

Nate DeMaagd, Kurt O'Hearn

MTH 499-02

February 18, 2013

Our outline for today is...

Our outline for today is...

- Development of Horner's Algorithm

Our outline for today is...

- Development of Horner's Algorithm
- Pseudocode

Our outline for today is...

- Development of Horner's Algorithm
- Pseudocode
- Applications and Connections to Root Finding

Our outline for today is...

- Development of Horner's Algorithm
- Pseudocode
- Applications and Connections to Root Finding
- Related Ideas: Julia Sets and Fractals

Horner's Algorithm (Polynomials)

- Consider a polynomial p with degree n and some point x_0 . We form the degree $n - 1$ polynomial q as:

$$q(x) = \frac{p(x) - p(x_0)}{x - x_0}.$$

Horner's Algorithm (Polynomials)

- Consider a polynomial p with degree n and some point x_0 . We form the degree $n - 1$ polynomial q as:

$$q(x) = \frac{p(x) - p(x_0)}{x - x_0}.$$

- Rearranging, we have

$$p(x) = (x - x_0)q(x) + p(x_0). \tag{1}$$

Horner's Algorithm (Polynomials)

- Consider a polynomial p with degree n and some point x_0 . We form the degree $n - 1$ polynomial q as:

$$q(x) = \frac{p(x) - p(x_0)}{x - x_0}.$$

- Rearranging, we have

$$p(x) = (x - x_0)q(x) + p(x_0). \quad (1)$$

- Expanding p and q :

$$p(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-2}x^{n-2} + a_{n-1}x^{n-1} + a_nx^n$$

$$q(x) = b_0 + b_1x + b_2x^2 + \cdots + b_{n-2}x^{n-2} + b_{n-1}x^{n-1}.$$

- Equating like coefficients in (1), we have

- Equating like coefficients in (1), we have

$$\text{constant : } p(x_0) = a_0 + b_0x_0$$

$$x : b_0 = a_1 + b_1x_0$$

$$x^2 : b_1 = a_2 + b_2x_0$$

$$\vdots$$

$$x^{n-1} : b_{n-2} = a_{n-1} + b_{n-1}x_0$$

$$x^n : b_{n-1} = a_n.$$

- Equating like coefficients in (1), we have

$$\text{constant : } p(x_0) = a_0 + b_0x_0$$

$$x : b_0 = a_1 + b_1x_0$$

$$x^2 : b_1 = a_2 + b_2x_0$$

$$\vdots$$

$$x^{n-1} : b_{n-2} = a_{n-1} + b_{n-1}x_0$$

$$x^n : b_{n-1} = a_n.$$

- Notice: repeated substitution starting with b_{n-1} to find all b_j and $p(x_0)$

- Equating like coefficients in (1), we have

$$\text{constant : } p(x_0) = a_0 + b_0 x_0$$

$$x : b_0 = a_1 + b_1 x_0$$

$$x^2 : b_1 = a_2 + b_2 x_0$$

$$\vdots$$

$$x^{n-1} : b_{n-2} = a_{n-1} + b_{n-1} x_0$$

$$x^n : b_{n-1} = a_n.$$

- Notice: repeated substitution starting with b_{n-1} to find all b_j and $p(x_0)$
- Question: why should we care?

```
1: function HORNER( $n, [a_0, \dots, a_n], x_0$ )
2:    $b_{n-1} \leftarrow a_n$ 
3:   for  $k = 0$  to  $n - 1$  step  $-1$  do
4:      $b_{k-1} \leftarrow a_k + b_k x_0$ 
5:   return  $[b_{-1}, \dots, b_{n-1}]$ 
```

where

n : degree of p

a_i : coefficients of i^{th} degree term in p (degree n)

b_j : coefficients of j^{th} degree term in q (degree $n - 1$)

x_0 : evaluation point

Applications

Applications

- Function evaluation and Deflation

Applications

- Function evaluation and Deflation
- Taylor Expansions

Applications

- Function evaluation and Deflation
- Taylor Expansions
- Newton's method

- Recall: Given p (degree n) and some x_0 , Horner's algorithm finds all coefficients b_j of q (degree $n - 1$) and $p(x_0)$

- Recall: Given p (degree n) and some x_0 , Horner's algorithm finds all coefficients b_j of q (degree $n - 1$) and $p(x_0)$
- Example: Deflate

$$p(x) = x^3 - 2x^2 - 5x + 6, \quad x_0 = -1$$

- Recall: Given p (degree n) and some x_0 , Horner's algorithm finds all coefficients b_j of q (degree $n - 1$) and $p(x_0)$
- Example: Deflate

$$p(x) = x^3 - 2x^2 - 5x + 6, \quad x_0 = -1$$

- Use synthetic division table:

- Recall: Given p (degree n) and some x_0 , Horner's algorithm finds all coefficients b_j of q (degree $n - 1$) and $p(x_0)$
- Example: Deflate

$$p(x) = x^3 - 2x^2 - 5x + 6, \quad x_0 = -1$$

- Use synthetic division table:

	a_n	a_{n-1}	a_{n-2}	\cdots	a_1	a_0
x_0		$a_n x_0$	$b_{n-1} x_0$	\cdots	$b_1 x_0$	$b_0 x_0$
	a_n	$a_{n-1} + a_n x_0$	$a_{n-2} + b_{n-1} x_0$	\cdots	$a_1 + b_1 x_0$	$a_0 + b_0 x_0$

- Recall: Given p (degree n) and some x_0 , Horner's algorithm finds all coefficients b_j of q (degree $n - 1$) and $p(x_0)$
- Example: Deflate

$$p(x) = x^3 - 2x^2 - 5x + 6, \quad x_0 = -1$$

- Use synthetic division table:

	a_n	a_{n-1}	a_{n-2}	\cdots	a_1	a_0
x_0		$a_n x_0$	$b_{n-1} x_0$	\cdots	$b_1 x_0$	$b_0 x_0$
	a_n	$a_{n-1} + a_n x_0$	$a_{n-2} + b_{n-1} x_0$	\cdots	$a_1 + b_1 x_0$	$a_0 + b_0 x_0$

	1	-2	-5	6
-1				

- Recall: Given p (degree n) and some x_0 , Horner's algorithm finds all coefficients b_j of q (degree $n - 1$) and $p(x_0)$
- Example: Deflate

$$p(x) = x^3 - 2x^2 - 5x + 6, \quad x_0 = -1$$

- Use synthetic division table:

	a_n	a_{n-1}	a_{n-2}	\cdots	a_1	a_0
x_0		$a_n x_0$	$b_{n-1} x_0$	\cdots	$b_1 x_0$	$b_0 x_0$
	a_n	$a_{n-1} + a_n x_0$	$a_{n-2} + b_{n-1} x_0$	\cdots	$a_1 + b_1 x_0$	$a_0 + b_0 x_0$

	1	-2	-5	6
-1				
	1			

- Recall: Given p (degree n) and some x_0 , Horner's algorithm finds all coefficients b_j of q (degree $n - 1$) and $p(x_0)$
- Example: Deflate

$$p(x) = x^3 - 2x^2 - 5x + 6, \quad x_0 = -1$$

- Use synthetic division table:

	a_n	a_{n-1}	a_{n-2}	\cdots	a_1	a_0
x_0		$a_n x_0$	$b_{n-1} x_0$	\cdots	$b_1 x_0$	$b_0 x_0$
	a_n	$a_{n-1} + a_n x_0$	$a_{n-2} + b_{n-1} x_0$	\cdots	$a_1 + b_1 x_0$	$a_0 + b_0 x_0$

	1	-2	-5	6
-1		-1		
	1			

- Recall: Given p (degree n) and some x_0 , Horner's algorithm finds all coefficients b_j of q (degree $n - 1$) and $p(x_0)$
- Example: Deflate

$$p(x) = x^3 - 2x^2 - 5x + 6, \quad x_0 = -1$$

- Use synthetic division table:

	a_n	a_{n-1}	a_{n-2}	\cdots	a_1	a_0
x_0		$a_n x_0$	$b_{n-1} x_0$	\cdots	$b_1 x_0$	$b_0 x_0$
	a_n	$a_{n-1} + a_n x_0$	$a_{n-2} + b_{n-1} x_0$	\cdots	$a_1 + b_1 x_0$	$a_0 + b_0 x_0$

	1	-2	-5	6
-1		-1		
	1	-3		

- Recall: Given p (degree n) and some x_0 , Horner's algorithm finds all coefficients b_j of q (degree $n - 1$) and $p(x_0)$
- Example: Deflate

$$p(x) = x^3 - 2x^2 - 5x + 6, \quad x_0 = -1$$

- Use synthetic division table:

	a_n	a_{n-1}	a_{n-2}	\cdots	a_1	a_0
x_0		$a_n x_0$	$b_{n-1} x_0$	\cdots	$b_1 x_0$	$b_0 x_0$
	a_n	$a_{n-1} + a_n x_0$	$a_{n-2} + b_{n-1} x_0$	\cdots	$a_1 + b_1 x_0$	$a_0 + b_0 x_0$

	1	-2	-5	6
-1		-1	3	
	1	-3		

- Recall: Given p (degree n) and some x_0 , Horner's algorithm finds all coefficients b_j of q (degree $n - 1$) and $p(x_0)$
- Example: Deflate

$$p(x) = x^3 - 2x^2 - 5x + 6, \quad x_0 = -1$$

- Use synthetic division table:

	a_n	a_{n-1}	a_{n-2}	\cdots	a_1	a_0
x_0		$a_n x_0$	$b_{n-1} x_0$	\cdots	$b_1 x_0$	$b_0 x_0$
	a_n	$a_{n-1} + a_n x_0$	$a_{n-2} + b_{n-1} x_0$	\cdots	$a_1 + b_1 x_0$	$a_0 + b_0 x_0$

	1	-2	-5	6
-1		-1	3	
	1	-3	-2	

- Recall: Given p (degree n) and some x_0 , Horner's algorithm finds all coefficients b_j of q (degree $n - 1$) and $p(x_0)$
- Example: Deflate

$$p(x) = x^3 - 2x^2 - 5x + 6, \quad x_0 = -1$$

- Use synthetic division table:

	a_n	a_{n-1}	a_{n-2}	\cdots	a_1	a_0
x_0		$a_n x_0$	$b_{n-1} x_0$	\cdots	$b_1 x_0$	$b_0 x_0$
	a_n	$a_{n-1} + a_n x_0$	$a_{n-2} + b_{n-1} x_0$	\cdots	$a_1 + b_1 x_0$	$a_0 + b_0 x_0$

	1	-2	-5	6
-1		-1	3	2
	1	-3	-2	

- Recall: Given p (degree n) and some x_0 , Horner's algorithm finds all coefficients b_j of q (degree $n - 1$) and $p(x_0)$
- Example: Deflate

$$p(x) = x^3 - 2x^2 - 5x + 6, \quad x_0 = -1$$

- Use synthetic division table:

	a_n	a_{n-1}	a_{n-2}	\cdots	a_1	a_0
x_0		$a_n x_0$	$b_{n-1} x_0$	\cdots	$b_1 x_0$	$b_0 x_0$
	a_n	$a_{n-1} + a_n x_0$	$a_{n-2} + b_{n-1} x_0$	\cdots	$a_1 + b_1 x_0$	$a_0 + b_0 x_0$

	1	-2	-5	6
-1		-1	3	2
	1	-3	-2	8

- Recall: Given p (degree n) and some x_0 , Horner's algorithm finds all coefficients b_j of q (degree $n - 1$) and $p(x_0)$
- Example: Deflate

$$p(x) = x^3 - 2x^2 - 5x + 6, \quad x_0 = -1$$

- Use synthetic division table:

	a_n	a_{n-1}	a_{n-2}	\cdots	a_1	a_0
x_0		$a_n x_0$	$b_{n-1} x_0$	\cdots	$b_1 x_0$	$b_0 x_0$
	a_n	$a_{n-1} + a_n x_0$	$a_{n-2} + b_{n-1} x_0$	\cdots	$a_1 + b_1 x_0$	$a_0 + b_0 x_0$

	1	-2	-5	6
-1		-1	3	2
	1	-3	-2	8

- Thus, $p(x) = (x + 1)(x^2 - 3x - 2) + 8$

- Consider: n term Taylor expansion of $p(x)$ about $x = x_0$:

- Consider: n term Taylor expansion of $p(x)$ about $x = x_0$:

$$p(x) = p(x_0) + \frac{p'(x_0)}{1!}(x - x_0) + \dots + \frac{p^{(n)}(x_0)}{n!}(x - x_0)^n$$

- Consider: n term Taylor expansion of $p(x)$ about $x = x_0$:

$$p(x) = p(x_0) + \frac{p'(x_0)}{1!}(x - x_0) + \dots + \frac{p^{(n)}(x_0)}{n!}(x - x_0)^n$$

- Relabeling coefficients, we have

- Consider: n term Taylor expansion of $p(x)$ about $x = x_0$:

$$p(x) = p(x_0) + \frac{p'(x_0)}{1!}(x - x_0) + \dots + \frac{p^{(n)}(x_0)}{n!}(x - x_0)^n$$

- Relabeling coefficients, we have

$$p(x) = c_n(x - x_0)^n + c_{n-1}(x - x_0)^{n-1} + \dots + c_0$$

- Consider: n term Taylor expansion of $p(x)$ about $x = x_0$:

$$p(x) = p(x_0) + \frac{p'(x_0)}{1!}(x - x_0) + \dots + \frac{p^{(n)}(x_0)}{n!}(x - x_0)^n$$

- Relabeling coefficients, we have

$$p(x) = c_n(x - x_0)^n + c_{n-1}(x - x_0)^{n-1} + \dots + c_0$$

- Now, observe that repeatedly deflating p at x_0 yields

- Consider: n term Taylor expansion of $p(x)$ about $x = x_0$:

$$p(x) = p(x_0) + \frac{p'(x_0)}{1!}(x - x_0) + \dots + \frac{p^{(n)}(x_0)}{n!}(x - x_0)^n$$

- Relabeling coefficients, we have

$$p(x) = c_n(x - x_0)^n + c_{n-1}(x - x_0)^{n-1} + \dots + c_0$$

- Now, observe that repeatedly deflating p at x_0 yields

$$\begin{aligned} p(x) &= (x - x_0)q_1(x) + p(x_0) \\ &= (x - x_0)[(x - x_0)q_2(x) + q_1(x_0)] + p(x_0) \\ &\quad \vdots \\ &= \sum_{i=1}^n \left[(x - x_0)^i q_i(x_0) \right] + p(x_0). \end{aligned}$$

- Consider: n term Taylor expansion of $p(x)$ about $x = x_0$:

$$p(x) = p(x_0) + \frac{p'(x_0)}{1!}(x - x_0) + \dots + \frac{p^{(n)}(x_0)}{n!}(x - x_0)^n$$

- Relabeling coefficients, we have

$$p(x) = c_n(x - x_0)^n + c_{n-1}(x - x_0)^{n-1} + \dots + c_0$$

- Now, observe that repeatedly deflating p at x_0 yields

$$\begin{aligned} p(x) &= (x - x_0)q_1(x) + p(x_0) \\ &= (x - x_0)[(x - x_0)q_2(x) + q_1(x_0)] + p(x_0) \\ &\quad \vdots \\ &= \sum_{i=1}^n \left[(x - x_0)^i q_i(x_0) \right] + p(x_0). \end{aligned}$$

- Thus, the remainder terms from Horner's method are the coefficients we seek!

- Example: Consider

$$f(x) = x^3 - 2x^2 - 5x + 6.$$

Find Taylor expansion centered at $x = 2$.

- Example: Consider

$$f(x) = x^3 - 2x^2 - 5x + 6.$$

Find Taylor expansion centered at $x = 2$.

Table : Finding c_0

2	1	-2	-5	6
		2	0	-10
	1	0	-5	-4

- Example: Consider

$$f(x) = x^3 - 2x^2 - 5x + 6.$$

Find Taylor expansion centered at $x = 2$.

Table : Finding c_0

	1	-2	-5	6
2		2	0	-10
	1	0	-5	-4

Table : Finding c_1

	1	0	-5
2		2	4
	1	2	-1

- Example: Consider

$$f(x) = x^3 - 2x^2 - 5x + 6.$$

Find Taylor expansion centered at $x = 2$.

Table : Finding c_0

	1	-2	-5	6
2		2	0	-10
	1	0	-5	-4

Table : Finding c_1

	1	0	-5
2		2	4
	1	2	-1

Table : Finding c_2

	1	2
2		2
	1	4

- Example: Consider

$$f(x) = x^3 - 2x^2 - 5x + 6.$$

Find Taylor expansion centered at $x = 2$.

Table : Finding c_0

	1	-2	-5	6
2		2	0	-10
	1	0	-5	-4

Table : Finding c_1

	1	0	-5
2		2	4
	1	2	-1

Table : Finding c_2

	1	2
2		2
	1	4

- So, the Taylor expansion centered at $x = 2$ is

- Example: Consider

$$f(x) = x^3 - 2x^2 - 5x + 6.$$

Find Taylor expansion centered at $x = 2$.

Table : Finding c_0

2	1	-2	-5	6
		2	0	-10
	1	0	-5	-4

Table : Finding c_1

2	1	0	-5
		2	4
	1	2	-1

Table : Finding c_2

2	1	2
		2
	1	4

- So, the Taylor expansion centered at $x = 2$ is

$$f(x) = (x - 2)^3 + 4(x - 2)^2 - (x - 2) - 4.$$

- Recall: Newton's method is defined as

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$$

- Recall: Newton's method is defined as

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$$

- If f is a polynomial, use H.A. to evaluate $f(x_n)$ and $f'(x_n)$

- Recall: Newton's method is defined as

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$$

- If f is a polynomial, use H.A. to evaluate $f(x_n)$ and $f'(x_n)$
- How: H.A. can evaluate functions and Taylor series!

- Recall: Newton's method is defined as

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$$

- If f is a polynomial, use H.A. to evaluate $f(x_n)$ and $f'(x_n)$
- How: H.A. can evaluate functions and Taylor series!
- Code: evaluate both points simultaneously (pg. 115)

- Consider p and its roots ξ_i

- Consider p and its roots ξ_i
- Basin of attraction of ξ_i : set of all points which converge to ξ_i when iterated (say, by N.M.)

- Consider p and its roots ξ_i
- Basin of attraction of ξ_i : set of all points which converge to ξ_i when iterated (say, by N.M.)
- Question: what about the points that don't converge to any root?

- Consider p and its roots ξ_i
- Basin of attraction of ξ_i : set of all points which converge to ξ_i when iterated (say, by N.M.)
- Question: what about the points that don't converge to any root?
- Answer: form Julia set

- Consider p and its roots ξ_i
- Basin of attraction of ξ_i : set of all points which converge to ξ_i when iterated (say, by N.M.)
- Question: what about the points that don't converge to any root?
- Answer: form Julia set
- Demo!