

Are We There Yet?

Iterative Methods for Solving Linear Systems

Nate DeMaagd, Kurt O'Hearn

MTH 499-02

April 8, 2013

- Overview of methods of solution: direct, indirect

- Overview of methods of solution: direct, indirect
- Theory on iterative methods for solving linear systems

- Overview of methods of solution: direct, indirect
- Theory on iterative methods for solving linear systems
- Description and examples of Richardson, Jacobi, and Gauss-Seidel methods

- Direct methods

- Direct methods
 - Execute a predetermined number of computations to produce a result

- Direct methods
 - Execute a predetermined number of computations to produce a result
 - Example: Gaussian elimination with scaled partial pivoting

- Direct methods
 - Execute a predetermined number of computations to produce a result
 - Example: Gaussian elimination with scaled partial pivoting
 - Solves $Ax = b$ in a finite number of steps

- Direct methods
 - Execute a predetermined number of computations to produce a result
 - Example: Gaussian elimination with scaled partial pivoting
 - Solves $Ax = b$ in a finite number of steps
 - Gives exact answer (within roundoff errors)

- Direct methods
 - Execute a predetermined number of computations to produce a result
 - Example: Gaussian elimination with scaled partial pivoting
 - Solves $Ax = b$ in a finite number of steps
 - Gives exact answer (within roundoff errors)
- Indirect methods

- Direct methods
 - Execute a predetermined number of computations to produce a result
 - Example: Gaussian elimination with scaled partial pivoting
 - Solves $Ax = b$ in a finite number of steps
 - Gives exact answer (within roundoff errors)
- Indirect methods
 - Generate a sequence of intermediate results which (hopefully) produce the desired final result

- Direct methods
 - Execute a predetermined number of computations to produce a result
 - Example: Gaussian elimination with scaled partial pivoting
 - Solves $Ax = b$ in a finite number of steps
 - Gives exact answer (within roundoff errors)
- Indirect methods
 - Generate a sequence of intermediate results which (hopefully) produce the desired final result
 - Typically yield approximate solutions within some tolerance (ϵ) or after a prescribed number of iterations

- Direct methods
 - Execute a predetermined number of computations to produce a result
 - Example: Gaussian elimination with scaled partial pivoting
 - Solves $Ax = b$ in a finite number of steps
 - Gives exact answer (within roundoff errors)
- Indirect methods
 - Generate a sequence of intermediate results which (hopefully) produce the desired final result
 - Typically yield approximate solutions within some tolerance (ϵ) or after a prescribed number of iterations
 - Almost always are iterative in nature

- Direct methods
 - Execute a predetermined number of computations to produce a result
 - Example: Gaussian elimination with scaled partial pivoting
 - Solves $Ax = b$ in a finite number of steps
 - Gives exact answer (within roundoff errors)
- Indirect methods
 - Generate a sequence of intermediate results which (hopefully) produce the desired final result
 - Typically yield approximate solutions within some tolerance (ϵ) or after a prescribed number of iterations
 - Almost always are iterative in nature
 - Example: finding solutions to linear systems (today's topic!)

Developing A Simple Iterative Method

Developing A Simple Iterative Method

- Want to solve: $Ax = b$ for x with $A \in \mathbb{R}^{n \times n}$, A invertible

Developing A Simple Iterative Method

- Want to solve: $Ax = b$ for x with $A \in \mathbb{R}^{n \times n}$, A invertible
- Introduce an invertible “splitting matrix” Q and rearrange to get

$$Qx = (Q - A)x + b$$

Developing A Simple Iterative Method

- Want to solve: $Ax = b$ for x with $A \in \mathbb{R}^{n \times n}$, A invertible
- Introduce an invertible “splitting matrix” Q and rearrange to get

$$Qx = (Q - A)x + b$$

- Now define our iterative process as

$$Qx^{(k)} = (Q - A)x^{(k-1)} + b$$

where $k > 1$ denotes the k^{th} step in the process

Developing A Simple Iterative Method

- Want to solve: $Ax = b$ for x with $A \in \mathbb{R}^{n \times n}$, A invertible
- Introduce an invertible “splitting matrix” Q and rearrange to get

$$Qx = (Q - A)x + b$$

- Now define our iterative process as

$$Qx^{(k)} = (Q - A)x^{(k-1)} + b$$

where $k > 1$ denotes the k^{th} step in the process

- Want: each successive iteration to produce a better approximation for x (i.e., converge)

Developing A Simple Iterative Method

- Want to solve: $Ax = b$ for x with $A \in \mathbb{R}^{n \times n}$, A invertible
- Introduce an invertible “splitting matrix” Q and rearrange to get

$$Qx = (Q - A)x + b$$

- Now define our iterative process as

$$Qx^{(k)} = (Q - A)x^{(k-1)} + b$$

where $k > 1$ denotes the k^{th} step in the process

- Want: each successive iteration to produce a better approximation for x (i.e., converge)
- Also want: algorithms which guarantee convergence after satisfying some conditions

Developing A Simple Iterative Method

- Want to solve: $Ax = b$ for x with $A \in \mathbb{R}^{n \times n}$, A invertible
- Introduce an invertible “splitting matrix” Q and rearrange to get

$$Qx = (Q - A)x + b$$

- Now define our iterative process as

$$Qx^{(k)} = (Q - A)x^{(k-1)} + b$$

where $k > 1$ denotes the k^{th} step in the process

- Want: each successive iteration to produce a better approximation for x (i.e., converge)
- Also want: algorithms which guarantee convergence after satisfying some conditions
- To achieve these ends, we seek Q such that:
 - $\|x - x^{(k)}\| \rightarrow 0$ rapidly, and
 - $[x^{(k)}]$ is easy to compute

Developing A Simple Iterative Method

- Want to solve: $Ax = b$ for x with $A \in \mathbb{R}^{n \times n}$, A invertible
- Introduce an invertible “splitting matrix” Q and rearrange to get

$$Qx = (Q - A)x + b$$

- Now define our iterative process as

$$Qx^{(k)} = (Q - A)x^{(k-1)} + b$$

where $k > 1$ denotes the k^{th} step in the process

- Want: each successive iteration to produce a better approximation for x (i.e., converge)
- Also want: algorithms which guarantee convergence after satisfying some conditions
- To achieve these ends, we seek Q such that:
 - $\|x - x^{(k)}\| \rightarrow 0$ rapidly, and
 - $[x^{(k)}]$ is easy to compute
- Note: often the initial vector $x^{(0)}$ is an estimate of the solution or arbitrary ($x = 0$)

- Recall iterative process $Qx^{(k)} = (Q - A)x^{(k-1)} + b$

- Recall iterative process $Qx^{(k)} = (Q - A)x^{(k-1)} + b$
- Letting $k \rightarrow \infty$, solution is $Qx = (Q - A)x + b$

- Recall iterative process $Qx^{(k)} = (Q - A)x^{(k-1)} + b$
- Letting $k \rightarrow \infty$, solution is $Qx = (Q - A)x + b$
- Assume Q and A nonsingular. So, Q^{-1} and A^{-1} exist and

$$x^{(k)} = (I - Q^{-1}A)x^{(k-1)} + Q^{-1}b$$

- Recall iterative process $Qx^{(k)} = (Q - A)x^{(k-1)} + b$
- Letting $k \rightarrow \infty$, solution is $Qx = (Q - A)x + b$
- Assume Q and A nonsingular. So, Q^{-1} and A^{-1} exist and

$$x^{(k)} = (I - Q^{-1}A)x^{(k-1)} + Q^{-1}b$$

- Taking limit of this gives solution

$$x = (I - Q^{-1}A)x + Q^{-1}b$$

- Recall iterative process $Qx^{(k)} = (Q - A)x^{(k-1)} + b$
- Letting $k \rightarrow \infty$, solution is $Qx = (Q - A)x + b$
- Assume Q and A nonsingular. So, Q^{-1} and A^{-1} exist and

$$x^{(k)} = (I - Q^{-1}A)x^{(k-1)} + Q^{-1}b$$

- Taking limit of this gives solution

$$x = (I - Q^{-1}A)x + Q^{-1}b$$

- Thus $x^{(k)} - x = (I - Q^{-1}A)(x^{(k-1)} - x)$

- Recall iterative process $Qx^{(k)} = (Q - A)x^{(k-1)} + b$
- Letting $k \rightarrow \infty$, solution is $Qx = (Q - A)x + b$
- Assume Q and A nonsingular. So, Q^{-1} and A^{-1} exist and

$$x^{(k)} = (I - Q^{-1}A)x^{(k-1)} + Q^{-1}b$$

- Taking limit of this gives solution

$$x = (I - Q^{-1}A)x + Q^{-1}b$$

- Thus $x^{(k)} - x = (I - Q^{-1}A)(x^{(k-1)} - x)$
- Select vector norm and subordinate norm so that by repeated
[[repeated what?]]

$$\|x^{(k)} - x\| \leq \|I - Q^{-1}A\|^k \|x^{(0)} - x\|$$

- Recall iterative process $Qx^{(k)} = (Q - A)x^{(k-1)} + b$
- Letting $k \rightarrow \infty$, solution is $Qx = (Q - A)x + b$
- Assume Q and A nonsingular. So, Q^{-1} and A^{-1} exist and

$$x^{(k)} = (I - Q^{-1}A)x^{(k-1)} + Q^{-1}b$$

- Taking limit of this gives solution

$$x = (I - Q^{-1}A)x + Q^{-1}b$$

- Thus $x^{(k)} - x = (I - Q^{-1}A)(x^{(k-1)} - x)$
- Select vector norm and subordinate norm so that by repeated [[repeated what?]]

$$\|x^{(k)} - x\| \leq \|I - Q^{-1}A\|^k \|x^{(0)} - x\|$$

- Thus, if $\|I - Q^{-1}A\| < 1$, then $\lim_{k \rightarrow \infty} \|x^{(k)} - x\| = 0$ [[How does this follow from the last step?]]