

- Lowpass FIR filter

simple MA filter -


$$h[n] = \begin{cases} 1/m & , 0 \leq n \leq m-1 \\ 0 & , \text{otherwise} \end{cases}$$

Let $m=2$

$$H_0(z) = \sum_{n=0}^1 \frac{1}{2} z^{-n} = \frac{1}{2} (1 + z^{-1}) = \frac{z+1}{2z}$$


pole at $z=0$

zero at $z=-1$



Pole vector magnitude is unity for all ω . Zero vector magnitude decreased as ω approaches π .

$$|H_0(e^{j0})| = 1, \quad |H_0(e^{j\pi})| = 0$$



$$H_0(e^{j\omega}) = e^{-j\omega/2} \cos(\omega/2)$$

What is cutoff freq? $|H_0(e^{j\omega})| = \frac{1}{\sqrt{2}} |H_0(e^{j0})|$

Use dB -

$$20 \log_{10} |H_0(e^{j\omega})| - 20 \log_{10} \sqrt{2} \approx -3.01 \text{ dB}$$

$$|H_0(e^{j\omega_c})|^2 = \cos^2\left(\frac{\omega_c}{2}\right) = \frac{1}{2}$$

$$\Rightarrow \omega_c = \pi/2 \quad \text{power is halved at } \pi/2$$

- Highpass FIR filters



Replace z w/ z^{-1} in MA filter

$$H_1(z) = \frac{1}{2} (1 - z^{-1}) = \frac{z-1}{2}$$

zero at $z=1$
pole at $z=0$

Freq. Response -

$$H_1(e^{j\omega}) = j e^{-j\omega/2} \sin(\omega/2)$$



- Inverse z -transform

Recover DT sequence from its z -transform

$$x[n] = \frac{1}{2\pi j} \oint_C X(z) z^{n-1} dz$$

Two useful methods to avoid contour integration -

1. Partial Fraction Expansion

For rational functions of z , simple poles

Assume $N > M$
 $P \rightarrow Q$

$$X(z) = \frac{\sum_{k=0}^{QM} b_k z^{-k}}{\sum_{k=0}^{PN} a_k z^{-k}}$$

$$\Rightarrow X(z) = \sum_{k=1}^N \frac{A_k}{1 - d_k z^{-1}} \quad \Rightarrow A_k = X(z) (1 - d_k z^{-1}) \big|_{z=d_k}$$

Example $X(z) = \frac{1}{(1 - \frac{1}{4}z^{-1})(1 - \frac{1}{2}z^{-1})}$ $|z| > \frac{1}{2} \Rightarrow$ right sided sequence.

$$X(z) = \frac{A_1}{1 - \frac{1}{4}z^{-1}} + \frac{A_2}{1 - \frac{1}{2}z^{-1}}$$

$$A_1 = (1 - \frac{1}{2}z^{-1}) X(z) \Big|_{z=1/4} = \frac{(1 - \frac{1}{2}z^{-1})}{(1 - \frac{1}{4}z^{-1})(1 - \frac{1}{2}z^{-1})} \Big|_{z=1/4} = -1$$

$$A_2 = (1 - \frac{1}{4}z^{-1}) X(z) \Big|_{z=1/2} = \frac{(1 - \frac{1}{4}z^{-1})}{(1 - \frac{1}{4}z^{-1})(1 - \frac{1}{2}z^{-1})} \Big|_{z=1/2} = 2$$

so, $X(z) = \frac{-1}{(1 - \frac{1}{4}z^{-1})} + \frac{2}{(1 - \frac{1}{2}z^{-1})} \Rightarrow x[n] = 2\left(\frac{1}{2}\right)^n u[n] - \left(\frac{1}{4}\right)^n u[n]$

2- Power series Expansion - The z-transform is a power series expansion:-

$$X(z) = \sum_{n=-\infty}^{\infty} x[n] z^{-n} = \dots x[-2]z^2 + x[-1]z^1 + x[0] + x[1]z^{-1} + \dots$$

Find power series expansion form of $X(z)$ and then find OT sequence by inspection.

Example $X(z) = \frac{z^{-1}}{(1 - z^{-1})^2}$ $|z| > 1$ use binomial expansion.

Binomial Expansion $(1+x)^\alpha = 1 + \alpha x + \frac{\alpha(\alpha-1)}{2!} x^2 + \dots$

$$\alpha = -2, \quad x = -z^{-1}$$

$$\Rightarrow \frac{1}{(1-z^{-1})^2} = 1 + 2z^{-1} + 3z^{-2} + 4z^{-3} + \dots$$

$$\Rightarrow \frac{z}{(1-z^{-1})^2} = z^{-1} + 2z^{-2} + 3z^{-3} + 4z^{-4} + \dots$$

So, $x[n] = 1\delta[n-1] + 2\delta[n-2] + 3\delta[n-3] + 4\delta[n-4] + \dots$

$$x[n] = \begin{cases} n, & n \geq 0 \\ 0, & n < 0 \end{cases}$$

3- Contour Integration

Cauchy's Integral Theorem - If C is a closed contour that encircles the origin in a counter-clockwise direction -

$$\frac{1}{2\pi j} \oint_C z^{-k} dz = \begin{cases} 1 \\ 0 \end{cases}$$

- Linear phase systems

An LTI system has linear phase if

$$H(e^{j\omega}) = |H(e^{j\omega})| e^{-j\alpha\omega} \quad \alpha \text{ is real valued}$$

Linear phase systems have constant group delay

$$\tau_g = -\frac{d\phi(\omega)}{d\omega} = -\frac{d(-\alpha\omega)}{d\omega} = \alpha$$

Note: In order for causal system w/ rational transfer function to have linear phase, the impulse response must be finite in length.

\Rightarrow IIR filters cannot have linear phase!

- 4 types of linear phase FIR systems.

Type I - Symmetrical impulse response, and N is even

$$h[n] = h[N-n] \quad 0 \leq n \leq N$$

• The delay through the filter is $N/2$

Type II - Symmetrical impulse response, and N is odd
• delay is $N/2$

Type III - anti symmetrical impulse response, and N is even

$$h[n] = -h[N-n] \quad 0 \leq n \leq N$$

• delay is $N/2$

Type IV - anti symmetrical impulse response, and N is odd

• delay is $N/2$

- Locations of zeros for FIR Linear phase systems.

Type I & II -
$$H(z) = \sum_{n=0}^N h[n] z^{-n} = \sum_{k=0}^N h[N-k] z^{-k}$$

Let $m = N - n$
$$= \sum_{m=0}^N h[m] z^{-N+m} = z^{-N} \sum_{m=0}^N h[m] z^m = z^{-N} H(z^{-1})$$

If z_0 is a zero, $H(z_0) = z_0^{-N} H(z_0^{-1}) = 0$
 so, z_0^{-1} must be a zero.

Note; For $h[n]$ w/ real-valued coeff., the zeros occur in complex conjugate pairs. Hence, a zero at $z = z_0$ is associated with a set of four zeros given by $z_0, z_0^*, z_0^{-1}, z_0^{-1*}$.

Important case - zero at $z = -1$

If N is even $H(-1) = (-1)^N H(-1) = H(-1)$

If N is odd $H(-1) = (-1) H(-1) \Rightarrow H(-1)$ must be zero - always!

For symmetric $h[n]$ with odd N , system must have zero at $z = -1$. Freq Resp is constrained to 0 at $\omega = \pi$.
 Do not use for highpass filter.

Type III & IV -
$$H(z) = -z^{-N} H(z^{-1})$$

Both $z = 1$ & $z = -1$ are special cases.

If $z = 1$ $H(1) = -H(1) \Rightarrow$ must have zero at $z = 1$
 for both N even and odd

If $z = -1$ $H(-1) = (-1)^{-N} H(-1) \Rightarrow z = -1$ must be zero for N even