

# CHAPTER 3: Number Systems

---

## **The Architecture of Computer Hardware and Systems Software & Networking: An Information Technology Approach**


**5th Edition, Irv Englander**  
**John Wiley and Sons ©2013**

PowerPoint slides authored by Angela Clark, University of South Alabama  
PowerPoint slides for the 4<sup>th</sup> edition were authored by Wilson Wong, Bentley  
University





# Counting and Arithmetic


- Decimal or base 10 number system
  - Origin: counting on the fingers
  - “Digit” from the Latin word *digitus* meaning “finger”
- **Base**: the number of different digits, including zero, in the number system
  - Example: Base 10 has 10 digits, 0 through 9
- **Binary** or **base 2**
  - **Bit** (short for *binary digit*): 2 digits, 0 and 1
- **Octal** or **base 8**: 8 digits, 0 through 7
- **Hexadecimal** or **base 16**: 16 digits, 0 through F
- Examples:  $10_{10} = A_{16}$ ;  $11_{10} = B_{16}$



# Why Binary?

- Early computer design was decimal
  - Mark I and ENIAC
- John von Neumann proposed binary data processing (1945)
  - Simplified computer design
  - Used for both instructions and data
- Natural relationship between on/off switches and calculations using Boolean logic


	
On	Off
True	False
Yes	No
1	0



# Keeping Track of the Bits


---

- Bits commonly stored and manipulated in groups
  - 8 bits = 1 *byte*
  - 4 bytes = 1 *word* (in many systems)
- Number of bits used in calculations
  - Affects accuracy of results
  - Limits size and range of numbers manipulated by the computer



# Numbers: Physical Representation

- Different numerals, same number of oranges
  - Cave dweller: IIII
  - Roman: V
  - Arabic: 5
- Different bases, same number of oranges
  - $5_{10}$
  - $101_2$
  - $12_3$





# Number Systems

---

- Roman: position *independent*, simple *additive* system
  - Example: XII is a ten and two ones, i.e. 12
- Modern systems: based on positional notation (place value)
  - Decimal system: based on powers of 10
  - Binary system: based powers of 2
  - Octal system: based on powers of 8
  - Hexadecimal system: based powers of 16



# Positional Notation: Base 10

$$527_{10} = 5 \times 10^2 + 2 \times 10^1 + 7 \times 10^0$$

100's place

10's place




1's place

Place	$10^2$	$10^1$	$10^0$
Value	100	10	1
Evaluate	$5 \times 100$	$2 \times 10$	$7 \times 1$
Sum	500	20	7



# Positional Notation: Octal

$$624_8 = 404_{10}$$

	64's place	8's place	1's place
			
Place	$8^2$	$8^1$	$8^0$
Value	64	8	1
Evaluate	6 x 64	2 x 8	4 x 1
Sum for Base 10	384	16	4





# Positional Notation: Hexadecimal

$$6,704_{16} = 26,372_{10}$$

4,096's place      256's place      16's place      1's place



Place	$16^3$	$16^2$	$16^1$	$16^0$
Value	4,096	256	16	1
Evaluate	6 x 4,096	7 x 256	0 x 16	4 x 1
Sum for Base 10	24,576	1,792	0	4



# Positional Notation: Binary


$$1101\ 0110_2 = 214_{10}$$

Place	$2^7$	$2^6$	$2^5$	$2^4$	$2^3$	$2^2$	$2^1$	$2^0$
Value	128	64	32	16	8	4	2	1
Evaluate	1 x 128	1 x 64	0 x 32	1 x 16	0 x 8	1 x 4	1 x 2	0 x 1
Sum for Base 10	128	64	0	16	0	4	2	0



# Range of Possible Numbers

- $R = B^K$ 
  - $R$  = range
  - $B$  = base
  - $K$  = number of digits
- Example #1: Base 10, 2 digits
  - $R = 10^2 = 100$  different numbers (0...99)
- Example #2: Base 2, 16 digits
  - $R = 2^{16} = 65,536$  or 64K
  - 16-bit number can represent 65,536 different number values



# Decimal Range for Bit Widths

Bits	Digits	Range
1	0+	2 (0 and 1)
4	1+	16 (0 to 15)
8	2+	256
10	3	1,024 (1K)
16	4+	65,536 (64K)
20	6	1,048,576 (1M)
32	9+	4,294,967,296 (4G)
64	19+	Approx. $1.6 \times 10^{19}$
128	38+	Approx. $2.6 \times 10^{38}$



# Different Number Bases

---

- Base:
  - The number of different symbols required to represent any given number
- The *larger* the base, the *more* symbols are required
  - Base 10: 0,1,2,3,4,5,6,7,8,9
  - Base 2: 0,1
  - Base 8: 0,1,2,3,4,5,6,7
  - Base 16: 0,1,2,3,4,5,6,7,8,9,A,B,C,D,E,F



# Number of Symbols vs. Number of Digits

---

- For a given number, the *larger* the base
  - the *more* symbols required
  - but the *fewer* digits needed
- Example #1:
  - $65_{16}$      $101_{10}$      $145_8$      $110\ 0101_2$
- Example #2:
  - $11C_{16}$      $284_{10}$      $434_8$      $1\ 0001\ 1100_2$



# Counting in Base 2

Binary Number	Equivalent				Decimal Number
	8's ( $2^3$ )	4's ( $2^2$ )	2's ( $2^1$ )	1's ( $2^0$ )	
0				$0 \times 2^0$	0
1				$1 \times 2^0$	1
10			$1 \times 2^1$	$0 \times 2^0$	2
11			$1 \times 2^1$	$1 \times 2^0$	3
100		$1 \times 2^2$			4
101		$1 \times 2^2$		$1 \times 2^0$	5
110		$1 \times 2^2$	$1 \times 2^1$		6
111		$1 \times 2^2$	$1 \times 2^1$	$1 \times 2^0$	7
1000	$1 \times 2^3$				8
1001	$1 \times 2^3$			$1 \times 2^0$	9
1010	$1 \times 2^3$		$1 \times 2^1$		10



# Base 10 Addition Table

$$3_{10} + 6_{10} = 9_{10}$$

+	0	1	2	3	4	5	6	7	8	9
0	0	1	2	3	4	5	6	7	8	9
1	1	2	3	4	5	6	7	8	9	10
2	2	3	4	5	6	7	8	9	10	11
3	3	4	5	6	7	8	9	10	11	12
4	4	5	6	7	8	9	10	11	12	13






# Base 8 Addition Table

$$3_8 + 6_8 = 11_8$$

+	0	1	2	3	4	5	6	7
0	0	1	2	3	4	5	6	7
1	1	2	3	4	5	6	7	10
2	2	3	4	5	6	7	10	11
3	3	4	5	6	7	10	11	12
4	4	5	6	7	10	11	12	13
5	5	6	7	10	11	12	13	14
6	6	7	10	11	12	13	14	15
7	7	10	11	12	13	14	15	16

(no 8 or 9,  
of course)



# Base 10 Multiplication Table

$$3_{10} \times 6_{10} = 18_{10}$$

x	0	1	2	3	4	5	6	7	8	9
0				0						
1		1	2	3	4	5	6	7	8	9
2		2	4	6	8	10	12	14	16	18
3		3	6	9	12	15	18	21	24	27
4	0	4	8	12	16	20	24	28	32	36
5		5	10	15	20	25	30	35	40	45
6		6	12	18	24	30	36	42	48	54
7		7	14	21	28	35	42	49	56	63



# Base 8 Multiplication Table

$$3_8 \times 6_8 = 22_8$$

x	0	1	2	3	4	5	6	7
0	0	1	2	3	4	5	6	7
1	1	2	4	6	10	12	14	16
2	2	4	6	10	12	14	16	17
3	3	6	11	14	17	22	25	26
4	4	10	14	20	24	30	34	35
5	5	12	17	24	31	36	43	44
6	6	14	22	30	36	44	52	53
7	7	16	25	34	43	52	61	62



# Addition

Base	Problem	Largest Single Digit
Decimal	$\begin{array}{r} 6 \\ +3 \\ \hline \end{array}$	9
Octal	$\begin{array}{r} 6 \\ +1 \\ \hline \end{array}$	7
Hexadecimal	$\begin{array}{r} 6 \\ +9 \\ \hline \end{array}$	F
Binary	$\begin{array}{r} 1 \\ +0 \\ \hline \end{array}$	1




# Addition

Base	Problem	Carry	Answer
Decimal	$\begin{array}{r} 6 \\ +4 \\ \hline \end{array}$	Carry the 10	10
Octal	$\begin{array}{r} 6 \\ +2 \\ \hline \end{array}$	Carry the 8	10
Hexadecimal	$\begin{array}{r} 6 \\ +A \\ \hline \end{array}$	Carry the 16	10
Binary	$\begin{array}{r} 1 \\ +1 \\ \hline \end{array}$	Carry the 2	10



# Binary Arithmetic

$$\begin{array}{rcccccccc} & 1 & 1 & & 1 & & 1 & & 1 & & & & & & \\ & & & & 1 & & 1 & & 0 & & 1 & & 1 & & 0 & & 1 & \\ + & & & & & & & & 1 & & 0 & & 1 & & 1 & & 1 & & 0 & \\ \hline 1 & & 0 & & 0 & & 0 & & 0 & & 0 & & 0 & & 1 & & 1 & & & \end{array}$$



# Binary Arithmetic

- Addition

- Boolean using XOR and AND

- Multiplication

- AND
- Shift *left*

- Division

- Subtraction or complement
- Shift *right*

		0	1
+			
0		0	1
1		1	10

		0	1
x			
0		0	0
1		0	1



# Binary Arithmetic: Boolean Logic

- Boolean logic without performing arithmetic
  - *EXCLUSIVE-OR*
    - Output is “1” only if either input, but **not** both inputs, is a “1”
  - *AND (carry bit)*
    - Output is “1” if and only both inputs are a “1”

1	1	1	1	1			
	1	1	0	1	1	0	1
+			1	0	1	1	0
<hr/>							
1	0	0	0	0	0	1	1





# Binary Multiplication

- Boolean logic without performing arithmetic
  - *AND (carry bit)*
    - ▢ Output is “1” if and only both inputs are a “1”
  - *Shift*
    - ▢ Shifting a number in any base *left* one digit *multiplies* its value by the base
    - ▢ Shifting a number in any base *right* one digit *divides* its value by the base
    - ▢ Examples:
      - ▢  $10_{10}$  shift *left* =  $100_{10}$       ▢  $10_{10}$  shift *right* =  $1_{10}$
      - ▢  $10_2$  shift *left* =  $100_2$       ▢  $10_2$  shift *right* =  $1_2$



# Binary Multiplication

$$\begin{array}{r} 1101 \\ 101 \\ \hline 1101 \quad \text{1's place} \\ 0 \quad \text{2's place} \\ 1101 \quad \text{4's place (bits shifted to line up with 4's place of multiplier)} \\ \hline 1000001 \quad \text{Result (AND)} \end{array}$$



# Converting from Base 10

## ■ Powers Table

Power Base	8	7	6	5	4	3	2	1	0
2	256	128	64	32	16	8	4	2	1
8				32,768	4,096	512	64	8	1
16					65,536	4,096	256	16	1



# From Base 10 to Base 2

$$42_{10} = 101010_2$$

Power Base	6	5	4	3	2	1	0
2	64	32	16	8	4	2	1
		1	0	1	0	1	0
Integer		$42/32 = 1$	$10/16 = 0$	$10/8 = 1$	$2/4 = 0$	$2/2 = 1$	$0/1 = 0$
Remainder		10	10	2	2	0	0



# From Base 10 to Base 2

Base 10    42

Quotient    2 ) 42 ( 0    Remainder  
                    2 ) 21 ( 1    Least significant bit  
                    2 ) 10 ( 0  
                    2 ) 5 ( 1  
                    2 ) 2 ( 0  
                    2 ) 1  
Base 2    101010

Most significant bit



# From Base 10 to Base 16

$$5,735_{10} = 1667_{16}$$

Power Base	4	3	2	1	0
16	65,536	4,096	256	16	1
		1	6	6	7
Integer		$5,735 / 4,096$ $= 1$	$1,639 / 256$ $= 6$	$103 / 16$ $= 6$	7
Remainder		$5,735 - 4,096$ $= 1,639$	$1,639 - 1,536$ $= 103$	$103 - 96$ $= 7$	



# From Base 10 to Base 16

**Base 10 8,039**

Remainder

Quotient

16 )	8,039	( 7	Least significant bit
16 )	502	( 6	
16 )	31	( 15	
16 )	1	( 1	Most significant bit
16 )	0		

**Base 16 1F67**



# From Base 8 to Base 10

$$7263_8 = 3,763_{10}$$

Power	$8^3$	$8^2$	$8^1$	$8^0$
	512	64	8	1
	x 7	x 2	x 6	x 3
Sum for Base 10	3,584	128	48	3





# From Base 8 to Base 10

$$7263_8 = 3,763_{10}$$

$$\begin{array}{r} 7 \\ \times 8 \\ \hline 56 \end{array} + 2 = 58$$

$$\begin{array}{r} \phantom{4} \times 8 \\ \hline 464 \end{array} + 6 = 470$$

$$\begin{array}{r} \phantom{37} \times 8 \\ \hline 3760 \end{array} + 3 = 3,763$$



# From Base 16 to Base 2

- Why hexadecimal?
  - Hex easier to read and write than binary
  - Modern computer operating systems and networks present variety of troubleshooting data in hex format
- The **nibble** approach

Base 16	1	F	6	7
Base 2	0001	1111	0110	0111



# Fractions

---

- Number point or radix point
  - Decimal point in base 10
  - Binary point in base 2
- No exact relationship between fractional numbers in different number bases
  - Exact conversion may be impossible, such as in the case of  $1/3$  or  $0.3333333...$



# Decimal Fractions

- Move the number point one place to the right
  - Effect: multiplies the number by the base number
  - Example:  $139.0_{10} \longrightarrow 1390_{10}$
- Move the number point one place to the left
  - Effect: divides the number by the base number
  - Example:  $139.0_{10} \longrightarrow 13.9_{10}$



# Fractions: Base 10 and Base 2

$$.2589_{10} = 2 \times 10^{-1} + 5 \times 10^{-2} + 8 \times 10^{-3} + 9 \times 10^{-4}$$

Place	$10^{-1}$	$10^{-2}$	$10^{-3}$	$10^{-4}$
Value	1/10	1/100	1/1000	1/10000
Evaluate	$2 \times 1/10$	$5 \times 1/100$	$8 \times 1/1000$	$9 \times 1/1000$
Sum	.2	.05	.008	.0009

$$.101011_2 = 0.671875_{10}$$

Place	$2^{-1}$	$2^{-2}$	$2^{-3}$	$2^{-4}$	$2^{-5}$	$2^{-6}$
Value	1/2	1/4	1/8	1/16	1/32	1/64
Evaluate	$1 \times 1/2$	$0 \times 1/4$	$1 \times 1/8$	$0 \times 1/16$	$1 \times 1/32$	$1 \times 1/64$
Sum	.5		0.125		0.03125	0.015625



# Fractions: Base 10 and Base 2

---

- No general relationship between fractions of types  $1/10^k$  and  $1/2^k$ 
  - Therefore a number representable in base 10 may not be representable in base 2
  - But the converse is true: all fractions of the form  $1/2^k$  can be represented in base 10
- Fractional conversions from one base to another are stopped
  - If there is a rational solution or
  - When the desired accuracy is attained



# Mixed Number Conversion

---

- Integer and fractional parts must be converted separately
- Radix point: fixed reference for the conversion
  - Digit to the left is a unit digit in every base
  - $B^0$  is always equal to 1 regardless of the base



# Copyright 2013 John Wiley & Sons

---

All rights reserved. Reproduction or translation of this work beyond that permitted in section 117 of the 1976 United States Copyright Act without express permission of the copyright owner is unlawful. Request for further information should be addressed to the Permissions Department, John Wiley & Sons, Inc. The purchaser may make back-up copies for his/her own use only and not for distribution or resale. The Publisher assumes no responsibility for errors, omissions, or damages caused by the use of these programs or from the use of the information contained herein.”