

Supporting Information:

**Rotational dynamics of proteins from spin
relaxation times and molecular dynamics
simulations**

O. H. Samuli Ollila,^{*,†,‡} Harri Heikkinen,[†] and Hideo Iwai[†]

[†]Institute of Biotechnology, University of Helsinki

*[‡]Institute of Organic Chemistry and Biochemistry, Czech Academy of Sciences, Prague 6,
Czech Republic*

E-mail: samuli.ollila@helsinki.fi

S1 Supplementary Figures




Figure S1: Mean square angle deviations of inertia tensor axes calculated from *PaTonB-96* simulation with tip4p water model at 310K. The data shown with linear (left) and logarithmic scale (right).




Figure S2: Mean square angle deviations of inertia tensor axes calculated from *PaTonB-96* simulation with tip4p water model at 298K. The data shown with linear (left) and logarithmic scale (right).

Table S1: Scaling factors of rotational diffusion for different proteins simulated with different water models. The last row shows the ratio of simulated and experimental water self-diffusion constant.

	tip3p	tip4p	OPC4
<i>Hp</i> TonB-92	2.9	1	-
<i>Pa</i> TonB-96	-	1.2	-
Calcium recoverin ¹	3.2	-	-
CBM-64 ²	-	-	1.3
65K C-RRM ³	-	1	-
Water self-diffusion ⁴	2.4	1.1	1

References

- (1) Timr, S.; Kadlec, J.; Srb, P.; Ollila, O. H. S.; Jungwirth, P. Calcium Sensing by Recoverin: Effect of Protein Conformation on Ion Affinity. In Preparation.
- (2) Heikkinen, H. A.; Iwai, H. In Preparation.
- (3) Norppa, A. J.; Kauppala, T. M.; Heikkinen, H. A.; Verma, B.; Iwai, H.; Frilander, M. J. Mutations in the U11/U12-65K protein associated with isolated growth hormone deficiency lead to structural destabilization and impaired binding of U12 snRNA. 2017; 10.1261/rna.062844.117.
- (4) Izadi, S.; Anandakrishnan, R.; Onufriev, A. V. Building Water Models: A Different Approach. *J. Phys. Chem. Lett.* **2014**, 5, 3863–3871.