



# Ostrich

## Journal of African Ornithology

ISSN: 0030-6525 (Print) 1727-947X (Online) Journal homepage: <http://www.tandfonline.com/loi/tost20>

## Managing human disturbance: factors influencing flight-initiation distance of birds in a West African nature reserve

Bukola Braimoh, Soladoye Iwajomo, Mark Wilson, Adams Chaskda, Afan Ajang & Will Cresswell

To cite this article: Bukola Braimoh, Soladoye Iwajomo, Mark Wilson, Adams Chaskda, Afan Ajang & Will Cresswell (2017): Managing human disturbance: factors influencing flight-initiation distance of birds in a West African nature reserve, Ostrich, DOI: [10.2989/00306525.2017.1388300](https://doi.org/10.2989/00306525.2017.1388300)

To link to this article: <https://doi.org/10.2989/00306525.2017.1388300>



Published online: 13 Dec 2017.



Submit your article to this journal [↗](#)



Article views: 14



View related articles [↗](#)



View Crossmark data [↗](#)

# Managing human disturbance: factors influencing flight-initiation distance of birds in a West African nature reserve

Bukola Braimoh<sup>1,2\*</sup>, Soladoye Iwajomo<sup>3</sup>, Mark Wilson<sup>4</sup>, Adams Chaskda<sup>1</sup>, Afan Ajang<sup>1</sup> and Will Cresswell<sup>1,5</sup>

<sup>1</sup> AP Leventis Ornithological Research Institute, University of Jos, Jos, Nigeria

<sup>2</sup> Animal Demography Unit, Department of Biological Sciences, University of Cape Town, Cape Town, South Africa

<sup>3</sup> Department of Zoology, Faculty of Science, University of Lagos, Akoka, Yaba, Nigeria

<sup>4</sup> British Trust for Ornithology, University of Stirling, Stirling, FK9 4LA, UK

<sup>5</sup> Centre for Biological Diversity, University of St Andrews, St Andrews, Fife KY16 9TH, UK

\* Corresponding author, email: [godsbatlax@yahoo.ca](mailto:godsbatlax@yahoo.ca)

Escape behaviour in response to perceived predators can be employed as a guide when designating protected areas around sensitive bird species to minimise the impact of human disturbance. A key measure of escape response is flight-initiation distance (FID), the distance at which a prey animal initiates its escape when approached by a potential predator. We tested the predictions of optimal escape theory by determining the factors that influence FID of bird species in a Nigerian reserved area and its surrounding habitats, and so the potential utility of FID in managing human disturbance on birds, for the first time within a West African context. We tested how FID varied with group size, proximity to vegetation acting as protective cover, levels of human use, and survival rate, and whether these relationships varied by species. We collected 504 FIDs for seven bird species in Amurum Forest Reserve and its surrounding habitats (Jos, Nigeria). The FID was lower in larger groups and when species were closer to protective cover. The FID was lower outside of the protected area because animals in sites with higher levels of human presence and use may become habituated. The FID was higher for species with higher survival, being consistent with predictions from life history theory. Overall, birds perceived humans as a potential threat and responded in accordance to the predictions of optimal escape theory, with FID increasing with increased cost of staying. Reserve managers in Africa could use species- and context-specific FIDs to designate buffer distances for the protection of wildlife from human disturbance.

## Gestion des perturbations humaines: facteurs influençant la distance d'initiation au vol des oiseaux dans une réserve naturelle d'Afrique de l'Ouest

Le comportement de fuite en réponse aux prédateurs perçus peut être utilisé comme guide lors de la désignation d'aires protégées autour d'espèces d'oiseaux sensibles afin de minimiser l'impact des perturbations humaines. Une mesure clé de la réponse de fuite est la distance d'initiation de vol « flight-initiation distance » (FID), la distance à laquelle une proie déclenche sa fuite lorsqu'elle est approchée par un prédateur potentiel. Nous avons testé les prédictions de la théorie de la fuite optimale en déterminant les facteurs qui influencent le FID des espèces d'oiseaux dans une zone réserve nigériane et ses habitats environnants et donc l'utilité potentielle du FID dans la gestion des perturbations humaines sur les oiseaux, pour la première fois dans un contexte d'Afrique de l'Ouest. Nous avons testé la façon dont le FID a varié avec la taille du groupe, la proximité de la végétation agissant comme couverture protectrice, les niveaux d'utilisation humaine et le taux de survie, et si ces relations variaient selon les espèces. Nous avons recueilli 504 FID pour sept espèces d'oiseaux dans la Réserve forestière d'Amurum et ses habitats environnants (Jos, Nigeria). Le FID était plus faible dans les grands groupes et lorsque les espèces étaient plus proches de la couverture protectrice. Le FID était plus faible à l'extérieur de la zone protégée parce que les animaux dans les sites ayant des niveaux plus élevés de présence et d'utilisation humaine peuvent être habitués. Le FID était plus élevé pour les espèces ayant une survie plus élevée, ce qui était conforme aux prédictions de la théorie de l'histoire de la vie. Dans l'ensemble, les oiseaux ont perçu l'homme comme une menace potentielle et ont répondu en fonction des prédictions de la théorie de fuite optimale, le FID augmentant avec un coût accru de séjour. Les gestionnaires des réserves en Afrique pourraient utiliser des FID spécifiques aux espèces et au contexte pour désigner des distances tampon pour la protection de la faune contre les perturbations humaines.

**Keywords:** conservation buffer, escape distance, flight-initiation distance, human disturbance, setback distance

## Introduction

Optimal escape theory predicts that prey animals should optimise their escape from an approaching predator by initiating escape as soon as the costs of staying

(e.g. being killed or injured) is higher than the costs of escape (e.g. reduced foraging time, energetic costs and lost mating opportunities) (Ydenberg and Dill 1986). A

prey animal reduces these costs by first assessing risk and responding to different levels of risk by varying the distance at which it commences escape when approached by the potential predator (Stankowich and Blumstein 2005; Eason et al. 2006); this distance is called the flight–initiation distance (FID; Ydenberg and Dill 1986). Generally, animals increase FID with increased level of risk and reduce FID with reduced level of risk (Ydenberg and Dill 1986). Consequently, factors that influence a prey animal's perception and assessment of risk should typically influence FID, and so it can be used by wildlife managers to quantify how tolerant animals are to human activities (Burger 1981; Frid and Dill 2002; Rodgers and Schwikert 2002; Pease et al. 2005).

The FID has been widely studied across different animal taxa (Cooper and Blumstein 2015), e.g. insects (Dill and Ydenberg 1987; Nelson and Formanowicz 2005), fish (Januchowski-Hartley et al. 2011), amphibians (Tidwell and Hayes 2013), reptiles (Cooper 2003, 2008), birds (Blumstein 2003; Thiel et al. 2007; Burger et al. 2010) and mammals (Nyahongo 2007; Reimers et al. 2009; Tarakini et al. 2014). It has been applied in several conservation contexts, such as to measure the degree of an animal's fearfulness of humans (Stankowich and Blumstein 2005) and delineate buffer zones around sensitive species, habitats and human activities (Rodgers and Smith 1997; Rodgers and Schwikert 2002; Weston et al. 2012; Guay et al. 2016). It has also been used to indicate hunting pressure on animals (Tarakini et al. 2014; Sreekar et al. 2015) and to protect water birds from being disturbed by boats (Burger et al. 2010; Glover et al. 2011). The FID has also been extensively studied across different continents of the world (Møller et al. 2014, 2016; Livezey et al. 2016). For instance, FID has been collected for as many as 352 bird species and applied to manage human-induced disturbances in Australia (Glover et al. 2011; Weston et al. 2012; Guay et al. 2016). Similarly, several studies focusing on FID have been conducted on European birds across different urbanisation and latitudinal gradients in Europe (Díaz et al. 2013; Samia et al. 2017).

However, there are only a few published studies of FID within an African context (Dill 1990; Nyahongo 2007; Magige et al. 2009; Tarakini et al. 2014) and none that we are aware of in a West African context. This is particularly relevant because of the very rapidly increasing human populations in the region and the close contact of people with animals, even within reserved areas. We thus investigated the factors that might influence FID in several common bird species in the Amurum Forest Reserve in Nigeria, West Africa. Optimal escape theory makes a number of predictions relevant to FID:

- (1) Group size: FID is expected to decrease because individual risk is diluted in larger groups (Foster and Treherne 1981) or because predators are less likely to make a successful hunt on multiple targets – the confusion effect (Neill and Cullen 1974; Milinski 1977). However, FID might also increase as group size increases because probability of predator detection increases with group size (Pulliam 1973; Siegfried and Underhill 1975; Lazarus 1979; Roberts 1996).
- (2) Distance to vegetation cover: if prey animals perceive vegetation cover as a place to seek refuge (Dill

and Houtman 1989; Lima and Dill 1990), the risk of predation is expected to increase with the distance between the approached bird and vegetation used as a refuge (Cooper 2007; Lagos et al. 2009; Clucas and Marzluff 2012). We predicted that FID should increase with increasing distance from vegetation cover.

- (3) Degree of human presence and activities (reserve versus outer farmed areas): animals in sites with high levels of human presence and activities are often reported to become habituated and have lower FIDs (Stankowich and Blumstein 2005; Webb and Blumstein 2005; Møller 2010; Grolle et al. 2014; Tarakini et al. 2014). Likewise, urban birds are known to be more habituated to human activities than are rural ones (Samia et al. 2017). However, it has been reported that in areas where human presence is linked with activities such as hunting, birds respond with longer FIDs (Sreekar et al. 2015). We predicted that FID of birds would be higher in the reserve compared with the surrounding farmed areas, characterised by higher human presence and activities. Although the reserve experiences a level of human exposure because it is used for research and educational activities, disturbance is much lower than surrounding lands. The reserve is effective as a nature haven for wildlife in the area because it is protected against anthropogenic disturbances such as farming, grazing, hunting, logging and indiscriminate burning, and contains several species of birds, plants and mammals, e.g. Tantalus Monkey *Cercopithecus (aethiops) tantalus* and Rock Hyrax *Procavia capensis* (Abalaka 2009; Daru et al. 2015; Ali et al. 2016), that do not occur outside the reserve because of harvesting. In contrast, the area surrounding the reserve is a settlement area, and exposed to uncontrolled human access, activities such as farming, and feral animals.
- (4) Life history: a key component to risk evaluation by birds is life-history strategy, where longer-lived species are expected to take fewer risks than shorter-lived species (Bennett and Owens 2002). We therefore expect differences in basal levels of risk taking between species as a result of differences in life histories between them. For instance, tropical birds differ from temperate birds in several life-history traits. Tropical birds live longer (Peach et al. 2001; Lloyd et al. 2014), have smaller clutch sizes (Ghalambor and Martin 2001; Martin 2004) and attempt first breeding at a relatively older age (Martin 2004). Møller and Liang (2013) showed that tropical birds take smaller risks than closely related temperate taxa in order not to jeopardise their prospects of long-term survival. We thus predicted that species with higher survival rates would have larger FIDs, i.e. take less risk as an optimal strategy to maximise lifetime reproductive success than would species with lower estimates of survival (Martin 1995; Ghalambor and Martin 2001; Martin 2002).

We also tested whether any of the relationships between FID and group size, distance to vegetation cover or levels of human use depend on species by including species in the model as well as the interaction terms (group size\*species, distance to cover\*species, site\*species) in the

analysis. We predict that these relationships are likely to be species specific, hence our inclusion of the interaction terms in the model.

Finally, we considered ‘alert distance’ and ‘starting distance’ and their interaction as confounding variables. Detection distance, the distance at which prey animals detect an approaching predator, is important to the process of making decisions to initiate escape (Lima and Dill 1990; Blumstein 2003). Alert distance, the distance at which an approached animal exhibits alert behaviour in response to the approaching perceived predator, is used as a surrogate measure of detection distance (Stankowich and Coss 2005; Samia et al. 2017). This is because it is difficult to know for sure at what point prey animals detect approaching predators (Ydenberg and Dill 1986). The premise on which alert distance can serve as surrogate for detection distance is the fact that animals are unlikely to respond to stimuli they have not yet become aware of (detected). The fact that an animal shows alert behaviour towards an approaching predator serves as evidence that it has detected the predator, so long as precautions have been taken to ensure that no other external influence, such as other predators or loud sounds, exist during an approach. The limitation of using alert behaviour as a proxy is that it is restricted to animal species that show alert behaviours to approaching predators (Ydenberg and Dill 1986; Cooper 2008). Since all of the focal species in this study exhibited alert behaviours, we have measured and used alert distance as a surrogate for detection distance. FID increases with alert distance because animals that notice the approaching perceived predator at large distances incur costs associated with prolonged monitoring and are less likely to show a default, surprise, instantaneous escape response (Stankowich and Coss 2005; Cooper et al. 2015; Samia et al. 2017). Starting distance is the distance at which a perceived predator begins its approach towards the prey, and this significantly influences FID in many species with few exceptions (Blumstein 2003; Engelhardt and Weladji 2011; Clucas and Marzluff 2012; Tarakini et al. 2014).

## Methods

### Study area

This study was conducted in the Amurum Forest Reserve, Laminga (09°53' N, 08°59' E), Laminga, Jos East, Nigeria, West Africa. The reserve holds one of the few remnants of natural vegetation typical of the Jos plateau. The reserve was established in 2001 as an Important Bird and Biodiversity Area partly because it contains the endemic Jos Plateau Indigobird *Vidua maryae* and its brood-host the Rock Firefinch *Lagonosticta sanguinodorsalis* (Ezealor 2002). It consists of three distinct habitat types: dry savanna, rocky outcrops and gallery forests. Observations were made in the rocky outcrop and savanna habitats within the reserve as well as rocky outcrop and savanna habitats in outer farmed areas surrounding the reserve.

### Bird species

Study species were selected because they occurred in large numbers within and outside the reserve of the study area. All mass measurements of focal birds were extracted

from the ringing database of the AP Leventis Ornithological Research Institute (APLORI) and Borrow and Demey (2008). Species chosen were:

- (1) Rock Firefinch *Lagonosticta sanguinodorsalis*. This species is endemic to the Jos Plateau area of Nigeria. It is a bird of conservation concern belonging to the Estrildidae family (Barshep et al. 2005; Brandt and Cresswell 2008; Abalaka and Hansson 2014). It favours wooded grassland and rocky outcrops, and has a body length of 10 cm on average and a body mass of 10.6 g. It feeds on small grass seeds and occurs singly, in pairs or in small family groups (Fry and Keith 2004; Borrow and Demey 2008).
- (2) Red-cheeked Cordon-bleu *Uraeginthus bengalus*. This bird species is also a member of the Estrildidae family that is widespread and abundant within the northern Guinea and Sudan Savanna agro-ecological zones (Fry and Keith 2004). It favours degraded savanna, gardens as well as farms, and has a body length of 13 cm and mass of 10.1 g (Borrow and Demey 2008). Individuals of this species forage singly, in pairs or in small groups. It feeds on small seeds of annual grasses and termites (Fry and Keith 2004).
- (3) Familiar Chat *Cercomela familiaris*. This species belongs to the Muscicapidae family, it favours rocky areas in savanna, notably over much of Jos Plateau (Elgood et al. 1994). It has an average body length of 14 cm and mass of 17.3 g (Keith et al. 1992; Borrow and Demey 2008). Individuals are often seen singly, in pairs or larger groups, and forage on insects and other invertebrates on the ground.
- (4) Common Bulbul *Pycnonotus barbatus*. This species belongs to the Pycnonotidae family, and is Africa's ubiquitous, quite conspicuous and familiar bird found in almost all habitat types except closed forest and treeless desert (Borrow and Demey 2008). It has a mean body length of 18–20 cm and mass of 38.2 g (Keith et al. 1992). Individuals are usually found in pairs or small groups (Keith et al. 1992). Food includes wild and cultivated fruits, nectar, seeds and insects.
- (5) Laughing Dove *Streptopelia senegalensis*. This species belongs to the Columbidae family. It is abundant and ubiquitous throughout most of sub-Saharan Africa (Urban et al. 1986). It inhabits open woodland and all kinds of human settlements, has an average body length of 24 cm and mass of 93 g (Urban et al. 1986; Borrow and Demey 2008). Individuals often form flocks of 3–4 or more, and feed primarily on seeds and insects (Urban et al. 1986).

The above-mentioned five species, used in the main analysis, belong to two distinct avian orders: Passeriformes (Rock Firefinch, Red-cheeked Cordon-bleu, Familiar Chat, Common Bulbul) and Columbiformes (Laughing Dove). There was no survival estimate found for Laughing Dove in Stevens et al. (2013) or elsewhere in the literature. Hence, to increase the number of species for the survival analysis, FID data were taken from two additional common bird species for which survival estimates were available: sample sizes, however, were not sufficient to include these species in the main analysis. The two additional species mentioned below belong to the avian order Coliiformes (Speckled



Mousebird *Colius striatus*) and Passeriformes (African Thrush *Turdus pelios*).

1. Speckled Mousebird *Colius striatus*. This species belongs to the Coliidae family, it is highly gregarious, occurring usually in small, noisy parties. It is found in various open and wooded habitats, and feeds mainly on leaves and fruits. It has a body length of 30–36 cm and an average body mass of 52.8 g (Borrow and Demey 2008).
2. African Thrush *Turdus pelios*. This species belongs to the Turdidae family. It is a ground-feeding species, occurring solitarily or in small groups, that mostly feeds on plant matter, earthworms and terrestrial arthropods. It has a body length of about 23 cm and mass of 65.2 g on average (Akinpelu and Oyedipe 2004; Borrow and Demey 2008).

### Field methods

Field observations were made using binoculars (8 × 42 mm) to identify focal species between 06:30 and 09:30, and 16:00 and 18:00 from 5 May to 27 June 2014. The observer (BB) walked slowly through existing paths within the study area searching for focal species, as has been done in other FID studies (Blumstein 2003; Cooper 2003; Fernández-Juricic et al. 2006). To eliminate possible effects of observer size and observer bias as well as to avoid variation that might occur through different colours of clothes (Eason et al. 2006), only one observer conducted all trials, wearing the same colour of clothes throughout the survey period. Observations were only made when there was no other person or other confounding factors such as a flying aerial predator or other flying birds between the observer and the focal bird (Pease et al. 2005; Eason et al. 2006; Weston et al. 2012). Approaches were made only on individuals at positions where no obstructing vegetation prevented the observer and focal bird from seeing each other (Fernández-Juricic et al. 2006; Rodriguez-Prieto et al. 2008). Individuals perched above 6 m were excluded, because during a preliminary survey it was discovered that individuals perched at such heights do not usually respond to the observer's approach. Birds that were alert already were excluded regardless of starting distance.

Most approached birds were not individually marked; however, resampling individuals was avoided by focusing on birds in different locations and not resampling the same location more than once per species on each visit. Some individuals of focal species had colour rings on them as part of previous and ongoing studies, and so the colour combinations of the colour-ringed individual were noted during this study. We did not encounter any ringed individual twice during this study, further suggesting that it is unlikely that we repeatedly sampled the same individuals. Furthermore, data collection was alternated daily between the reserve and outer farmed areas outside the reserve (Webb and Blumstein 2005).

On locating an individual of a focal bird species, FID measurements, the predictors (group size, distance to vegetation cover and position in or outside of the reserve), and several potential confounding variables (habitat, species, activity, orientation, social status, position, time of day and perch height) were recorded. Group size was determined by counting the number of individuals within

a 10 m radius of the focal individual (Clucas and Marzluff 2012), before beginning the approach and at the end of each approach. Distance to the nearest vegetation cover (cover was defined as any cluster of trees, shrubs or bushes into which the focal bird can escape) was measured as the distance between the position of the bird and the nearest vegetation cover to it (Cooper 2003); initially by eye to the nearest metre and then checked where necessary by pacing after the response was concluded. Habitat was recorded as either rocky outcrop or savanna, species, as was activity (preening, resting, foraging or calling) and orientation (facing, backing, having side view to the left, and side view to the right) of the focal bird in respect to the observer as well as flock type (social status; either conspecifics or heterospecifics), position of the birds (on the ground, rock, grass, shrub or tree), time of day (morning or evening) and perch height to the nearest 0.2 m (measured using a calibrated range pole). The point at which approach was initiated was marked before the observer commenced the approach at a slow, steady speed of  $\sim 0.5 \text{ ms}^{-1}$  (Blumstein 2003; Fernández-Juricic et al. 2006) while maintaining a direct gaze at the focal bird (Blumstein 2003; Eason et al. 2006; Clucas and Marzluff 2012). The points at which the focal bird detected the observer and began to exhibit alert behaviour, such as increased head-ups, alarm calls, body movements or tail flicks, were marked by the observer by dropping a marker on the ground (Eason et al. 2006). The approach continued until the focal bird escaped. The observer stopped when the bird escaped and measured the FID as the distance between the position of the observer and the position of the bird (Blumstein 2003; Eason et al. 2006; Fernández-Juricic et al. 2006; Bregnballe et al. 2009). Starting distance was measured as the distance between the first marked point, where the observer initiated the approach and the position of the focal individual (Blumstein 2003). Alert distance was measured as the distance between the position of the bird and the second marked point, at which the bird detected the observer and began to exhibit alert behaviour (Fernández-Juricic et al. 2001; Eason et al. 2006).

All distance measurements were taken in metres using a laser rangefinder (Nikon Laser 800; Nikon, Tokyo, Japan with measuring range of 10–730 m) by pointing the laser beam at the position (point) from which the focal bird escaped (Fernández-Juricic et al. 2001; Clucas and Marzluff 2012). Given that the laser rangefinder used could not take distance measurements of less than 10 m, all such distances were measured directly with a metre-length stick.

### Data analysis

All data collected were analysed using general linear mixed models with a normal error distribution in R statistical package version 3.0.2 (2013-09-25) using the lme4 and lmerTest libraries (R Development Core Team 2013). An individual-based approach was used, using multiple measurements from a species and then controlling for the repeated measures using a mixed model. Starting distance, distance to cover, group size, perch height and alert distance were entered as covariates; while species (five levels), habitat type (two levels), site (two levels), time of day (two levels), flock type (two levels), position

(six levels), activity (four levels) and orientation (four levels) were entered as categorical variables alongside all the two-way interactions among the above-mentioned explanatory variables. Random effects of species nested in family nested in order were included to control for phylogenetic effects. Following examples of Thiel et al. (2007), Reimers et al. (2009), Clucas and Marzluff (2012) and Tarakini et al. (2014), variables were selected using backward elimination as recommended in Crawley (2005) by starting with full models containing all biologically plausible main effects and interaction terms. Variables with the highest  $p$ -values were removed starting with the non-significant interactions and the procedure repeated until the final model was attained. The final model was selected using Akaike's information criterion (AIC), as the model with the lowest AIC value, and with all terms retained being statistically significant. Final models were checked by reinsertion of all variables removed, singly back into the model. All the variables reintroduced back into the final model remained non-significant and the variables that were significant in the final model remained significant. We present both starting models and final models to demonstrate that model simplification did not change the overall biological significance of the results. The FID measurements were square-root transformed prior to the analysis to improve the normal distribution of the residuals of the model: final models were visually checked for not significantly violating assumptions according to criteria in Crawley (2005). All FID data were back-transformed when illustrating predicted results from the final model. Interactions between species and the predictors were also added to the model.

A further species-based general linear model assuming a normal distribution was tested to investigate the relationship between survival and FID. This model only contained six data points (one for each species) and so was deliberately kept simply as  $FID \sim \text{survival}$ . This was an *a priori* hypothesis of interest and limited power prevented the addition of other variables to the model. Note also that limited power prevented phylogenetic control within the analysis and so any trends revealed may largely reflect traits within Passerines, because five of the six species belonged to this order. Mean values of variables are presented as the mean  $\pm$  standard error.

## Results

At the end of the survey, a total of 465 approaches to focal birds were recorded for use in the analyses. Of these, 133 approaches were conducted on Common Bulbul, 122 were on Red-cheeked Cordon-bleu, 92 on Familiar Chat, 77 on Laughing Dove, 41 on Rock Firefinch, 20 on African Thrush and 19 for Speckled Mousebird. Responses to 193 approaches were recorded inside the reserve and 272 recorded outside the reserve. Approaches were initiated at distances ranging from 15 to 94 m ( $32.0 \pm 0.5$  m). Focal individuals exhibited alert behaviour at distances ranging from 5 to 54 m ( $18.3 \pm 0.3$  m) while FID ranged from 2 to 41 m ( $14.2 \pm 0.3$  m). Individuals of focal species approached occurred in groups ranging from a single individual to 22 individuals ( $2.5 \pm 0.1$ ) and on average focal species were

$4.8 \pm 0.3$  m away from cover. Species-specific summaries are included in Appendix 1.

The model with the lowest AIC value, hence the best model, retained time of day, site, habitat type, species, position, group size, starting distance, alert distance, distance to cover and the interaction term starting distance\*alert distance (Table 1). The FID significantly decreased with increasing group size in all species (Table 1, Figure 1). Meanwhile, FID increased significantly with increasing distance to cover (Table 1, Figure 2). The FID was significantly higher in the reserve than outside (Table 1). The FID varied little depending on taxonomy (i.e. between species and families); marginal  $R^2$  for the final model in Table 1 was 0.64, and overall  $R^2$  accounting for phylogeny was 0.65 (Table 1, Appendix 2), i.e. only 0.01 of the total variance was accounted for by taxonomic differences. The relationship between FID and group size, FID and distance to cover as well as FID by sites did not differ by species because the interaction terms species\*group size, species\*distance to cover and species\*site were not significant in any cases (Table 1). Activity, orientation, flock type and perch height were not significantly related to FID in this study.

The model testing the relationship between survival estimates and FID indicated that FID in focal species significantly increased with survival ( $F_{1,4} = 13.4$ ,  $p < 0.02$ ,  $B = 23.47$ , adj.  $R^2 = 0.71$ ; or Spearman's rank correlation coefficient  $\rho = 0.89$ ,  $p = 0.033$ ; Figure 3); see Appendix 2 for species-specific survival and FID values used.

## Discussion

The present results generally confirmed all our predictions, although there was little variation found across the species we sampled. The FID decreased with increasing group size, confirming that FID is likely to decrease because individual risk is diluted in larger groups and the fact that predators are less likely to make a successful hunt on multiple targets due to the confusion effect (Neill and Cullen 1974; Milinski 1977). Our finding, therefore, agrees with Foster and Treherne (1981) and Pulliam (1973) that individuals in a group may benefit through risk dilution, and so perceive a lower risk when in larger groups. However, positive relationships have been reported between FID and group size in Dark-bellied Brent Geese *Branta bernicla bernicla* (Owens 1977), shorebirds (Glover et al. 2011) and fish (Januchowski-Hartley et al. 2011).

We found a positive linear relationship between FID and distance to cover, indicating that focal birds perceived higher levels of threat with increasing distance from refuge (Dear et al. 2015; Monclús et al. 2015). Refuge could vary depending on species involved; for example, some birds use water as refuge and thus have longer FIDs when further from water (Guay et al. 2013; Dear et al. 2015). For animals that utilise vegetation cover during escape from their predators, risk increases with distance to vegetation cover (Dill and Houtman 1989) and so animals far from vegetation cover usually have longer FIDs than those closer to it (Ydenberg and Dill 1986). This finding is consistent across many different taxa, e.g. reptiles (Cooper 2007), birds (Clucas and Marzluff 2012) and mammals (Dill


**Table 1:** Relationship between square-root transformed flight-initiation distance and explanatory variables in five bird species within Amurum Forest Reserve and outside the reserve. The full model (marginal  $R^2 = 0.52$ , overall  $R^2 = 0.74$ ) and the model with the lowest AIC value (final model) (marginal  $R^2 = 0.64$ , overall  $R^2 = 0.65$ ) are shown. Random effects of species nested in family nested in order were included to control for phylogenetic effects. Reserve and Common Bulbul take the value of the intercept. Significant  $p$ -values and variables are in bold

Explanatory variable	Full model					Final model				
	Est.	SE	df	$t$	$p$	Est.	SE	df	$T$	$P$
(Intercept)	1.370	0.495	427	2.8	<0.001	1.258	0.198	191	6.4	<0.001
Time of day morning	-0.059	0.046	427	-1.3	0.208	—	—	—	—	—
Outside reserve	-0.121	0.089	427	-1.4	0.177	-0.115	0.048	404	-2.4	<b>0.018</b>
Habitat type savanna	-0.074	0.051	427	-1.4	0.151	—	—	—	—	—
Position ground	-0.080	0.119	427	-0.7	0.504	—	—	—	—	—
Position rock	-0.027	0.110	427	-0.2	0.803	—	—	—	—	—
Position shrub	0.001	0.081	427	0.0	0.991	—	—	—	—	—
Position thicket	0.118	0.112	427	1.0	0.296	—	—	—	—	—
Position tree	-0.107	0.093	427	-1.1	0.254	—	—	—	—	—
Activity foraging	0.101	0.104	427	1.0	0.335	—	—	—	—	—
Activity preening	0.135	0.112	427	1.2	0.228	—	—	—	—	—
Activity resting	0.127	0.082	427	1.5	0.125	—	—	—	—	—
Orientation facing	-0.019	0.060	427	-0.3	0.754	—	—	—	—	—
Orientation side-view-left	-0.013	0.080	427	-0.2	0.869	—	—	—	—	—
Orientation side-view-right	0.053	0.068	427	0.8	0.439	—	—	—	—	—
Flock type	0.029	0.062	427	0.5	0.645	—	—	—	—	—
Group size	-0.016	0.016	427	-1.0	0.321	-0.031	0.012	457	-2.7	<b>0.008</b>
Starting distance	0.020	0.006	427	3.4	<b>0.001</b>	0.019	0.006	449	3.5	<0.001
Alert distance	0.147	0.010	427	13.4	<0.001	0.147	0.010	434	14.3	<0.001
Distance to cover	0.010	0.007	427	1.4	0.151	0.014	0.004	458	3.5	<0.001
Perch height	-0.012	0.020	427	-0.6	0.551	—	—	—	—	—
Familiar Chat	-0.140	0.606	0	-0.2	1.000	—	—	—	—	—
Laughing Dove	-0.176	0.618	0	-0.3	1.000	—	—	—	—	—
Red-cheeked Cordon-bleu	-0.075	0.608	0	-0.1	1.000	—	—	—	—	—
Rock Firefinch	-0.151	0.617	0	-0.2	1.000	—	—	—	—	—
Familiar chat:Group size	-0.049	0.039	427	-1.3	0.211	—	—	—	—	—
Laughing Dove:Group size	-0.043	0.032	427	-1.3	0.184	—	—	—	—	—
Red-cheeked Cordon-bleu:Group size	-0.055	0.031	427	-1.7	0.086	—	—	—	—	—
Rock Firefinch:Group size	-0.052	0.061	427	-0.8	0.401	—	—	—	—	—
Familiar Chat:Distance to cover	0.015	0.015	427	1.0	0.334	—	—	—	—	—
Laughing Dove:Distance to cover	0.017	0.011	427	1.5	0.123	—	—	—	—	—
Red-cheeked Cordon-bleu:Distance to cover	0.003	0.010	427	0.3	0.752	—	—	—	—	—
Rock Firefinch:Distance to cover	0.018	0.032	427	0.6	0.574	—	—	—	—	—
Outside:Familiar Chat	-0.053	0.154	427	-0.3	0.729	—	—	—	—	—
Outside:Laughing Dove	-0.040	0.154	427	-0.3	0.798	—	—	—	—	—
Outside:RC Cordon-bleu	0.067	0.148	427	0.5	0.649	—	—	—	—	—
Outside:Rock Firefinch	0.093	0.198	427	0.5	0.641	—	—	—	—	—
Starting distance:Alert distance	-0.001	0.0003	427	-5.1	<0.001	0.001	0.0002	455	-5.3	<0.001


and Houtman 1989; Lagos et al. 2009). However, vegetation might have inhibitory effects on visibility and thus may prevent accurate processing of visual information from the environment, thereby decreasing the probability of early detection (Javůrková et al. 2012). Furthermore, predation risk can increase with proximity to cover for prey animals whose predators attack from cover rather than open (Cresswell 1994). The relationship between FID and distance to cover may thus be confounded by these factors.

We recorded a significantly higher FID in areas with lower human presence (reserve) compared with areas characterised by higher human presence (outside the reserve). This suggests at least to some degree a higher level of habituation in birds that occur outside the reserve of the study area. Habituation and experience with humans significantly influences the escape response of animals during animal-human interactions (Stankowich and Blumstein

2005). Generally, animals are able to habituate to the level of human presence in two ways, depending on the behaviour of humans toward them (Cooper and Blumstein 2013). In areas where human presence is frequent and non-threatening, animals lose their fear of humans and respond with shorter FIDs (Stankowich and Blumstein 2005; Webb and Blumstein 2005; Grolle et al. 2014; Tarakini et al. 2014; Samia et al. 2017). On the other hand, in areas where human presence is associated with threatening or discouraging behaviour such as hunting, animals respond with exaggerated FIDs (Januchowski-Hartley et al. 2011; Clucas and Marzluff 2012; Sreekar et al. 2015). In certain cases, habituation to humans can cause problems through also affecting habituation to natural predators and because of dietary shifts as animals feed in close proximity to humans (Hines 2011; Geffroy et al. 2015). Given that the focal species observed in this study are not under any




**Figure 1:** Predicted relationship between flight-initiation distance and group size plotted from the final model. Shaded area indicates one standard error. ○ Common Bulbul, △ Familiar Chat, + Laughing Dove, × Red-cheeked Cordon-bleu, ◇ Rock Firefinch



**Figure 2:** Predicted relationship between flight-initiation distance and distance to cover plotted from the final model. Shaded area indicates one standard error. ○ Common Bulbul, △ Familiar Chat, + Laughing Dove, × Red-cheeked Cordon-bleu, ◇ Rock Firefinch

hunting pressure in the study sites, the significantly lowered mean FID in the settlement area outside the reserve compared with that observed inside the reserve may perhaps be attributed to habituation of focal bird species to non-lethal, repeated exposure to humans.

We found a strong positive linear correlation between mean FID and survival estimates across the focal bird species we sampled. Although our sample size is small, it was nonetheless a strong relationship, confirming our prediction that species with higher survival rates (and/or life-history traits that correlate with this) would have larger FIDs. Species that initiate flight early when approached by a predator are assumed to take less risk, whereas those



**Figure 3:** Predicted relationship between mean flight-initiation distance of six bird species and mean adult survival estimated for each species

that delay escape are held to take more risk (Møller and Liang 2013). Life-history theory proposes that longer-lived species take fewer risks compared with shorter-lived species in order to optimise their overall fitness (Martin 1995; Ghalambor and Martin 2001; Bennett and Owens 2002; Martin 2002). This then implies that longer-lived species increase their probability of future breeding by taking less risk, whereas shorter-lived species take more risk to boost their current reproductive output. Our focal species differed in their mean FID, indicating that they associated different levels of risk to the observer. This could possibly be attributed to differences in ecology and life-history traits between them. For example, variation in size among species have been shown to drive avian interspecific differences in FID and body size is one of the best predictors of survival rate (Blumstein et al. 2005; Fernández-Juricic et al. 2006; Bregnballe et al. 2009). Larger species are known to have higher visual acuity (Brooke et al. 1999; Kiltie 2000) and higher wing loading (Burns and Ydenberg 2002), both of which influence flight manoeuvrability and possibly escape strategy (Fernández-Juricic et al. 2001). These imply that, under comparable risk of predation, the benefits of early escape will be higher for larger species than for smaller species. Furthermore, differences in other ecology and life-history traits (Bregnballe et al. 2009), such as foraging habits (Blumstein et al. 2005), energy requirements (Blumstein et al. 2005), escape strategy (Cresswell 1993), levels of exposure and so habituation to humans (Burger 1981), are the probable explanations for the interspecific variation in FID among bird species.

Although there were species differences in the absolute levels of FID as stated above, we found no strong evidence for any of the relationships between FID and group size or distance to cover to depend on species, nor for the difference in FID between the reserve and outside of the reserve to depend on species. This suggests that, for at least the limited range of species we tested, that FID is broadly affected in the same way by group size, proximity to vegetation cover acting as a refuge and habituation to humans. Furthermore, this suggests the generality



of the FID approach, despite ecological and life-history differences, at least for the species we sampled.

Other results with respect to the confounding variables are also consistent with predictions of optimal escape theory. Starting distance was related to FID consistent with the findings of Blumstein (2003) and confirms that starting distances should be included in FID studies as a covariate. The FID increased with alert distance consistent with the many other studies that have reported strong positive correlations between FID and alert distance (Stankowich and Coss 2005; Eason et al. 2006; Lagos et al. 2009), probably because prey animals that detect approaching predators at greater distances may initiate escape at longer distances to minimise the costs associated with monitoring. There was also evidence that the relationship between FID and starting distance depends on alert distance. This can be best explained by incorporating the perceptual limit hypothesis, which states that escape from an approaching predator is constrained by the prey animal's ability to detect predators and assumes that prey animals flee as soon as predators are detected (Ydenberg and Dill 1986), and the economic escape theory, which predicts that individuals do not usually flee as soon as potential predators are detected, but delay their decision until the costs of staying outweighs the benefit of staying (Ydenberg and Dill 1986). Both hypotheses may be true depending on the distance at which the approached individual initially detects the approaching predator.

In conclusion, this study provides the first empirical data for testing the predictions of optimal escape theory in West Africa. Overall, the focal birds perceived humans as a potential threat and responded in accordance to the predictions of optimal escape theory. We show that bird species differ in FID, probably because of life-history traits and their ecology. Consequently, some species are expected to be less able to deal with human disturbance (that does not actually have immediate fitness costs) than others and so identification of these through FID on a case-by-case basis will allow reserve managers to manage human access and viewing of wildlife in nature reserves. For instance, the endemic Rock Firefinch that showed greater sensitivity will require more management attention than would the Red-cheeked Cordon-bleu. Safe buffer zones may be built using the highest mean FID so the needs of the less-demanding species, and likely species where FID is unknown, are also met. However, care must be taken to avoid excessively long buffers that threaten birdwatching, educational activities and research. Hence, alternatives such as formed path buffers around human access routes can be used to manage disturbance. Furthermore, larger buffers may be built around certain sensitive habitats for the conservation of birds with special habitat needs.

**Acknowledgements** — The authors would like to thank the AP Leventis Ornithological Research Institute (APLORI) for help with logistics. We acknowledge field visits made by Manu Shiiwua and Ulf Ottosson. We also acknowledge the SCCS Miriam Rothschild Travel Bursary Programme for travel funds during the period of writing this paper. Thanks to the Leventis Conservation Foundation for providing full funding for this study. The study was carried out in Nigeria where no licences are required for the procedures used. Nevertheless, this study was carried out under

the ethical guidelines of the APLORI Scientific Committee (APLORI is the only ornithological research institute in Nigeria) based on the Association for the Study of Animal Behaviour guidelines. This paper is number 125 published by the AP Leventis Ornithological Research Institute.

## References

- Abalaka J. 2009. How important are figs in the conservation effort at Amurum forest reserve, Laminga, Nigeria. Unpublished report to the Rufford Small Grants Foundation. Available at [http://www.rufford.org/files/32.12.07 Detailed Final Report.pdf](http://www.rufford.org/files/32.12.07%20Detailed%20Final%20Report.pdf).
- Abalaka J, Hansson B. 2014. Microsatellite markers for population genetic studies of the Rock Firefinch, *Lagonosticta sanguinodorsalis*. *African Zoology* 49: 301–306.
- Akinpelu AI, Oyedipe OA. 2004. A twelve-month field study of the West African Thrush *Turdus pelios* (Passeriformes: Muscicapidae). Part 1: food and feeding ecology. *Revista de biología tropical* 52: 1001–1007.
- Ali AD, Elisha EB, Abiem I, Habila S, Okeke OM. 2016. Hygrophytes and wetland angiospermic macrophyte in gallery forest of Amurum forest reserve, Jos, Plateau. *Research in Plant Sciences* 4: 10–15.
- Barshap Y, Manu S, Brandt M. 2005. Resource partitioning in the Rock Firefinch *Lagonosticta sanguinodorsalis* and the Red-billed Firefinch *Lagonosticta senegala*. *Journal of Ornithology* 29: 5711–5725.
- Bennett PM, Owens IPF. 2002. *Evolutionary ecology of birds: life histories, mating systems, and extinction*. Oxford: Oxford University Press.
- Blumstein DT. 2003. Flight-initiation distance in birds is dependent on intruder starting distance. *Journal of Wildlife Management* 67: 852–857.
- Blumstein DT, Fernández-Juricic E, Zollner PA, Garity SC. 2005. Inter-specific variation in avian responses to human disturbance. *Journal of Applied Ecology* 42: 943–953.
- Borrow N, Demey R. 2008. *Birds of western Africa*. London: Christopher Helm.
- Brandt MJ, Cresswell W. 2008. Breeding behaviour, home range and habitat selection in Rock Firefinches *Lagonosticta sanguinodorsalis* in the wet and dry season in central Nigeria. *Ibis* 150: 495–507.
- Bregnballe T, Aaen KIM, Fox AD. 2009. Escape distances from human pedestrians by staging waterbirds in a Danish wetland. *Wildfowl Special Issue* 2: 115–130.
- Brooke MDL, Hanley S, Laughlin SB. 1999. The scaling of eye size with body mass in birds. *Proceedings of the Royal Society B* 266: 405–412.
- Burger J. 1981. The effect of human activity on birds at a coastal bay. *Biological Conservation* 21: 231–241.
- Burger J, Gochfeld M, Jenkins CD, Lesser F. 2010. Effect of approaching boats on nesting Black Skimmers: using response distances to establish protective buffer zones. *Journal of Wildlife Management* 74: 102–108.
- Burns JG, Ydenberg RC. 2002. The effects of wing loading and gender on the escape flights of Least Sandpipers (*Calidris minutilla*) and Western Sandpipers (*Calidris mauri*). *Behavioral Ecology and Sociobiology* 52: 128–136.
- Clucas B, Marzluff JM. 2012. Attitudes and actions toward birds in urban areas: human cultural differences influence bird behavior. *The Auk* 129: 8–16.
- Cooper WE Jr. 2003. Risk factors affecting escape behavior by the Desert Iguana, *Dipsosaurus dorsalis*: speed and directness of predator approach, degree of cover, direction of turning by a predator, and temperature. *Canadian Journal of Zoology* 81: 979–984.
- Cooper WE Jr. 2007. Escape and its relationship to pursuit-deterrent

- signalling in the Cuban curly-tailed lizard, *Leiocephalus carinatus*. *Herpetologica* 63: 144–150.
- Cooper WE Jr. 2008. Strong artifactual effect of starting distance on flight initiation distance in the actively foraging lizard *Aspidooscelis exsanguis*. *Herpetologica* 64: 200–206.
- Cooper WE Jr, Blumstein DT. 2013. Novel effects of monitoring predators on costs of fleeing and not fleeing explain flushing early in economic escape theory. *Behavioral Ecology* 25: 44–52.
- Cooper WE Jr, Blumstein DT. 2015. *Escaping from predators: an integrative view of escape decisions*. Cambridge: Cambridge University Press.
- Cooper WE Jr, Samia DSM, Blumstein DT. 2015. Fear, spontaneity, and artifact in economic escape theory: a review and prospectus. *Advances in the Study of Behavior* 47: 147–179.
- Crawley MJ. 2005. *Statistics: an introduction using R*. Chichester: John Wiley and Sons.
- Cresswell W. 1993. Escape responses by Redshanks, *Tringa totanus*, on attack by avian predators. *Animal Behaviour* 46: 609–11.
- Cresswell W. 1994. Age-dependent choice of Redshank (*Tringa totanus*) feeding location: profitability or risk? *Journal of Animal Ecology* 63: 589–600.
- Daru BH, Yessoufou K, Nuttman C, Abalaka J. 2015. A preliminary study of bird use of fig *Ficus* species in Amurum forest reserve, Nigeria. *Malimbus* 37: 1–15.
- Dear EJ, Guay P-J, Robinson RW, Weston MA. 2015. Distance from shore positively influences alert distance in three wetland bird species. *Wetlands Ecology and Management* 23: 315–318.
- Diaz M, Möller AP, Flensted-Jensen E, Grim T, Ibáñez-Álamo JD, Jokimäki J, Markó G, Tryjanowski P. 2013. The geography of fear: a latitudinal gradient in anti-predator escape distances of birds across Europe. *PLoS ONE* 8: e64634.
- Dill LM. 1990. Distance-to-cover and the escape decisions of an African cichlid fish, *Melunochromis chipokae*. *Environmental Biology of Fishes* 27: 147–152.
- Dill LM, Houtman R. 1989. The influence of distance to refuge on flight initiation distance in the Gray Squirrel. *Canadian Journal of Zoology* 67: 233–235.
- Dill LM, Ydenberg RC. 1987. The group size – flight distance relationship in water striders (*Gerris remigis*). *Canadian Journal of Zoology* 65: 223–226.
- Eason PK, Sherman PT, Rankin O, Coleman B. 2006. Factors affecting flight initiation distance in American robins. *Journal of Wildlife Management* 70: 1796–1800.
- Elgood J, Heigham JB, Moore AM, Sharland RE, Skinner NJ. 1994. *The birds of Nigeria* (2nd edn). BOU Checklist no. 4. Tring: British Ornithologists' Union.
- Engelhardt SC, Weladji RB. 2011. Effects of levels of human exposure on flight initiation distance and distance to refuge in foraging eastern gray squirrels (*Sciurus carolinensis*). *Canadian Journal of Zoology* 89: 823–830.
- Ezealor AU. 2002. *Critical sites for biodiversity conservation in Nigeria*. Lagos: Nigerian Conservation Foundation.
- Fernández-Juricic E, Blumstein DT, Abrica G, Manriquez L, Adams LB, Adams R, Daneshrad M, Rodriguez-Prieto I. 2006. Relationships of anti-predator escape and post-escape responses with body mass and morphology: a comparative avian study. *Evolutionary Ecology Research* 8: 731–752.
- Fernández-Juricic E, Jimenez MD, Lucas E. 2001. Alert distance as an alternative measure of bird tolerance to human disturbance: implications for park design. *Environmental Conservation* 28: 263–269.
- Foster WA, Treherne JE. 1981. Evidence for the dilution effect in the selfish herd from fish predation on a marine insect. *Nature* 293: 466–467.
- Frid A, Dill L. 2002. Human-caused disturbance stimuli as a form of predation risk. *Conservation Ecology* 6: 11–27.
- Fry CH, Keith S. 2004. *The birds of Africa*, vol. 7. London: Academic Press.
- Geffroy B, Samia DSM, Bessa E, Blumstein DT. 2015. How nature-based tourism might increase prey vulnerability to predators. *Trends in Ecology and Evolution* 30: 755–765.
- Ghalambor CK, Martin TE. 2001. Fecundity-survival trade-offs and parental risk-taking in birds. *Science* 292: 494–497.
- Glover HK, Weston MA, Maguire GS, Miller KK, Christie BA. 2011. Towards ecologically meaningful and socially acceptable buffers: response distances of shorebirds in Victoria, Australia, to human disturbance. *Landscape and Urban Planning* 103: 326–334.
- Grolle EK, Lopez MC, Marina MG. 2014. Flight initiation distance differs between populations of Western fence lizards (*Sceloporus occidentalis*) at a rural and an urban site. *Bulletin of the Southern California Academy of Science* 113: 42–46.
- Guay P-J, van Dongen WFD, Robinson RW, Blumstein DT, Weston MA. 2016. AvianBuffer: an interactive tool for characterising and managing wildlife fear responses. *Ambio* 45: 841–851.
- Guay P-J, Lorenz RDA, Robinson RW, Symonds MRE, Weston MA. 2013. Distance from water, sex and approach direction influence flight distances among habituated Black Swans. *Ethology* 119: 552–558.
- Hines KN. 2011. Effects of ecotourism on endangered Northern Bahamian Rock Iguanas (*Cyclura cychlura*). *Herpetological Conservation and Biology* 6: 250–259.
- Januchowski-Hartley FA, Graham NAJ, Feary DA, Morove T, Cinner J. 2011. Fear of fishers: human predation explains behavioral changes in coral reef fishes. *PLoS ONE* 6: e22761.
- Javůrková V, Šizling AL, Kreisinger J, Albrecht T. 2012. An alternative theoretical approach to escape decision-making: The role of visual cues. *PLoS ONE* 7: e32522.
- Keith S, Urban EK, Fry CH. 1992. *The birds of Africa*, vol. 4. London: Academic Press.
- Kiltie RA. 2000. Scaling of visual acuity with body size. *Functional Ecology* 14: 226–234.
- Lagos P, Meier A, Tolhuysen LO, Castro R, Bozinovic F, Ebersperger L. 2009. Flight initiation distance is differentially sensitive to the costs of staying and leaving food patches in a small-mammal prey. *Canadian Journal of Zoology* 87: 1016–1023.
- Lazarus J. 1979. The early warning function of flocking in birds: an experimental study with captive quelea. *Animal Behaviour* 27: 855–865.
- Lima SL, Dill LM. 1990. Behavioral decisions made under the risk of predation: a review and prospectus. *Canadian Journal of Zoology* 68: 619–640.
- Livezey KB, Fernández-Juricic E, Blumstein DT. 2016. Database of bird flight initiation distances to assist in estimating effects from human disturbance and delineating buffer areas. *Journal of Fish and Wildlife Management* 7: 181–191.
- Lloyd P, Abadi F, Altwegg R, Martin TE. 2014. South temperate birds have higher apparent adult survival than tropical birds in Africa. *Journal of Avian Biology* 45: 493–500.
- Magige FJ, Holmern T, Stokke S, Mlingwa C, Røskoft E. 2009. Does illegal hunting affect density and behaviour of African grassland birds? A case study on Ostrich (*Struthio camelus*). *Biodiversity and Conservation* 18: 1361–1373.
- Martin TE. 1995. Avian life history evolution in relation to nest sites, nest predation, and food. *Ecological Monographs* 65: 101–127.
- Martin TE. 2002. A new view of avian life-history evolution tested on an incubation paradox. *Proceedings of the Royal Society B* 269: 309–316.
- Martin TE. 2004. Avian life-history evolution has an eminent past: does it have a bright future? *The Auk* 121: 289–301.
- Milinski M. 1977. Do all members of a swarm suffer the same predation? *Zeitschrift für Tierpsychologie* 43: 311–325.
- Møller AP. 2010. Interspecific variation in fear responses predicts urbanization in birds. *Behavioural Ecology* 21: 365–371.

- Møller AP, Liang W. 2013. Tropical birds take small risks. *Behavioral Ecology* 24: 267–272.
- Møller AP, Samia DSM, Weston MA, Guay P-J, Blumstein DT. 2014. American exceptionalism: population trends and flight initiation distances in birds from three continents. *PLoS ONE* 9: e107883.
- Møller AP, Samia DSM, Weston MA, Guay PJ, Blumstein DT. 2016. Flight initiation distances in relation to sexual dichromatism and body size in birds from three continents. *Biological Journal of the Linnean Society* 117: 823–831.
- Monclús R, Anderson AM, Blumstein DT. 2015. Do Yellow-bellied Marmots perceive enhanced predation risk when they are farther from safety? An experimental study. *Ethology* 121: 831–839.
- Neill SRJ, Cullen JM. 1974. Experiments on whether schooling by their prey affects the hunting behaviour of cephalopods and fish predators. *Journal of Zoology (London)* 172: 549–569.
- Nelson MK, Formanowicz DR. 2005. Relationship between escape speed and flight distance in a wolf spider, *Hogna carolinensis* (Walckenaer 1805). *Journal of Arachnology* 33: 153–158.
- Nyahongo JW. 2007. Flight initiation distances of five herbivores to approaches by vehicles in the Serengeti National Park, Tanzania. *African Journal of Ecology* 46: 227–229.
- Owens N. 1977. Responses of wintering Brent Geese to human disturbance. *Wildfowl* 28: 5–14.
- Peach WJ, Hanmer DB, Oatley TB. 2001. Do southern African songbirds live longer than their European counterparts? *Oikos* 93: 235–249.
- Pease ML, Rose RK, Butler MJ. 2005. Effects of human disturbances on the behavior of wintering ducks. *Wildlife Society Bulletin* 33: 103–112.
- Pulliam HR. 1973. On the advantages of flocking. *Journal of Theoretical Biology* 38: 419–422.
- R Development Core Team. 2013. R: a language and environment for statistical computing. Vienna: R Foundation for Statistical Computing.
- Reimers E, Loe LE, Eftestøl S, Colman JE, Dahle B. 2009. Effects of hunting on response behaviors of Wild Reindeer. *Journal of Wildlife Management* 73: 844–851.
- Roberts G. 1996. Why individual vigilance declines as group size increases. *Animal Behaviour* 51: 1077–1086.
- Rodgers JA, Schwikert ST. 2002. Buffer-zone distances to protect foraging and loafing waterbirds from disturbance by personal watercraft and outboard-powered boats. *Conservation Biology* 16: 216–224.
- Rodgers JA, Smith HT. 1997. Buffer zone distances to protect foraging and loafing waterbirds from human disturbance in Florida. *Wildlife Society Bulletin* 25: 139–145.
- Rodriguez-Prieto I, Fernández-Juricic E, Martín J. 2008. To run or to fly: low cost versus low risk escape strategies in blackbirds. *Behaviour* 145: 1125–1138.
- Samia DSM, Blumstein DT, Díaz M, Grim T, Ibáñez-Álamo JD, Jokimäki J, Tätté K, Markó G, Tryjanowski P, Møller AP. 2017. Rural-urban differences in escape behavior of European birds across a latitudinal gradient. *Frontiers in Ecology and Evolution* 5: 66.
- Siegfried WR, Underhill LG. 1975. Strategy in doves. *Animal Behaviour* 23: 504–508.
- Sreekar R, Goodale E, Harrison RD. 2015. Flight initiation distance as behavioral indicator of hunting pressure: a case study of the Sooty-headed Bulbul (*Pycnonotus aurigaster*) in Xishuangbanna, SW China. *Tropical Conservation Science* 8: 505–512.
- Stankowich T, Blumstein DT. 2005. Fear in animals: a meta-analysis and review of risk assessment. *Proceedings of the Royal Society B* 272: 2627–2634.
- Stankowich T, Coss RG. 2005. Effects of predator behavior and proximity on risk assessment by Columbian black-tailed deer. *Behavioural Ecology* 17: 246–254.
- Stevens MC, Ottoson U, McGregor R, Brandt M, Cresswell W. 2013. Survival rates in West African savanna birds. *Ostrich* 84: 11–25.
- Tarakini T, Crosmay W, Fritz H, Mundy P. 2014. Flight behavioural responses to sport hunting by two African herbivores. *South African Journal of Wildlife Research* 44: 76–83.
- Thiel D, Ménoni E, Brenot J, Jenni L. 2007. Effects of recreation and hunting on flushing distance of Capercaillie. *Journal of Wildlife Management* 71: 1784–1792.
- Tidwell KS, Hayes MP. 2013. Difference in flight initiation distance between recently metamorphosed Oregon spotted frogs (*Rana pretiosa*) and American bullfrogs (*Lithobates catesbeianus*). *Herpetological Conservation and Biology* 8: 426–434.
- Urban EK, Fry CH, Keith S. 1986. *The birds of Africa*, vol. 2. London: Academic Press.
- Webb N V, Blumstein DT. 2005. Variation in human disturbance differentially affects predation risk assessment in Western Gulls. *The Condor* 107: 178–181.
- Weston MA, McLeod E m., Blumstein DT, Guay P. 2012. A review of flight initiation distances and their application to managing disturbance to Australian birds. *Emu* 112: 269–286.
- Ydenberg RC, Dill LM. 1986. The economics of fleeing from predators. *Advances in the Study of Behavior* 16: 229–249.

**Appendix 1:** Species-specific summaries of main variable for the five species used in the main flight-initiation distance analysis

Variable	Common Bulbul (N = 133)		Familiar Chat (N = 92)		Laughing Dove (N = 77)		Red-cheeked Cordon-bleu (N = 122)		Rock Firefinch (N = 41)	
	Mean	Range	Mean	Range	Mean	Range	Mean	Range	Mean	Range
<b>Continuous variables</b>										
FID	17 ± 0.5	5–41	15 ± 0.7	5–34	15 ± 0.7	531.5	10 ± 0.4	2–25	12 ± 0.7	5–26
Alert distance	21 ± 0.5	6–45	20 ± 0.7	7–36	21 ± 0.8	7.5–54	14 ± 0.4	5–25	16 ± 0.9	6–29.5
Starting distance	35 ± 0.9	20–71	33 ± 1.2	16–94	36 ± 1.3	20.5–75	26 ± 0.7	15–55	27 ± 1.2	16–46
Group size	3 ± 0.3	1–22	2 ± 0.16	1–7	3 ± 0.2	1–12	2 ± 0.2	1–10	2 ± 0.2	1–6
Distance to cover	5 ± 0.6	0–32	5 ± 0.4	0–19	7 ± 0.7	0–29	4 ± 0.5	0–38	3 ± 0.4	0–8
Perch height	3 ± 0.2	0–6	5 ± 0.2	0–6	2 ± 0.2	0–6	1 ± 0.1	0–6	1 ± 2.1	0–5
<b>Categorical variables</b>										
Time section	Morning N = 60	Evening N = 73	Morning N = 34	Evening N = 58	Morning N = 31	Evening N = 46	Morning N = 52	Evening N = 70	Morning N = 15	Evening N = 26
Habitat	Rock N = 49	Savanna N = 84	Rock N = 76	Savanna N = 16	Rock N = 46	Savanna N = 31	Rock N = 47	Savanna N = 75	Rock N = 35	Savanna N = 6
Site	Reserve N = 48	Outside N = 85	Reserve N = 73	Outside N = 19	Reserve N = 21	Outside N = 56	Reserve N = 21	Outside N = 101	Reserve N = 30	Outside N = 11

**Appendix 2:** Mean survival and mean flight-initiation distance (FID) for each bird species used in the survival analysis. All survival estimates, number of captures and recaptures as well as 95% CI were extracted from Stevens et al. (2013)

Species <sup>1</sup>	Scientific name	Mean survival ± SE	Number of captures	Number of recaptures	95% CI of survival estimates	Mean FID ± SE (m)
FCHAT	<i>Cercomela familiaris</i>	0.54 ± 0.06	45	37	0.41–0.66	15.0 ± 0.8
RECCB	<i>Uraeginthus bengalus</i>	0.46 ± 0.04	498	226	0.38–0.54	10.4 ± 0.7
ROCFF	<i>Lagonosticta sanguinodorsalis</i>	0.56 ± 0.04	305	300	0.48–0.63	12.4 ± 1.0
COMBU	<i>Pycnonotus barbatus</i>	0.67 ± 0.05	311	111	0.56–0.76	17.1 ± 0.5
AFRTH	<i>Turdus pelios</i>	0.80 ± 0.06	190	87	0.66–0.89	18.5 ± 2.8
SPEMO	<i>Colius striatus</i>	0.62 ± 0.05	223	95	0.51–0.72	17.25 ± 2.1

<sup>1</sup> FCHAT = Familiar Chat, RECCB = Red-cheeked Cordon-bleu, ROCFF = Rock Firefinch, COMBU = Common Bulbul, AFRTH = African Thrush, SPEMO = Speckled Mousebird