

OKI2018 RefSeq a XP_081641266.1; b XP_081642886.1. Transcripts (top to bottom): TRINITY_DN6371_c0_g1_i1, TRINITY_DN6371_c0_g1_i4, TRINITY_DN6371_c0_g1_i3.
OSKA2016 ANISEED S8.g08683.01.t. Transcripts (top to bottom; 4 of the 7 aligned across the anchor codon, most introns first): comp20351_c1_seq1, comp20351_c1_seq5, comp20351_c1_seq3, comp20351_c1_seq2.
OdB3 GenBank CBY09053.1. Transcripts (top to bottom; 4 of the 13 aligned across the anchor codon, most introns first): KT0AAA134YG12RM1, KT0AAA276YO11RM1, KT0AAA145YJ12RM1, KT0AAA417YN19RM1.
proteins human Q9HD40; C. robusta XP_002120570.1.

exon: coding (thick), UTR (thin) — public model — gold-standard model — coding sequence in another frame
start codon of the gold-standard model (x axis: nucleotides from it, gene orientation) — ◂ n number of exons beyond the drawn region
a b public models of the same row, as named in the paragraph
transcript aligned to the assembly (exons) — bases of the transcript missing from the assembly (number when more than one)
bases of the assembly missing from the transcript — mismatches
intron splice sites (drawn length unchanged): GT-AG (colour of the model) — GA-AG — GC-AG — GG-AG — other site
Pfam domain of the first reference — residues not aligned to the first reference
92 % similar residues to the OdB3 protein (BLOSUM62 > 0; global alignment, aligned part) — human: UniProt; Ciona: RefSeq