

OKI2018 RefSeq XP_081645007.1. Transcripts (top to bottom; 4 of the 5 that hold the anchor codon, most introns first): TRINITY_DN748_c0_g1_i7, TRINITY_DN748_c0_g1_i4, TRINITY_DN748_c0_g1_i5, TRINITY_DN748_c0_g1_i3.
OSKA2016 ANISEED S5.g06617.01.t. Gold standard: no model: no Selenoprofiles prediction of a second SEPHS copy in this assembly. Transcripts (top to bottom): comp24718_c1_seq1.
OdB3 GenBank a GSOID_T00009332001; b CBY21562.1; c CBY21563.1. Gold standard: no model: no Selenoprofiles prediction of a second SEPHS copy in this assembly.
proteins human P49903; C. intestinalis Selenoprofiles prediction, glycine form.

exon: coding (thick), UTR (thin) — public model — gold-standard model — coding sequence in another frame
start codon of the gold-standard model (x axis: nucleotides from it, gene orientation) — n number of exons beyond the drawn region
a b public models of the same row, as named in the paragraph
transcript aligned to the assembly (exons) — bases of the transcript missing from the assembly (number when more than one)
bases of the assembly missing from the transcript — mismatches
intron splice sites (drawn length unchanged): GT-AG (colour of the model) — GA-AG — other site
Pfam domain of the first reference — residues not aligned to the first reference
92 % similar residues to the OdB3 protein (BLOSUM62 > 0; global alignment, aligned part) human: UniProt; C. intestinalis: RefSeq