

OKI2018 RefSeq a XP_081629121.1; b XP_081629120.1. Transcripts (top to bottom; 4 of the 6 aligned across the anchor codon, most introns first): TRINITY_DN4400_c0_g1_i4 (SL), TRINITY_DN4400_c0_g1_i1 (SL), TRINITY_DN4400_c0_g1_i8 (SL), TRINITY_DN4400_c0_g1_i11 (SL).
OSKA2016 ANISEED S28.g15187.01.t. Transcripts (top to bottom): comp23981_c0_seq1 (SL).
OdB3 GenBank CBY07707.1. Transcripts (top to bottom; 4 of the 5 aligned across the anchor codon, most introns first): KT0AAA4YE16NM1, KT0AAA399YH18RM1, KT0AAA331YJ10RM1, KT0AAA400YK11RM1.
proteins human P57772; C. robusta XP_002124067.1.

exon: coding (thick), UTR (thin) — public model — gold-standard model — coding sequence in another frame
start codon of the gold-standard model (x axis: nucleotides from it, gene orientation) — ◂ n number of exons beyond the drawn region
a b public models of the same row, as named in the paragraph
transcript aligned to the assembly (exons) — bases of the transcript missing from the assembly (number when more than one)
bases of the assembly missing from the transcript — mismatches — SL transcript that starts with the spliced leader
intron splice sites (drawn length unchanged): GT-AG (colour of the model) — GA-AG — GC-AG — other site
Pfam domain of the first reference — residues not aligned to the first reference
92 % similar residues to the OdB3 protein (BLOSUM62 > 0; global alignment, aligned part) — human: UniProt; Ciona: RefSeq