


**OKI2018** RefSeq XP_081645007.1. Transcripts (top to bottom; 4 of the 5 aligned across the anchor codon, most introns first): TRINITY_DN748_c0_g1_i7 (SL), TRINITY_DN748_c0_g1_i4 (SL), TRINITY_DN748_c0_g1_i5, TRINITY_DN748_c0_g1_i3 (SL).

**OSKA2016** ANISEED S5.g06617.01.t. Gold standard: no Selenoprofiles prediction; anchor from an alignment of the OKI2018 protein (curation/machinery_extra_anchors.tsv); exon chain of the transcript. Transcripts (top to bottom): comp24718_c1_seq1 (SL).

**OdB3** GenBank a GSOID_T00009332001; b CBY21562.1; c CBY21563.1. Gold standard: no Selenoprofiles prediction; anchor from an alignment of the OKI2018 protein; locus split by two assembly gaps (100 and 675 N): 34 residues read from the first gap (X), no EST; protein left out of the trees.

**proteins** human P49903; C. intestinalis Selenoprofiles prediction, glycine form.

exon: coding (thick), UTR (thin) — public model — gold-standard model — coding sequence in another frame

start codon of the gold-standard model (x axis: nucleotides from it, gene orientation) — ◂ n number of exons beyond the drawn region

a b public models of the same row, as named in the paragraph

transcript aligned to the assembly (exons) — bases of the transcript missing from the assembly (number when more than one)

bases of the assembly missing from the transcript — mismatches — SL transcript that starts with the spliced leader

intron splice sites (drawn length unchanged): GT-AG (colour of the model) — GA-AG — other site

Pfam domain of the first reference — residues not aligned to the first reference

92 % similar residues to the OdB3 protein (BLOSUM62 > 0; global alignment, aligned part) — human: UniProt; Ciona: RefSeq