


OKI2018 RefSeq XP_081642556.1: first three exons only; reads on into the third intron (110 codons, 98 of them as in the gold standard). Gold standard: third and fourth introns with splice sites outside the intron classes (TG-AG, AG-AA), as in the three supporting transcripts. Transcripts (top to bottom): TRINITY_DN16768_c0_g1_i1 (SL), TRINITY_DN16768_c0_g1_i3 (SL), TRINITY_DN16768_c0_g1_i4.

OSKA2016 ANISEED S10.g09825.01.t: starts in the fourth exon and continues past the stop codon into seven exons downstream (604 codons). Gold standard: third and fourth introns with splice sites outside the intron classes (AA-AG, TT-GT), as in the supporting transcripts; the third intron is retained in two of them. Transcripts (top to bottom): comp20361_c0_seq1 (SL), comp20361_c0_seq5 (SL), comp20361_c0_seq6, comp20361_c0_seq4.

OdB3 GenBank CBY23135.1: same first three exons; other first site of the fourth exon, and no last exon (178 codons). Gold standard: exon chain set by the curator: the GenBank exons 1 to 3, the acceptor of the third intron (GA-AG; retained in both ESTs) from the prediction, and the last intron of the ESTs (AT-AG). Transcripts (top to bottom): KT0AAA259YN02RM1 (SL), KT0AAA109YN17RM1. Also starting with the SL, not drawn: FP831546, FP830122.

proteins human Q8IV42; C. intestinalis XP_002120985.2.

C Equivalent placements of introns

Third intron


OKI2018	cgaataaTGCAGGGT … TTATCAGtctttatt	TG-AG	GC-GT	
KUM-M3-7f	cgaataaTGCAGGGT … TTATCAGtctttatt	TG-AG	GC-GT	
	R I I F I			
OSKA2016	cgaatAGAAGGGT … TTCAACGagttttc	AG-CG	GA-GA	AA-AG
AOM-5-5f	cgaatAGAAGGGT … TTCAACGagttttc	AG-CG	GA-GA	AA-AG
OdB3	cgaatAGGAGGGT … TTCAACGagttttc	AG-CG	GG-GA	GA-AG
Bar2_p4	cgaatAGGAGGGT … TTCAACGagttttc	AG-CG	GG-GA	GA-AG
	R I V F			

Fourth intron

OKI2018	cacttAGTTTGGG … TTTTGAAagtagaa	AG-AA	GT-AA	TT-AG	TT-GT
KUM-M3-7f	cacttAGTTTGGG … TTTTGAAagtagaa	AG-AA	GT-AA	TT-AG	TT-GT
OSKA2016	catttAGTTTGGG … GTGTGACagtagaa	AG-AC	GT-CA	TT-AG	TT-GT
AOM-5-5f	catttAGTTTGGG … GTATGACagtagaa	AG-AC	GT-CA	TT-AG	TT-GT
OdB3	catttAGATCGGG … CTGTGTCagtggaa	AG-TC	GA-CA	AT-AG	
Bar2_p4	catttAGATCGGG … CTGTGTCagtggaa	AG-TC	GA-CA	AT-AG	
	H L V E				

Gene strand, cased for the leftmost placement: lowercase grey, exon; uppercase, intron. Shaded: bases found at both ends, across which the intron can be moved without changing the spliced sequence. Letters: amino acids under the first base of each codon (the codon split by the intron included), one line under the rows that translate alike; exon flanks trimmed to whole codons. Right: dinucleotides of each placement, from left to right; blue: placement on an AG acceptor (phase 1); ▲: placement named by the gold-standard script. Rows in italics: test assemblies (other individuals of the same species), exons carried from the gold standard.

D Simplified tree of PSTK, more eukaryotes and archaea (PMSF)



Nodes supported by less than 50% collapsed; ultrafast bootstrap supports of 70% or more shown. Wedges: clades of sequences of one group, from their ancestor to the deepest sequence; "unresolved": sequences of the other groups in one polytomy, drawn as one tip. Sequences added to the tree of Figure TREES: see Methods.

exon: coding (thick), UTR (thin) — public model — gold-standard model — coding sequence in another frame

start codon of the gold-standard model (x axis: nucleotides from it, gene orientation) — ◂n number of exons beyond the drawn region

a b public models of the same row, as named in the paragraph

transcript aligned to the assembly (exons) — bases of the transcript missing from the assembly (number when more than one)

bases of the assembly missing from the transcript — mismatches — SL transcript that starts with the spliced leader

intron splice sites (drawn length unchanged): GT-AG (colour of the model) — GA-AG — GG-AG — other site

Pfam domain of the first reference — residues not aligned to the first reference

92 % similar residues to the OdB3 protein (BLOSUM62 > 0; global alignment, aligned part) — human: UniProt; Ciona: RefSeq