SEPHS2, N-terminal domain, LG+R6
groups of one taxon collapsed; nodes with ultrafast bootstrap < 50 collapsed; supports >= 70 shown

| | length (aa) |
|---|---|
| Bacteria (2) | 361 (360-362) |
| Alveolates (2) | 690 (570-811) |
| sps1_Aureococcus_anophagefferens_2008-20 | 795 |
| gi_190358310_sp_O67139.3_SELD_AQUAE_gi_1 | 336 |
| gi_212283264_sp_Q9L4E1.2_SELD1_EUBAC_gi_ | 347 |
| Campylobacter_hominis_ATCC_BAA-381_gi_15 | 378 |
| Archaea (3) | 348 (335-356) |
| Fbor_SEPHS2.1 (selenocysteine) | 369 |
| Rhodnius_prolixus.SPS2 | 381 |
| Caenorhabditis_elegans_gi_17543426_ref_N | 378 |
| Monosiga_brevicollis_SPSsec_Monosiga_bre | 296 |
| Dictyostelium_discoideum_gi_1657427_gb_A | 370 |
| Kinetoplastids (3) | 393 (377-398) |
| sps2_Perkinsus_marinus_2008-2009_sps2_Pe | 392 |
| Stramenopiles (2) | 374 (356-392) |
| Pediculus_humanus.SPS2 | 377 |
| Mosquitoes (5) | 365 (323-371) |
| Flies (4) | 398 (398-398) |
| Olon_gold_SEPHS_b | 424 |
| Olon_SEPHS2.5 (unaligned) | 287 |
| OKI_gold_SEPHS_b | 635 |
| Ovan_gold_SEPHS_b | 334 |
| Oalb_gold_SEPHS_b | 545 |
| Olon_SEPHS2.2 (selenocysteine) | 380 |
| Ovan_SEPHS2.10 (selenocysteine) | 376 |
| Oalb_SEPHS2.3 (selenocysteine) | 315 |
| Olon_SEPHS2.4 (selenocysteine) | 379 |
| OSKA_gold_SEPHS2 | 391 |
| OKI_gold_SEPHS2 | 393 |
| OdB3_gold_SEPHS2 | 392 |
| Oikopleura_dioica.SPS2 | 373 |
| Pmam_SEPHS2.3 (glycine) | 403 |
| Csav_SEPHS2.3 (glycine) | 483 |
| Crob_SEPHS2.6 (glycine) | 483 |
| Cint_SEPHS2.5 (glycine) | 483 |
| Hror_SEPHS2.5 (glycine) | 412 |
| Scla_SEPHS2.4 (glycine) | 408 |
| Bsch_SEPHS2.3 (glycine) | 417 |
| Botryllus_schlosseri.SPS1 | 367 |
| Tdem_SEPHS2.1 (selenocysteine) | 431 |
| Stho_SEPHS2.1 (pseudo) | 414 |
| Pege_SEPHS2.5 (selenocysteine) | 422 |
| Bsch_SEPHS2.1 (selenocysteine) | 477 |
| Botryllus_schlosseri.SPS2 | 367 |
| Scla_SEPHS2.2 (selenocysteine) | 369 |
| Scla_SEPHS2.1 (selenocysteine) | 480 |
| Hror_SEPHS2.1 (selenocysteine) | 468 |
| Helobdella_robusta.SPS2 | 366 |
| Priapulus_caudatus.SPS2 | 397 |
| Annelids (2) | 370 (370-370) |
| Strigamia_maritima.SPS2 | 411 |
| Chelicerates (2) | 410 (402-418) |
| Lepeophtheirus_salmonis.SPS2 | 413 |
| Catajapyx_aquilonaris.SPS2 | 420 |
| Drosophila_pseudoobscura_pseudoobscura_g | 409 |
| Crustaceans (2) | 416 (416-416) |
| PediculusA_PediculusA | 396 |
| Other insects (2) | 413 (413-413) |
| Other insects (2) | 413 (413-413) |
| Flies (4) | 398 (398-398) |
| Mosquitoes (4) | 397 (376-398) |
| Blan_SEPHS2.2 (selenocysteine) | 389 |
| Bflo_SEPHS2.5 (selenocysteine) | 389 |
| Petromyzon_marinus.SPS2 | 383 |
| Ray-finned fishes (4) | 398 (384-447) |
| Latimeria_chalumnae.SPS2 | 376 |
| Reptiles (2) | 376 (375-376) |
| Mammals (7), with human SEPHS2 | 448 (390-452) |
| Tetraodon_nigroviridis.SPS1 | 377 |
| Xiphophorus_maculatus.SPS1 | 381 |
| Ray-finned fishes (2) | 385 (380-390) |
| Latimeria_chalumnae.SPS1 | 376 |
| Anolis_carolinensis.SPS1 | 378 |
| Chrysemys_picta_bellii.SPS1 | 377 |
| Mammals (4), with human SEPHS1 | 387 (381-392) |

0 1 2 3 4 5 6 7

branch length

paralog of the named gene  clade of the human gene that names the family