

OKI2018 RefSeq XP_081630352.1: same exon chain; TGA read as stop.

OSKA2016 ANISEED S27.g15032.01.t: same coding exons; TGA read as stop. Gold standard: 5′ UTR exon from the transcripts.

OdB3 GenBank CBY22487.1: same exon chain; TGA read as stop.

proteins human O43396 (the O. dioica protein aligns with the PITH domain only, residues 120-283); C. intestinalis XP_078490413.1.



OKI2018 RefSeq XP_081632581.1: mRNA of another protein (173 aa) with this gene in its 3′ UTR.

OSKA2016 ANISEED S22.g13919.01.t: same exon chain; TGA read as stop.

OdB3 GenBank CBY21026.1: same exon chain; TGA read as stop.

proteins O. albicans SCLG01000111.1 (genome; open stretch of the comparative screen).

exon: coding (thick), UTR (thin) — public model — gold-standard model — coding sequence in another frame

selenocysteine TGA (x axis: nucleotides from it, gene orientation) — 37 best SECIS hit and bit score (Oikopleura model; open box: below 20 bits)

◀ n number of exons beyond the drawn region — a b public models of the same row, as named in the paragraph

intron splice sites (drawn length unchanged): GT-AG (colour of the model)

Pfam domain of the first reference — U selenocysteine aligned to one of the first reference — © other residue aligned to it

residues not aligned to the first reference — 92 % similar residues to the OdB3 protein (BLOSUM62 > 0; global alignment, aligned part)

human: UniProt; C. intestinalis: RefSeq; O. albicans: open stretch of the comparative screen on its genome (accessions in the paragraphs)