

OKI2018 RefSeq XP_081641373.1. Transcripts (top to bottom): TRINITY_DN5140_c0_g1_i2, TRINITY_DN5140_c0_g1_i3, TRINITY_DN5140_c0_g1_i1.
OSKA2016 ANISEED S1.g00233.01.t. Transcripts (top to bottom): comp4911_c0_seq1, comp4911_c0_seq2.
OdB3 GenBank CBY19450.1. Transcripts (top to bottom): KT0AAA377YB24RM1, KT0AAA213YM23RM1, KT0AAA149YI11RM1.
proteins human Q93073; C. intestinalis XP_078482116.1.

exon: coding (thick), UTR (thin) — public model — gold-standard model — coding sequence in another frame
start codon of the gold-standard model (x axis: nucleotides from it, gene orientation) — ◂ n number of exons beyond the drawn region
a b public models of the same row, as named in the paragraph
transcript aligned to the assembly (exons) — bases of the transcript missing from the assembly (number when more than one)
bases of the assembly missing from the transcript — mismatches
intron splice sites (drawn length unchanged): GT-AG (colour of the model) — GA-AG — GC-AG — GG-AG — other site
Pfam domain of the first reference — residues not aligned to the first reference
92 % similar residues to the OdB3 protein (BLOSUM62 > 0; global alignment, aligned part) human: UniProt; Ciona: RefSeq