TXNL1.b



OKI2018 RefSeq XP_081630352.1: same exon chain; TGA read as stop.
OSKA2016 ANISEED S27.g15032.01.t: same coding exons; TGA read as stop. Gold standard: 5′ UTR exon from the transcripts.
OdB3 GenBank CBY22487.1: same exon chain; TGA read as stop.
TXNL1.a OKI2018 RefSeq XP_081633344.1; OSKA2016 ANISEED S14.g11310.01.t (model of 756 residues; drawn: residues 478-756, aligned to human TXNL1); OdB3 GenBank CBY22771.1.
proteins human O43396 (the O. dioica protein aligns with the PITH domain only, residues 120-283); C. intestinalis XP_078490413.1.

KKSU



OKI2018 RefSeq XP_081632581.1: mRNA of another protein (173 aa) with this gene in its 3′ UTR.
OSKA2016 ANISEED S22.g13919.01.t: same exon chain; TGA read as stop.
OdB3 GenBank CBY21026.1: same exon chain; TGA read as stop.
proteins O. albicans SCLG01000111.1 (genome; open stretch of the comparative screen).

exon: coding (thick), UTR (thin) — public model — gold-standard model
TXNL1.a (public model; aligned on its codon aligned to the selenocysteine) — coding sequence in another frame
selenocysteine TGA (x axis: nucleotides from it, gene orientation) — 37 best SECIS hit and bit score (Oikopleura model; open box: below 20 bits)
◀ n number of exons beyond the drawn region — a b public models of the same row, as named in the paragraph
intron splice sites (drawn length unchanged): GT-AG (colour of the model)
Pfam domain of the first reference — U selenocysteine aligned to one of the first reference — © other residue aligned to it
residues not aligned to the first reference — 92 % similar residues to the OdB3 protein (BLOSUM62 > 0; global alignment, aligned part)
human: UniProt; C. intestinalis: RefSeq; O. albicans: open stretch of the comparative screen on its genome (accessions in the paragraphs)