

Chapter 5

Sequences, Mathematical Induction, and Recursion

Ojas Chaturvedi

Contents

5.1: Sequences	1
Notes	1
5.2: Mathematical Induction I	2
Notes	2
5.3: Mathematical Induction II	3
Different types of problems	3
Template for Mathematical Induction	6

5.1: Sequences

Notes

- Sequence: a function whose domain is either all the integers between two given integers, or all the integers greater than or equal to a given integer.

– Know subscript/index, initial and final term, infinite sequence, general/explicit formula

- Summation Notation:

$$\sum_{k=m}^n a_k = a_m + a_{m+1} + a_{m+2} + \cdots + a_n$$

where k is the index, m is the lower limit, and n is the upper limit.

- When the upper limit of a summation is a variable, an ellipsis is used to write the summation in expanded form. Expand the summation notation to first 3 or so, then put ellipsis and then variable form.

- Product Notation:

$$\prod_{k=m}^n a_k = a_m \cdot a_{m+1} \cdot a_{m+2} \cdots a_n$$

- Properties of Summations and Products (aka Theorem 5.1.1)

$$\sum_{k=m}^n a_k + \sum_{k=m}^n b_k = \sum_{k=m}^n (a_k + b_k) \quad (1)$$

$$c \cdot \sum_{k=m}^n a_k = \sum_{k=m}^n (c \cdot a_k) \quad (2)$$

$$\left(\prod_{k=m}^n a_k \right) \cdot \left(\prod_{k=m}^n b_k \right) = \prod_{k=m}^n (a_k \cdot b_k) \quad (3)$$

- When replacing a new variable into a summation or product, make sure to change the index variable to the new variable and the numbers by putting them into the equation of the new variable.
- Factorial: the quantity $n!$ is defined to be the product of all the integers from 1 to n :

$$n! = n \cdot (n-1) \cdots 3 \cdot 2 \cdot 1$$

and

$$0! = 1$$

Recursive definition:

$$n! = \begin{cases} 1 & \text{if } n = 0 \\ n \cdot (n-1)! & \text{if } n > 0 \end{cases}$$

- n choose r : the number of subsets (therefore an integer) of size r that can be chosen from a set of n elements.

$$\binom{n}{r} = \frac{n!}{r!(n-r)!}$$

for all integers n and r with $0 \leq r \leq n$.

5.2: Mathematical Induction I

Notes

- Principles of Mathematical Induction: Let $P(n)$ be a property that is defined for integers n , and let a be a fixed integer. Suppose the following two statements are true:
 1. Basis Step: Show that $P(a)$ is true.
 2. Inductive Step: For all integers $k \geq a$, if $P(k)$ is true, then $P(k + 1)$ is true.
 - To perform this step:
 - (a) Suppose that $P(k)$ is true for an arbitrary integer $k \geq a$, which is called the inductive hypothesis.
 - (b) Show that $P(k + 1)$ is true.
 - Remember that you need to prove each side of the equation separately. Otherwise, the proof is invalid.
 3. Conclusion: Then $P(n)$ is true for all integers $n \geq a$.
- Steps of Proof by Mathematical Induction:
 1. State the theorem to be proved.
 - Let the property $P(n)$ be the equation: problem goes here
 2. Prove the basis step.
 - Show that $P(a)$ is true.
 3. State the inductive hypothesis.
 - Show that for all integers $k \geq 1$, if $P(k)$ is true then $P(k + 1)$ is also true:
 4. Prove the inductive step.
 5. State the conclusion.
 - Therefore the equation $P(k + 1)$ is true *[as was to be shown]*. *[Since we have proved both the basis step and the inductive step, the conclusion follows by the principle of mathematical induction. Therefore the equation $P(n)$ is true for all integers $n \geq 1$.]*

- Sum of the first n integers is

$$1 + 2 + 3 + \cdots + n = \frac{n(n+1)}{2}$$

- Geometric sum of the first n integers is

$$\sum_{i=0}^n r^i = \frac{r^{n+1} - 1}{r - 1}$$

5.3: Mathematical Induction II

Different types of problems

Problem Type: Divisibility Property. For all integers $n \geq 0$, $2^{2^n} - 1$ is divisible by 3.

Proof. Let the property $P(n)$ be the sentence:

$$2^{2^n} - 1 \text{ is divisible by } 3$$

First, we must prove that $P(0)$ is true (basis step).

$$2^{2 \cdot 0} - 1 \text{ is divisible by } 3$$

$$\begin{aligned} 2^{2(0)} - 1 &= 2^0 - 1 \\ &= 1 - 1 \\ &= 0 \\ &= 3 \cdot 0 \end{aligned}$$

Thus, $P(0)$ is true.

Now, suppose that $P(k)$ is true for some integer $k \geq 0$ (inductive hypothesis). That is,

$$2^{2^k} - 1 \text{ is divisible by } 3$$

By the definition of divisibility, for some integer r ,

$$2^{2^k} - 1 = 3r$$

We must show that $P(k+1)$ is true (inductive step). That is,

$$2^{2^{(k+1)}} - 1 \text{ is divisible by } 3$$

The left-hand side of $P(k+1)$ is:

$$\begin{aligned} 2^{2^{(k+1)}} - 1 &= 2^{2^{k+2}} - 1 \\ &= 2^{2^k} \cdot 2^2 - 1 \\ &\quad \text{by the product rule for exponents} \\ &= 4 \cdot 2^{2^k} - 1 \\ &= 3 \cdot 2^{2^k} + 2^{2^k} - 1 \\ &= 3 \cdot 2^{2^k} + 3r \\ &\quad \text{by substituting the inductive hypothesis} \\ &= 3(2^{2^k} + r) \end{aligned}$$

$2^{2^k} + r$ is an integer since it is the sum of products of integers, so $2^{2^{(k+1)}} - 1$ can be written as $6m$ for some integer $m = (2^{2^k} + r)$.

By the definition of divisibility, $2^{2^{(k+1)}} - 1$ is divisible by 3, and thus, $P(k+1)$ is true.

Since we have proved both the basis step and the inductive step, we conclude that the statement is true using mathematical induction. \square

Problem Type: Inequality. For all integers $n \geq 3$, $2n + 1 < 2^n$.

Proof. Let the property $P(n)$ be the inequality:

$$2n + 1 < 2^n$$

First, we must prove that $P(3)$ is true (basis step).

$$\begin{aligned} 2(3) + 1 &< 2^3 \\ 7 &< 8 \end{aligned}$$

Thus, $P(3)$ is true.

Now, suppose that $P(k)$ is true for some integer $k \geq 3$ (inductive hypothesis). That is,

$$2k + 1 < 2^k$$

We must show that $P(k + 1)$ is true (inductive step). That is,

$$\begin{aligned} 2(k + 1) + 1 &< 2^{k+1} \\ 2k + 3 &< 2^{k+1} \end{aligned}$$

The left-hand side of $P(k + 1)$ is:

$$\begin{aligned} 2k + 3 &= 2k + 1 + 2 \\ &< 2^k + 2 && \text{by substitution of the inductive hypothesis} \\ &< 2^k + 2^k && \text{because } 2 < 2^k \text{ for all integers } k \geq 2 \\ &< 2 \cdot 2^k \\ &< 2^{k+1} && \text{by the product rule for exponents} \end{aligned}$$

Thus, the left-hand side of $P(k + 1)$ is less than the right-hand side of $P(k + 1)$, and $P(k + 1)$ is true. Since we have proved both the basis step and the inductive step, we conclude that the statement is true using mathematical induction. \square

Problem Type: Property of a Sequence. Define a sequence a_1, a_2, a_3, \dots as follows:

$$\begin{aligned} a_1 &= 2 \\ a_k &= 5a_{k-1} && \text{for all integers } k \geq 2 \end{aligned}$$

- Write the first four terms of the sequence.
- It is claimed that for each integer $n \geq 0$, the n th term of the sequence has the same value as that given by the formula $2 \cdot 5^{n-1}$. In other words, the claim is that the terms of the sequence satisfy the equation $a_n = 2 \cdot 5^{n-1}$. Prove that this is true.

Proof.

a. $a_1 = 2$

$$a_2 = 5a_{2-1} = 5a_1 = 5 \cdot 2 = 10$$

$$a_3 = 5a_{3-1} = 5a_2 = 5 \cdot 10 = 50$$

$$a_4 = 5a_{4-1} = 5a_3 = 5 \cdot 50 = 250$$

- b. Let a_1, a_2, a_3, \dots be the sequence defined by specifying that $a_1 = 2$ and $a_k = 5a_{k-1}$ for all integers $k \geq 2$. Let the property $P(n)$ be the equation:

$$a_n = 2 \cdot 5^{n-1}$$

First, we must prove that $P(1)$ is true (basis step).

$$a_1 = 2 \cdot 5^{1-1}$$

The left-hand side of $P(1)$ is

$$a_1 = 2$$

by the definition of a_1, a_2, a_3, \dots

The right-hand side of $P(1)$ is

$$\begin{aligned} 2 \cdot 5^{1-1} &= 2 \cdot 5^0 \\ &= 2 \cdot 1 \\ &= 2 \end{aligned}$$

Thus, the left-hand side of $P(1)$ is equal to the right-hand side of $P(1)$, and $P(1)$ is true. Now, suppose that $P(k)$ is true for some integer $k \geq 1$ (inductive hypothesis). That is,

$$a_k = 2 \cdot 5^{k-1}$$

We must show that $P(k+1)$ is true (inductive step). That is,

$$\begin{aligned} a_{k+1} &= 2 \cdot 5^{(k+1)-1} \\ a_{k+1} &= 2 \cdot 5^k \end{aligned}$$

The left-hand side of $P(k+1)$ is:

$$\begin{aligned} a_{k+1} &= 5a_k \\ &= 5(2 \cdot 5^{k-1}) \\ &= 2 \cdot 5^k \end{aligned}$$

Thus, the left-hand side of $P(k+1)$ is equal to the right-hand side of $P(k+1)$, and $P(k+1)$ is true.

Since we have proved both the basis step and the inductive step, we conclude that the statement is true using mathematical induction.

□

Template for Mathematical Induction

Let the property $P(n)$ be the equation/sentence/inequality:

{problem goes here}

First, we must prove that $P(\{\text{smallest possible number goes here}\})$ is true (basis step).

Show left-hand side = right-hand side of the equation

Thus, $P(\{\text{smallest possible number goes here}\})$ is true.

Now, suppose that $P(k)$ is true for some integer $k \geq \{\text{smallest possible number goes here}\}$ (inductive hypothesis). That is,

{problem with k substituted goes here}

We must show that $P(k+1)$ is true (inductive step). That is,

{problem with (k+1) substituted goes here}

The left-hand side of $P(k+1)$ is:

{work with reasoning goes here}

The right-hand side of $P(k+1)$ is:

{work with reasoning goes here}

Thus, the left-hand side of $P(k+1)$ is equal to the right-hand side of $P(k+1)$, and $P(k+1)$ is true.

Since we have proved both the basis step and the inductive step, we conclude that the statement is true using mathematical induction.