Asteroseismology provides a new range of internal rotation periods and ages in which to calibrate gyrochronology

EXPANDING THE GYROCHRONOLOGY RELATION WITH ASTEROSEISMIC ROTATION AND AGE



Oliver J. Hall, Guy R. Davies, Martin B. Nielsen
+ friends


INTRO


- The rotation of stars slows
 down as they age, at a rate that
 is a function of colour.
- We can calibrate this
 'gyrochronology' relation to
 help us estimate stellar age.
- van Saders+16 showed that some old stars stop slowing down at a certain point. Why?
- Asteroseisic measurements of age and interal rotation periods allow us to **study older stars** in this area of interest.

METHOD

- We use the Davies+16, Lund+17, Silva Aguirre+15,17 samples for their ages and locations of individual frequencies.
- We obtain rotation periods for 54 stars (so far!) by fitting a holistic model to the l = {0, 1, 2} p-modes, treating the mode frequencies as latent variables.
- We fit the classical Barnes+07 gyrochronology relation with Gaia BP-RP colours using latent variables to treat uncertainties in the three observeables.

RESULTS SO FAR






STELLAR ASTROPHYSICS CENTRE

ROTATIONAL SPLITTING

- The **rotation** and **inclination** of a star change how modes appear.
- The examples below are for a star with a period of 10 days and an inclination of 45°.



WHATS NEXT?

Frequency from centroid (μ Hz)

- Improving the fitting process with Gaussian Process priors on linewidth and height.
- Fitting an improved gyrochronology model that treats mass, metallicity, and the heteroskedastic uncertainties.
- Comparing our data to evolutionary models and clusters to figure out when (if?) rotation stops slowing.



