

New asteroseismic rotation rates of *Kepler* dwarfs show strong agreement with weakened magnetic braking on the late-age main sequence

Oliver J. Hall^{1,2,3,*}, Guy R. Davies^{2,3}, Jennifer van Saders^{4,5,6}, Martin B. Nielsen^{2,3}, Mikkel N. Lund³, William J. Chaplin^{2,3}, Rafael A. García^{7,8}, Louis Amard⁹, Angela A. Breimann⁹, Saniya Khan^{2,3}, Victor See⁹, Jamie Tayar^{4,10}

¹ European Space Agency (ESA), European Space Research and Technology Centre (ESTEC), Keplerlaan 1, 2201 AZ Noordwijk, The Netherlands

² School of Physics and Astronomy, University of Birmingham, Edgbaston, Birmingham, B15 2TT, UK

³ Stellar Astrophysics Centre, Department of Physics and Astronomy, Aarhus University, Ny Munkegade 120, 8000 Aarhus C, Denmark

⁴ Institute for Astronomy, University of Hawai'i, Honolulu, HI 96822

⁵ Observatories of the Carnegie Institution for Science, Pasadena, CA 91101

⁶ Department of Astrophysical Sciences, Princeton University, Princeton, NJ 08544

⁷ IRFU, CEA, Université Paris-Saclay, F-91191 Gif-sur-Yvette, France

⁸ AIM, CEA, CNRS, Université Paris-Saclay, Université Paris Diderot, Sorbonne Paris Cité, F-91191 Gif-sur-Yvette, France

⁹ University of Exeter Department of Physics and Astronomy, Stocker Road, Devon, Exeter, EX4 4QL, UK

¹⁰ Hubble Fellow

Studies using asteroseismic ages and rotation rates from star-spot rotation have indicated that standard age-rotation relations may break down roughly half-way through the main sequence lifetime, a phenomenon referred to as weakened magnetic braking. While rotation rates from spots can be difficult to determine for older, less active stars, rotational splitting of asteroseismic oscillation frequencies can provide rotation rates for both active and quiescent stars, and so can confirm whether this effect really takes place on the main sequence.

We obtained asteroseismic rotation rates of 91 main sequence stars showing high signal-to-noise modes of oscillation. Using these new rotation rates, along with effective temperatures, metallicities and seismic masses and ages, we built a hierarchical Bayesian mixture model to determine whether the ensemble more closely agreed with a standard rotational evolution scenario [1], or one where weakened magnetic braking takes place [2]. The weakened magnetic braking scenario was found to be 98.4% more likely for our stellar ensemble, adding to the growing body of evidence for this stage of stellar rotational evolution. This work represents the largest catalogue of seismic rotation on the main sequence to date, opening up possibilities for more detailed ensemble analysis of rotational evolution with *Kepler*.

*Email: oliver.hall@esa.int — Twitter: @asteronomer — GitHub: @ojhall94

Gyrochronology is the study of the relationship between a star’s rotation period and its age. As a star grows older along the main sequence, magnetic winds will cause it to lose angular momentum, slowing it down. Because the loss rate is related to temperature, the rotation period of a young star will rapidly settle on to a plane in age-colour-rotation space [3]. As a result, knowing the rotation and colour of a star provides an avenue to measure its age, to a precision of $\sim 10\%$ for stars most similar to the Sun [4]. Age can otherwise be difficult to come by, and so gyrochronology enables more in-depth studies of stellar populations and individual stars [5, 6].

Gyrochronology was previously calibrated on stellar clusters, which have well constrained ages, but only up to roughly 2.5 Gyr so far [4]. With the launch of the *Kepler* mission [7], ages of main sequence field stars (i.e. not in clusters) became more widely available through asteroseismology, the study of stellar variability [8]. When looking at these stars, disagreements were found between the observations and gyrochronology beyond the middle of stars’ main sequence lifetimes, which could not be reconciled with existing theories [9, 10, 11]. It was proposed that at some stage in a star’s evolution it undergoes *weakened magnetic braking* (WMB, [2]), where the efficiency of angular momentum loss rapidly drops. This would cause stars to keep fast rotation rates at old ages, which we would not expect from existing gyrochronology relations.

The mechanism by which weakened magnetic braking occurs has since been the subject of ongoing debate, and may be connected to changes in the magnetic field morphology [12, 13, 14, 15]. It may also be explained from an observational point of view. A large scale survey of stellar rotation rates measured using star-spots pointed out a lack of slowly rotating stars older than Sun [16]. As activity reduces with age, older stars with fewer star-spots are less likely to have their rotation measured [17]. The point at which the detection probability drops appears to lie at a similar level of activity as when the proposed weakened magnetic braking takes place, pitting these two possibilities against one another [18].

Determining whether weakened magnetic braking is a true phenomenon or a bias in star-spot observations requires new data, which can be provided by asteroseismology. A star’s rotation causes modes of oscillation to split into multiplets, making it possible to obtain an independent rate of rotation by measuring the splitting between oscillation frequencies [19]. Asteroseismic rotation is more difficult to obtain than spot rotation, but does not require spots to be visible, meaning that we can measure rotation for older, quiescent stars that would not have been present in existing rotation catalogues. This makes asteroseismic rotation rates the perfect avenue to explore whether weakened magnetic braking truly takes place on the main sequence. Here, we measured asteroseismic rotation rates of 91 main sequence stars across a broad range of ages, and evaluated this new ensemble to determine whether it agrees more closely with a classical rotational evolution scenario, or one where weakened magnetic braking takes place.

Asteroseismic main sequence targets: In order to obtain robust asteroseismic rotation rates, we required detections of multiples of both dipole (denoted as $\ell = 1$) and quadrupole ($\ell = 2$) p mode¹ oscillations for each star, of which the latter have significantly lower signal-to noise. Radial oscillations ($\ell = 0$), which are also visible, do not split. In this work, we

¹*Pressure* modes, the type of oscillation visible in main sequence stars.

studied a sample of 95 of the highest signal-to-noise targets observed by *Kepler*, combining the ‘Kages’ [8, 20] and LEGACY [21, 22] catalogues². Further details on these data can be found in **Methods**.

We separated our sample into three categories [23]: 64 main sequence stars (MS, $\log(g) > 4.0$ dex, $T_{\text{eff}} < 6250$ K); 4 potential sub-giant stars (SG, $\log(g) < 4.0$ dex, $T_{\text{eff}} < 6250$ K), which may have begun to evolve off the main sequence, complicating their rotational history³; and 24 ‘hot’ stars (H, $T_{\text{eff}} > 6250$ K), which lie above the Kraft break, the point at which convective envelopes are tenuous, and angular momentum loss through magnetic winds becomes inefficient [25]. Our conclusions are largely insensitive to these categorical assignments, and they are intended solely to help explore results category-by-category. The sample, which can be seen in Figure 1, spans surface gravities of $3.8 \text{ dex} < \log(g) < 4.6 \text{ dex}$, temperatures of $5000 \text{ K} < T_{\text{eff}} < 6700 \text{ K}$, and ages from 1 to ~ 13 Gyr.




Figure 1: Our sample of 95 stars from the ‘Kages’ and LEGACY catalogues coloured by asteroseismic age [8, 22]. The dashed lines indicate our classification boundaries: main sequence (circles, $T_{\text{eff}} < 6250$ K, $\log(g) < 4$ dex), potential sub-giants (squares, $T_{\text{eff}} < 6250$ K, $\log(g) > 4$ dex), and ‘hot’ stars (triangles, $T_{\text{eff}} > 6250$ K). The Sun is shown with the ‘ \odot ’ symbol, and has an age of 4.6 Gyr [26]. The solid lines are evolutionary tracks generated using MESA [27], for a metallicity of $Z = 0.01493$ and helium content of $Y = 0.26588$. Left to right, they represent masses of 1.5 , 1.25 , 1 and $0.75 M_{\odot}$. Vertical errorbars are too small to be visible [28].

²For these catalogues the mode extraction through frequency fitting [20, 21] and modelling using mode frequencies to obtain stellar parameters [8, 22] are covered in separate papers.

³It should be noted that none of these stars have identified mixed dipole modes, which indicate an evolved structure (see [24]).

Results of asteroseismic fitting: The stars in ‘Kages’ and LEGACY which form our stellar sample have been subject to detailed asteroseismic frequency fitting in the catalogue papers, but did not have reported asteroseismic rotation. Here, we repeated the mode frequency fitting with a Bayesian model that accounted for rotation in more detail, and improved the treatment of existing asteroseismic relations in frequency fitting techniques (see **Methods** and **Supplementary Information**). We fit our new model to *Kepler* power-spectra of 95 stars, which was successful in 94 cases. The summary statistics used throughout the rest of this paper were taken as the median of the posterior distribution on the model parameters, with the uncertainties being the 15.9th and 84.1st percentiles, meaning that some reported parameters (such as period) have asymmetric uncertainties.

We compared our results with both published and unpublished asteroseismic rotation rates [10, 11, 20, 21, 29], and found that the three stars with the lowest angles of inclination ($< 10^\circ$) were inconsistent with multiple other studies. To some degree this was expected, as for stars viewed near pole-on the split modes of oscillation have extremely low power, making them hard to detect reliably. We flagged our rotation rates for these stars, and did not use them in the next steps in our analysis, leaving a sample of 91 stars for which we considered our asteroseismic rotation measurements to be robust (for a full justification, see **Methods**).

Seismic versus surface rotation rates: Different methods of rotation measurement probe different depths of stars. Asteroseismology of main sequence stars probes internal rotation in the near surface layers, where the observed modes of oscillation are most sensitive [30]. Measurements of star-spot modulation instead probe the rotation rates of star-spots on the surface. Previous studies have shown that seismic rotation rates show no statistically significant deviation from spot rotation rates [10, 31]. We confirm that this holds true for our larger sample, in order to ensure that our seismic rotation periods can be reliably used to draw conclusions on the evolution of surface rotation.

Figure 2 shows a comparison between 48 stars from our sample for which spot rotation rates had previously been measured. Rotation rates from spot modulation are subject to measuring multiples of the true rotation rate, and also sometimes over-correct for this effect. This is not the case for asteroseismic rotation, and so some stars may appear at the 2:1 and 1:2 lines in the figure. We fit a line of the form $P_{\text{surf}} = m \times P_{\text{seis}}$, using the larger of the asymmetrical uncertainties on asteroseismic rotation, and excluding stars above the 1.8:1 line to avoid outliers (transparent in the Figure). Our fit found a value of $m = 0.96 \pm 0.03$, showing a close agreement ($< 2\sigma$) between asteroseismic and surface rotation rates on a population level. This overall agreement between spot and seismic rotation rates is in line with previous studies of Sun-like stars [10, 31, 32, 33], and so we concluded that our asteroseismic ensemble can be reliably used to draw inferences about gyrochronology.

Comparing gyrochronology models: To evaluate the implications of our seismic ensemble for gyrochronology, we compared our new asteroseismic rotation rates to two population models of rotational evolution [18]: a ‘standard’ model, which assumes a traditional angular momentum transport through magnetically driven stellar winds [35, 36], and one where weakened magnetic braking takes place (hereafter the ‘WMB model’) [2] The WMB model was identical to the standard model in its input physics, except for the condition that




Figure 2: Comparisons between asteroseismic and photometric measures of stellar rotation, for 48 stars. Literature values were taken, in order of priority, from: Garcia et al. (2014) [23] and Nielsen et al. (2013) [34] (i.e. if a value was reported in Nielsen et al. (2013) and also in Garcia et al. (2014), the latter was used). Stars used in the original van Saders et al. (2019) [2] study that first proposed weakened magnetic braking are highlighted. The four triangles represent stars included in the Nielsen et al. (2015) study [10] that previously measured this relationship. The solid line indicates 1:1 line, while the dash-dotted lines represent 2:1 and 1:2 lines. The shaded region around the 1:1 line is the 1σ credible interval of the fit to the data, the result of which is shown as the title of the figure. Stars that were not included in the fitting process are transparent. The errorbars on the asteroseismic rotation represent the 15.9th and 84.1st percentiles on the measured posterior distributions.

angular momentum loss ceases above a critical Rossby number of $Ro_{\text{crit}} = 1.97^4$.

Instead of comparing our rotation rates directly to models of individual stars, we instead compared them to synthetic TRILEGAL populations based on what the *Kepler* field would look like under these two different braking laws [18, 37]. These synthetic populations were then altered to match the distribution of existing catalogues of stars in the *Kepler* field in order to replicate the *Kepler* selection effect [38], and evolved following a standard or WMB braking law. For the full details on how these models were constructed, see **Methods**.

In order to ensure our results were based on the most robust rotation measurements, we focused on the stars with the best convergence metrics in the mode frequency fitting, excluding

⁴The Rossby number may be defined as $Ro \equiv P/\tau_{\text{cz}}$, where τ_{cz} is the convective turnover timescale, and P is the rotation. The Rossby number roughly scales inversely with stellar activity (and so increases as stars age).

those with a number of effective samples $n_{\text{eff}} < 1000$ and a Gelman-Rubin metric of $\hat{R} > 1.1$ [39, 40] (see **Extended Data** Table 1). Two stars with low metallicities (< -0.8 dex) that fell outside the boundaries of our stellar models were also excluded. The remaining sample of 73 stars contained 4 potential sub-giants, 22 ‘hot’ stars and 47 main sequence stars. Figure 3 shows these stars plotted over the top of both the standard and WMB population models in rotation-temperature space, alongside the stars not used in the gyrochronology analysis (shown without uncertainties).

We evaluated our ensemble against both population models simultaneously using a Bayesian mixture model, which treated the data as being drawn from a mixture made from Kernel Density Estimates (KDEs) of both populations, modulated by a mixing parameter P_s . In the limit that $P_s \rightarrow 1$, the data are most likely drawn from the standard model. In the limit $P_s \rightarrow 0$, the WMB model is more likely. This mixture model was evaluated in a five-dimensional parameter space, accounting for the population distributions in effective temperature (T_{eff}), metallicity ($[\text{Fe}/\text{H}]$), asteroseismic mass (M) and age (t) and our new rotation periods (P). For further detail, see **Methods**.

Results of comparing models: We fit our mixture model to each star individually, which yielded a posterior probability density for the value of P_s for that star. In order to assess the posterior probability for the full sample, we multiplied the individual posterior probabilities for P_s . This was done for the full ensemble of 73 stars together, as well as for the three different stellar types separately. These joint probabilities can be seen in Figure 4 in the left-hand plot. The cumulative probabilities can be seen in the right-hand plot. In both of these plots, the left-hand side holds 98.4% of the probability, indicating that the weakened magnetic braking model is preferred. When only considering the 47 main sequence stars, for which we expect the effects of weakened magnetic braking to be strongest, this rises to 99.2%.

Both hot and sub-giant stars do not strongly prefer one model over the other (both crossing the $P_s = 0.5$ line at roughly 45%), which is expected given that the rotation rates of both these stellar types are not expected to be strongly influenced by weakened magnetic braking. When performing this analysis on all 89 stars for which seismic rotation was measured (excluding the two stars with low metallicity), this the total probability below $P_s = 0.5$ is 96.6%.

Verifying consequences for gyrochronology: In order to verify that our results presented a meaningful confirmation of weakened magnetic braking, we ensured that the same results were obtained when using different subsets of the ensemble, and when exploring potential biases. We found that the joint posterior distribution did not change significantly when excluding stars with a metallicities that fell only slightly outside the model range, or those with extremely small uncertainties on asteroseismic mass (see **Methods**). Stars in our sample with known binary companions that may disrupt their rotational evolution were not found to contribute strongly to the joint posterior probability of P_s .

We checked that our results remained the same if the ‘Kages’ and LEGACY catalogues had underestimated the uncertainties on the asteroseismic mass and age, which form part of our mixture model analysis. We reran our analysis after inflating the uncertainties on these properties to align with the expected systematic uncertainty from stellar models [8, 22]. The results from this echoed what we found for the unaltered ensemble. We also accounted for the possibility that asteroseismic methods systematically overestimated stellar age, and reran




Figure 3: Stars for which rotation was measured in this work, plotted over TRILEGAL [37] stellar population models evolved under standard (top) and weakened magnetic braking (WMB, bottom) prescriptions of angular momentum evolution. The 21 Stars that were excluded from the gyrochronology analysis are marked with crosses and without uncertainties (see text). Circles, squares and triangles denote main sequence, sub-giant, and ‘hot’ stars respectively. Stars with a metallicity of $|\text{[Fe/H]}| > 0.4$ are shown in purple, to indicate that they fall outside the functional range of the stellar models [18]. Note that only period-temperature space is shown here, but when evaluating between the two model prescriptions we also considered mass, age and metallicity. The errorbars on the asteroseismic rotation represent the 15.9th and 84.1st percentiles on the measured posterior distributions.




Figure 4: Posterior estimates of the mixture model parameter P_s , broken down by stellar classification. A value of P_s close to zero indicates that the data are more consistent with a rotational evolution that includes weakened magnetic braking, whereas a value close to 1 indicates that the data are more consistent with a standard rotational evolution scenario. *Left:* the product of histograms of the posterior estimates for 73 stars, or subsets thereof for different stellar classifications. *Right:* the cumulative posterior probability for 73 stars, or subsets thereof for different stellar types. On the y-axis of the left hand plot, $p(P_s|\mathcal{D})$ represents the posterior distribution on P_s given the data. The product symbol represents the sum of the posteriors for individual stars i , for a total of N stars (which differs depending on the stellar type being shown).

our analysis after shifting the seismic ages younger by their systematic uncertainty. This resulted in 95.8% of the posterior probability to lie below $P_s = 0.5$, compared to 98.4% for the unaltered ensemble. From this, we can infer that our conclusions are robust against potential issues in seismic analyses. For a full discussion of these tests and other biases, see **Methods**.

Weakened magnetic braking was first proposed by van Saders et al. (2016) [2, hereafter the ‘van Saders’ study] in response to stars with spot rotation rates faster than expected from gyrochronology at their asteroseismic ages. This discrepancy was also indicated at around the same time in other studies of asteroseismic ages of main sequence stars [9, 10, 11]. The theory of weakened magnetic braking has been both reinforced by recent studies [41], as well as disputed [42], at least at the critical Rossby number originally proposed.

A recent study using spectroscopic rotation measurements of 14 solar twins, compared to grids of stellar models, found marginal evidence favouring a standard rotational evolution [42]. If weakened magnetic braking were to take place, this work proposed that it would most likely be outside the range of their sample, at $Ro_{\text{crit}} \geq 2.29$, compared to the value of 1.97 used in this work. Our sample does not overlap with theirs, so we can not directly compare to their results. Our work does find statistical agreement with a model of weakened magnetic braking at a critical Rossby number of 1.97 over one without any braking at all, indicating that traces of weakened magnetic braking do occur at Rossby numbers of around 2. A comparison to models of different Rossby numbers will be explored in future papers, using the LEGACY and ‘Kages’ rotation sample presented in this work.

As a sanity check that the evidence in favour of weakened magnetic braking presented in this work is independently robust, we recreated the joint posterior probability for P_s , this time looking only at MS stars, and excluding any of the 22 stars that were included in the original van Saders study used to detect weakened magnetic braking. Of the 47 MS stars used to distinguish between stellar models, 16 were included in the van Saders sample. Both the 16 stars in the original van Saders study as well as the remaining MS stars favoured the WMB model, although the van Saders stars did so more strongly, not including any stars that favoured the standard model. Specifically, when considering only the van Saders stars, 96.3% of the total joint probability lay below $P_s = 0.5$, compared to 91.7% when considering only those stars *not* included in the van Saders study. The stars initially used to propose the WMB model are still those most strongly in favour of it even when using asteroseismic rotation rates only, a result which is now further reinforced by an expanded sample that includes quiescent stars.

Conclusions: In this work, we obtained asteroseismic rotation rates for 91 stars using a hierarchical Bayesian model to fit for oscillation frequencies, convective background, and rotational splitting simultaneously. We validated these results against previously published and unpublished asteroseismic rotation measurements, making our new rotation catalogue the largest self-consistent sample of asteroseismic rotation in main sequence stars to date. We evaluated the stars in our sample against synthetic population models of the *Kepler* field evolved under either a ‘standard’ model of rotational evolution, or one where weakened magnetic braking takes place. This evaluation was done using a Bayesian mixture model to determine whether the ensemble preferred one model over another, and was done simultaneously in asteroseismic rotation, mass, age, and spectroscopic temperature and metallicity.

We leave the reader with the following conclusions:

1. We found that our ensemble was 98.4% more likely to be drawn from a weakened magnetic braking model than a standard model of rotational evolution [1], at a critical Rossby number of $Ro_{\text{crit}} = 1.97$. This work expands upon previous analysis [18] by including quiescent stars older than the Sun. This conclusion was found to be robust against change of ensemble members and potentially underestimated asteroseismic systematic uncertainties. A comparison to other braking laws [17] and other gyrochronology relations [43] using our asteroseismic ensemble will take place in future work.
2. We compared our new asteroseismic rotation rates with surface rotation measures from spot modulation. Our findings replicate those in similar studies [10, 31], in the sense that we find no statistically significant difference between seismic and spot modulation measures of stellar rotation in our ensemble.
3. We present new techniques for asteroseismic ‘peak-bagging’ frequency analyses, specifically: expressing well known empirical asteroseismic relations in a hierarchical modeling structure to account for small-scale effects not included in our model (such as acoustic glitches), and replacing uninformative empirical relations with Gaussian Process priors, detailed in the **Supplementary Information**.

In the near future our new asteroseismic rotation catalogue will be further complemented with high-precision surface rotation measurements and atmospheric parameters from spectroscopic surveys such as LAMOST [44], 4MOST [45] and WEAVE [46], and new asteroseismic measurements of age from the K2 and *TESS* missions. With these surveys, large scale ensemble asteroseismology will continue to increase the possibilities for understanding gyrochronology.

References

- [1] van Saders, J. L. & Pinsonneault, M. H. Fast Star, Slow Star; Old Star, Young Star: Subgiant Rotation as a Population and Stellar Physics Diagnostic. *The Astrophysical Journal* **776**, 67 (2013).
- [2] van Saders, J. L. *et al.* Weakened magnetic braking as the origin of anomalously rapid rotation in old field stars. *Nature* **529**, 181 (2016).
- [3] Barnes, S. A. Ages for Illustrative Field Stars Using Gyrochronology: Viability, Limitations, and Errors. *The Astrophysical Journal* **669**, 1167 (2007).
- [4] Meibom, S. *et al.* A spin-down clock for cool stars from observations of a 2.5-billion-year-old cluster. *Nature* **517**, 589 (2015).
- [5] Leiner, E., Mathieu, R. D., Vanderburg, A., Gosnell, N. M. & Smith, J. C. Blue Lurkers: Hidden Blue Stragglers on the M67 Main Sequence Identified from Their Kepler/K2 Rotation Periods. *The Astrophysical Journal* **881**, 47 (2019).

- [6] Claytor, Z. R. *et al.* Chemical Evolution in the Milky Way: Rotation-based ages for APOGEE-Kepler cool dwarf stars. *arXiv e-prints* arXiv:1911.04518 (2019).
- [7] Borucki, W. J., Koch, D. & Team, K. S. Kepler Planet Detection Mission: Highlights of the First Results. *AAS/Division for Planetary Sciences Meeting Abstracts #42* **42**, 47.03 (2010).
- [8] Silva Aguirre, V. *et al.* Ages and fundamental properties of Kepler exoplanet host stars from asteroseismology. *Monthly Notices of the Royal Astronomical Society* **452**, 2127 (2015).
- [9] Angus, R., Aigrain, S., Foreman-Mackey, D. & McQuillan, A. Calibrating gyrochronology using Kepler asteroseismic targets. *Monthly Notices of the Royal Astronomical Society* **450**, 1787 (2015).
- [10] Nielsen, M. B., Schunker, H., Gizon, L. & Ball, W. H. Constraining differential rotation of Sun-like stars from asteroseismic and starspot rotation periods. *Astronomy and Astrophysics* **582**, A10 (2015).
- [11] Davies, G. R. *et al.* Asteroseismic inference on rotation, gyrochronology and planetary system dynamics of 16 Cygni. *Monthly Notices of the Royal Astronomical Society* **446**, 2959 (2015).
- [12] Réville, V., Brun, A. S., Matt, S., Strugarek, A. & Pinto, R. The effect of magnetic topology on thermally-driven winds: Towards a general formulation of the braking law. *The Astrophysical Journal* **798**, 116 (2015). 1410.8746.
- [13] Garraffo, C., Drake, J. J. & Cohen, O. The Missing Magnetic Morphology Term in Stellar Rotation Evolution. *Astronomy & Astrophysics* **595**, A110 (2016). 1607.06096.
- [14] Metcalfe, T. S. *et al.* LBT/PEPSI Spectropolarimetry of a Magnetic Morphology Shift in Old Solar-type Stars. *arXiv:1912.01186 [astro-ph]* (2019). 1912.01186.
- [15] See, V. *et al.* Do Non-dipolar Magnetic Fields Contribute to Spin-down Torques? *The Astrophysical Journal* **886**, 120 (2019).
- [16] McQuillan, A., Mazeh, T. & Aigrain, S. Rotation Periods of 34,030 Kepler Main-Sequence Stars: The Full Autocorrelation Sample. *The Astrophysical Journal Supplement Series* **211**, 24 (2014). 1402.5694.
- [17] Matt, S. P., Brun, A. S., Baraffe, I., Bouvier, J. & Chabrier, G. The Mass-dependence of Angular Momentum Evolution in Sun-like Stars. *The Astrophysical Journal* **799**, L23 (2015).
- [18] van Saders, J. L., Pinsonneault, M. H. & Barbieri, M. Forward Modeling of the Kepler Stellar Rotation Period Distribution: Interpreting Periods from Mixed and Biased Stellar Populations. *The Astrophysical Journal* **872**, 128 (2019).
- [19] Ledoux, P. The Nonradial Oscillations of Gaseous Stars and the Problem of Beta Canis Majoris. *The Astrophysical Journal* **114**, 373 (1951).

- [20] Davies, G. R. *et al.* Oscillation frequencies for 35 Kepler solar-type planet-hosting stars using Bayesian techniques and machine learning. *Monthly Notices of the Royal Astronomical Society* **456**, 2183 (2016).
- [21] Lund, M. N. *et al.* Standing on the Shoulders of Dwarfs: The Kepler Asteroseismic LEGACY Sample. I. Oscillation Mode Parameters. *The Astrophysical Journal* **835**, 172 (2017).
- [22] Silva Aguirre, V. *et al.* Standing on the Shoulders of Dwarfs: The Kepler Asteroseismic LEGACY Sample. II. Radii, Masses, and Ages. *The Astrophysical Journal* **835**, 173 (2017).
- [23] García, R. A. *et al.* Rotation and magnetism of Kepler pulsating solar-like stars. Towards asteroseismically calibrated age-rotation relations. *Astronomy and Astrophysics* **572**, A34 (2014).
- [24] Bedding, T. R. *et al.* Solar-like Oscillations in Low-luminosity Red Giants: First Results from Kepler. *The Astrophysical Journal Letters* **713**, L176–L181 (2010).
- [25] Kraft, R. P. Studies of Stellar Rotation. V. The Dependence of Rotation on Age among Solar-Type Stars. *The Astrophysical Journal* **150**, 551 (1967).
- [26] Bonanno, A. & Fröhlich, H.-E. A Bayesian estimation of the helioseismic solar age. *Astronomy and Astrophysics* **580**, A130 (2015).
- [27] Paxton, B. *et al.* Modules for Experiments in Stellar Astrophysics (MESA): Convective Boundaries, Element Diffusion, and Massive Star Explosions. *arXiv:1710.08424 [astro-ph]* (2017). 1710.08424.
- [28] Hunter, J. D. Matplotlib: A 2D Graphics Environment. *Computing in Science and Engineering* **9**, 90–95 (2007).
- [29] Benomar, O. *et al.* Asteroseismic detection of latitudinal differential rotation in 13 Sun-like stars. *Science* **361**, 1231 (2018).
- [30] Lund, M. N., Miesch, M. S. & Christensen-Dalsgaard, J. Differential rotation in main-sequence solar-like stars: Qualitative inference from asteroseismic data. *The Astrophysical Journal* **790**, 121 (2014). 1406.7873.
- [31] Benomar, O., Takata, M., Shibahashi, H., Ceillier, T. & García, R. A. Nearly uniform internal rotation of solar-like main-sequence stars revealed by space-based asteroseismology and spectroscopic measurements. *Monthly Notices of the Royal Astronomical Society* **452**, 2654–2674 (2015).
- [32] Gizon, L. *et al.* Seismic constraints on rotation of Sun-like star and mass of exoplanet. *Proceedings of the National Academy of Science* **110**, 13267 (2013).
- [33] Chaplin, W. J. *et al.* Asteroseismic Determination of Obliquities of the Exoplanet Systems Kepler-50 and Kepler-65. *The Astrophysical Journal* **766**, 101 (2013).

- [34] Nielsen, M. B., Gizon, L., Schunker, H. & Karoff, C. Rotation periods of 12 000 main-sequence Kepler stars: Dependence on stellar spectral type and comparison with $v \sin i$ observations. *Astronomy and Astrophysics* **557**, L10 (2013).
- [35] Skumanich, A. TIME SCALES FOR Ca n EMISSION DECAY, ROTATIONAL BRAKING, AND LITHIUM DEPLETION. *ApJ*. . . 4 (1972).
- [36] Kawaler, S. D. Angular momentum loss in low-mass stars. *The Astrophysical Journal* **333**, 236 (1988).
- [37] Girardi, L. *et al.* TRILEGAL, a TRIdimensional modeL of thE GALaxy: Status and Future. *Astrophysics and Space Science Proceedings* **26**, 165 (2012).
- [38] Berger, T. A. *et al.* The \$Gaia\$-\$Kepler\$ Stellar Properties Catalog I: Homogeneous Fundamental Properties for 186,000 \$Kepler\$ Stars. *arXiv:2001.07737 [astro-ph]* (2020). 2001.07737.
- [39] Gelman, A. & Rubin, D. B. Inference from Iterative Simulation Using Multiple Sequences. *Statistical Science* **7**, 457–472 (1992).
- [40] Salvatier, J., Wiecki, T. V. & Fonnesbeck, C. Probabilistic programming in Python using PyMC3. *PeerJ Computer Science* **2**, e55 (2016).
- [41] Metcalfe, T. S. & Egeland, R. Understanding the Limitations of Gyrochronology for Old Field Stars. *The Astrophysical Journal* **871**, 39 (2019). 1811.11905.
- [42] Lorenzo-Oliveira, D. *et al.* Constraining the evolution of stellar rotation using solar twins. *Monthly Notices of the Royal Astronomical Society* **485**, L68 (2019).
- [43] Barnes, S. A. A Simple Nonlinear Model for the Rotation of Main-sequence Cool Stars. I. Introduction, Implications for Gyrochronology, and Color-Period Diagrams. *The Astrophysical Journal* **722**, 222 (2010).
- [44] Deng, L.-C. *et al.* LAMOST Experiment for Galactic Understanding and Exploration (LEGUE) — The survey’s science plan. *Research in Astronomy and Astrophysics* **12**, 735–754 (2012).
- [45] de Jong, R. S. *et al.* 4MOST: 4-metre Multi-Object Spectroscopic Telescope **9147**, 91470M (2014).
- [46] Dalton, G. *et al.* Project overview and update on WEAVE: The next generation wide-field spectroscopy facility for the William Herschel Telescope **9147**, 91470L (2014).

Acknowledgments: The authors would like to thank Sean Matt, Ellis Avallone, Alex Dixon, Warrick Ball and Brett Morris for helpful discussions. OJH, GRD and WJC acknowledge the support of the UK Science and Technology Facilities Council (STFC). JvS acknowledges support from the TESS Guest Investigator Program (80NSSC18K18584). MBN acknowledges support from the UK Space Agency This work has received funding from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (CartographY GA. 804752) Funding for the Stellar Astrophysics Centre is provided by The Danish National Research Foundation (Grant agreement no.: DNR106). LA, AB and VS acknowledge funding from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation program (grant agreement No. 682393 AWESoMeStars). AB also acknowledges the support of the College of Engineering, Mathematics and Physical Sciences at the University of Exeter. RAG acknowledges the support from the PLATO and GOLF CNES grants. JT acknowledges that support for this work was provided by NASA through the NASA Hubble Fellowship grant #51424 awarded by the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., for NASA, under contract NAS5-26555. The computations described in this paper were performed using the University of Birmingham’s BlueBEAR HPC service. This paper includes data collected by the Kepler mission and obtained from the MAST data archive at the Space Telescope Science Institute (STScI). Funding for the Kepler mission is provided by the NASA Science Mission Directorate. STScI is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5–26555.

Author Contributions: O.J.H. led the project, with help from G.R.D., J.V.S., M.B.N and W.J.C.. J.V.S. also led the development of the stellar population models. M.N.L., R.A.G. and S.K. provided data or stellar models and along with J.T. assessed the validity of our asteroseismic results. L.A., A.B. and V.S. provided the assessment of the theoretical implications of the gyrochronology results. All authors have contributed to the interpretation of the data and the results, and all discussed and provided comments for all drafts of the paper.

Competing Interests: The authors declare that they have no competing financial interests.

Correspondence: Correspondence and requests for materials should be addressed to O.J.H.. (email: oliver.hall@esa.int).

Methods

Contents

1	Asteroseismic Methods	15
1.1	Asteroseismic Data	15
1.2	Mode Frequency Fitting	16
2	Gyrochronology Methods	17
2.1	Stellar Models	17
2.2	Bayesian Mixture Model	18
3	Verifying asteroseismic results	20
3.1	Priors on rotational parameters	20
3.2	Comparisons to previous studies	21
3.3	Seismic vs spectroscopic rotation	22
3.4	Asteroseismic detection biases	23
4	Verifying consequences for gyrochronology	24
4.1	Limits of our stellar models	24
4.2	Systematic uncertainties from asteroseismology	25
4.3	Binaries and Planet Hosts	25

1 Asteroseismic Methods

1.1 Asteroseismic Data

For our asteroseismic power spectrum data we used the unweighted power spectra from the KASOC pipeline [1]⁵. We did not apply any additional treatment to these data. For 16 Cyg A & B (KIC 12069424 and KIC 12069449) we used the KEPSEISMIC lightcurves [2]⁶, which have significantly better signal-to-noise for these two stars.

For our asteroseismic ages, we used the ages obtained by **BASTA** [3, BAYesian STellar Algorithm] in the ‘Kages’ and LEGACY catalogues. These ages have been obtained by comparisons of measured oscillation properties to stellar models, accounting for an expanded range of metallicities. **BASTA** is thoroughly compared to four other seismic modelling techniques in [4]. While uncertainties found through **BASTA** are typically higher than for other techniques, only **BASTA** and **ASTFIT** [5, Aarhus STellar Evolution Code] recover the radius, mass and age of the Sun, when applied to solar data. Although the uncertainties on **ASTFIT** ages are overall lower, they are not published for the ‘Kages’ sample. In order to maintain an internally consistent stellar age sample, we used age results from **BASTA** for both the ‘Kages’ and LEGACY samples.

For our stellar masses we used asteroseismic model masses obtained by **BASTA** reported in ‘Kages’ and LEGACY, in order to maintain internal consistency with the age measurements.

⁵Obtainable from the KASOC webpage.

⁶Obtainable from MAST.

We note that age and mass posteriors from **BASTA** are correlated, but chose not to account for the unpublished correlations in this work.

As described in the catalogue papers, for ‘Kages’ stars atmospheric properties (T_{eff} and $[\text{Fe}/\text{H}]$) were measured through high-resolution spectroscopy [6]. For LEGACY stars, atmospheric properties were similarly taken from one study [7] for most stars in the catalogue, and complemented by other values from the literature for the remaining stars [4, see Table 3].

Other asteroseismic properties used in this study, such as those used as first guesses on the free parameters in our asteroseismic model, were taken from the ‘Kages’ and LEGACY catalogues [8, 9, 10]. In cases where both catalogues contained the same target, we used the stellar parameters reported in LEGACY [11]. This is also the case for the masses and ages described above.

1.2 Mode Frequency Fitting

In order to extract signatures of stellar rotation from the asteroseismic mode frequencies, we built a model that simultaneously treats the convective background, modes of oscillation, and white noise, while accounting for rotational splitting. The foundations of this approach follows best-practices in asteroseismology [12], fitting Harvey profiles for the background [13], and expressing the frequencies of the modes using an asymptotic expression [14, 15].

The first core improvement in this work come in the form of the use of hierarchical latent variable models [16, 17], which account for small-scale deviations in mode frequencies due to effects not explicitly accounted for in our model (such as acoustic glitches [18]). By improving the inference of mode frequencies in this way, we also improve the ability to resolve the rotational splitting. The second improvement comes in the form of priors on our parameters, and in particular the rotational inclination, which more accurately reflects the true distribution of angles than previous techniques [19]. A step-by-step breakdown of the model and all priors can be found in **Supplementary Materials**.

We separated our model fitting into two parts. First, we fit a model for the convective background and white noise only, to the region of the power spectra that did not contain modes of oscillation. This was done using **PyStan** [20, 21], for 10,000 iterations on each star⁷. Second, we fit our full model (including the convective background) to the region containing the modes of oscillation. The results of the first fit to the convective background were used as extremely informative priors on the background parameters in this second fit. This was done using **PyMC3** [22, 23, 24] for 2500 iterations each on 4 chains.

From this asteroseismic analysis, we report inclination angle (i), rotational splitting (ν_s), and rotation period (P) in **Extended Data Table 1**. The summary statistics on these parameters were taken as the median of the posterior distribution, with uncertainties being the 15.9th and 84.1st percentiles. For inclination angle ($i = \arccos(\cos(i))$) and rotation ($P = 1/\nu_s$), the full posterior samples were transformed before taking the summary statistics, as our model sampled in $\cos(i)$ and ν_s .

We flagged any sub-optimal conditions of the final fit. Of 94 stars, we flagged 5 for which the Gelman-Rubin convergence metric, \hat{R} , was greater than 1.01 and 2 stars for which it was

⁷These **PyStan** runs were initiated with a random seed of 11, as were all other random processes in this work

greater than 1.1 on inclination angle or rotation, where a $\hat{R} = 1$ indicates a converged result [25]. We also performed visual checks of the sampled chains on all hyperparameters and of the best-fit model compared to both the raw and smoothed asteroseismic data. We found no issues in the visual investigation of the 94 stars. KIC 8478994 was excluded due to both a poor fit as well as high \hat{R} on rotational parameters. KICs 6603624, 8760414 and 8938364 are reported in **Extended Data** Table 1, but were excluded from the gyrochronology analysis due to strong disagreement with multiple studies in the literature (see main body of paper). Finally, we flagged any stars with fewer than 1000 effective samples (n_{eff}) of ν_s .

2 Gyrochronology Methods

2.1 Stellar Models

The braking models used in this work have several parameters that inform the rotational evolution. These are: a normalization factor to reproduce the solar rotation (f_k); a disk locking timescale (T_{disk}) and period (P_{disk}) which together regulate the stellar angular velocity during the pre-main sequence; the critical angular velocity that marks the transition from saturated (rapidly rotating) to unsaturated (slowly rotating) regimes (ω_{crit}), and the critical Rossby number, above which stars conserve angular momentum (Ro_{crit}), mentioned above. T_{disk} , P_{disk} and ω_{crit} are calibrated to match the behavior in young open clusters, but have little impact on the rotational evolution beyond ≈ 1 Gyr in solar-mass stars. Both f_k and Ro_{crit} affect the late-time evolution. Both models adopt $\omega_{\text{crit}} = 3.4 \times 10^{-5} \text{ s}^{-1}$, $P_{\text{disk}} = 8.1$, $T_{\text{disk}} = 0.28$ and $f_k = 6.6$. In the weakened magnetic braking model, $Ro_{\text{crit}} = 1.97$. For further details, see [26] and [27, 28] for the standard and weakened magnetic braking cases respectively.

To construct a synthetic population of rotating stars, we expanded upon previous forward-models of the *Kepler* field with the purpose of studying weakened magnetic braking [28], with some important improvements. We started with a TRILEGAL [29] Milky-Way simulation of the *Kepler* field, using the simulation’s standard population values intended for this purpose. In order to replicate the *Kepler* selection effect, the TRILEGAL simulation was matched to the largest current catalogue of temperatures, luminosities and 2MASS *K*-band magnitudes of stars in the *Kepler* field [30]. This was done using a nearest-neighbours approach, based on the density of stars on the HR diagram. This ensured that the TRILEGAL population matched the density of stars actually observed by *Kepler*, replicating its selection effects, and improved upon previous efforts [28] by incorporating contemporary *Gaia* mission data [31].

We made further changes to account for possible binarity in the the matching sample [30]. If the first step is performed blindly, blended binaries in the sample cause an overestimation of the number of old stars. In order to overcome this, we:

1. blended the TRILEGAL stars with binary companions drawn from a flat mass-ratio distribution, using a known binary fraction [32],
2. recalculated the ‘observed’ luminosities and magnitudes assuming that each binary pair was blended, and
3. shifted these stars’ temperatures following a g - K magnitude relation [30].

This new distribution was used for the nearest-neighbour matching. Once drawn we dropped the binary companion and used the true TRILEGAL properties of those stars. For stars where the mass of the companion was $M < 0.4 M_{\odot}$, binary contributions were ignored. Every binary was assumed to result in a blend, regardless of separation. This results in slightly more young stars than reality, because young, blended binary systems contaminate regions of the HR diagram where one expects to find old stars, and the number of blends is overestimated by assuming every binary system is a blend.

Our asteroseismic sample of stars with short cadence observations are subject to additional selection functions not included in the creation of the model populations above. We did *not* explicitly account for these asteroseismic selection functions in our model, by design. Both the standard and WMB models contain stars with the same fundamental parameters (mass, radius, effective temperature, metallicity) but a different period based on the choice of rotational evolution prescription. Applying an asteroseismic selection function that depends on these fundamental parameters would apply an identical prior to both models, therefore providing no net effect on our posterior distribution [33]. Additionally, we expected any seismic selection function to be relatively flat (and therefore uninformative) on a star-by-star scale, on which we run our model analysis.

2.2 Bayesian Mixture Model

In order to determine whether weakened magnetic braking occurs on the main sequence, we compared our sample of seismic age and rotation, along with temperature, metallicity and mass, to the two stellar population models of the *Kepler* field [28], discussed above. Both stellar models were evaluated in a Bayesian framework, with the rationale of determining which of the two models (standard or WMB) is most likely to reproduce our observed data. Each model sample contained temperature (T_{eff}), mass (M), age (t), metallicity ($[\text{Fe}/\text{H}]$) and rotation (P) information.

In order to draw probabilistic inference about the models, we built a five-dimensional Kernel Density Estimate (KDE) of both model populations using the `statsmodels` package [34]. We used a band-width (setting the resolution of the KDE) of $0.02 M_{\odot}$ in mass, 10 K in T_{eff} , 0.01 dex in $\ln(t)$, 0.01 dex in $[\text{Fe}/\text{H}]$ and 0.01 dex in $\ln(P)$. Note that age and rotation were treated in log space, where the posterior estimates from asteroseismology more closely resemble normal distributions. This approach translates the population models to a probability distribution we can use in a Bayesian framework.

We evaluated our data against both models simultaneously by treating the data as being drawn from a mixture of both model KDEs. In this mixture model structure, the two KDEs were modulated by a weighting factor, P_s . In the limit $P_s \rightarrow 1$, the data are most likely drawn from the standard model. In the limit $P_s \rightarrow 0$, the data are most likely drawn from the WMB model.

The posterior probability of obtaining P_s and additional parameters θ given our data \mathcal{D} is $p(P_s, \theta | \mathcal{D})$. Using Bayes equation, we can express this as:

$$p(P_s, \theta | \mathcal{D}) \propto p(\mathcal{D} | \theta) p(\theta | P_s, \kappa_s, \kappa_{\text{WMB}}) p(P_s), \quad (1)$$

where κ_s and κ_{WMB} are the KDE functions for the standard and WMB models respectively, and θ here are parameters, $\theta = \{M, T_{\text{eff}}, \ln(t), [\text{Fe}/\text{H}], \ln(P)\}$. As done above for the astero-

seismic model, the parameters θ may be referred to as latent parameters, as they form a step between the parameter we want to infer (P_s) and our data. Using this approach allowed our model to properly take into account the observational uncertainties on the data.

The second component on the right hand side of Equation 1 describes the probability of obtaining our latent parameters θ given our KDEs and the mixture model weighting parameter P_s , and is described by the mixture model

$$p(\theta|P_s, \kappa_s, \kappa_{\text{WMB}}) = P_s \times \kappa_s(\theta) + (1 - P_s) \times \kappa_{\text{WMB}}(\theta), \quad (2)$$

where all parameters are as described above. This probability function describes a distribution that is a mixture of both KDEs. While the KDEs are constant, P_s is a free parameter, and so the shape of this distribution can vary. The latent parameters θ are drawn from this distribution, and therefore from some combination of the two stellar models.

The first component in Equation 1 describes the likelihood of obtaining the parameters θ given our data and their observational uncertainty. It takes the form

$$p(\mathcal{D}|\theta) = \mathcal{N}(\mathcal{D}|\theta, \sigma_{\mathcal{D}}), \quad (3)$$

a normal distribution evaluating the latent parameters θ against the observations, with observational uncertainty $\sigma_{\mathcal{D}}$. This approach means that in each parameter space (such as age), the age drawn from the stellar model mixture is entered into the likelihood equation with our static observations. The value of this equation (and thus the likelihood) will increase if θ is closer to the observations, and the mixture model will be modulated in a manner that maximises this probability, inferring whether one stellar model is more likely to produce our data than the other.

The final term, $p(P_s)$, represents the prior on the mixture model weight, which is uniform between 0 and 1. A visual representation of our model is shown in **Extended Data Figure 1**.

Typically, this model would evaluate all stars in our sample against the stellar models simultaneously for a single posterior estimate of P_s . At 95 stars, in 5 parameter spaces, this totals 476 free parameters to marginalise over. This is not an issue for Hamiltonian Monte Carlo [35, HMC], however the use of KDE functions, over which a probabilistic gradient can not be measured, reduces HMCs effectiveness. Alternative Markov Chain Monte Carlo techniques [36, MCMC] can more efficiently sample the KDE functions, but can not treat the large number of hierarchical parameters. To overcome this, we fit our model to each star to obtain an independent individual posterior distribution for P_s , and multiplied these post hoc to obtain a combined posterior. This comes with the benefit of easily allowing us to calculate the combined posterior for different stellar classifications, at the expense of the ability to marginalise for a single value of P_s directly.

The parameter space of the stellar models populations were reduced before calculating the KDEs. These cuts were made in M , T_{eff} , $\ln(t)$ and $[\text{Fe}/\text{H}]$, removing any stars in the models that fell more than $3 \times \sigma_{\mathcal{D}}$ outside the observations. Our observables M , t and P have asymmetric uncertainties from the Bayesian asteroseismic analysis. We used the larger of the reported uncertainties on each parameter as $\sigma_{\mathcal{D}}$ in each parameter space.

KIC 6278762 was excluded from this stellar model analysis, because its age fell more than 3σ outside of the highest age in the stellar models⁸, and KICs 7106245 and 8760414 were excluded for the same reason due to low metallicities outside the functional range of the stellar models (-0.99 and -0.92 respectively). We also excluded any stars with $n_{\text{eff}} < 1000$ for rotational splitting, and those with $\hat{R} > 1.1$. The remaining sample of 73 stars contained 4 sub-giants, 22 ‘hot’ stars, and 47 main sequence stars.

We fit our model Equation 1 using `emcee` [36, 37], using 32 walkers for a total of 7500 samples per walker, of which the first 2500 were discarded as a burn-in.

After fitting, we took a normalised histogram of the posterior estimate of P_s for each star, using 100 bins. In this histogram, each bin approximated the value of the posterior function for P_s . The array of 100 bins for all stars were multiplied, resulting in an approximation of the joint posterior probability function for P_s given all stars in the ensemble.

3 Verifying asteroseismic results

3.1 Priors on rotational parameters

In our Bayesian analysis, we have placed weakly informative priors on our sampled rotational parameters, $\nu_s \sin(i)$ and $\cos(i)$. The prior is especially important for the angle of inclination, which is hardest to infer from the data. We are able to validate the robustness of our asteroseismic results by confirming that their posterior distributions are data-dominated, and not prior-dominated. We can do so by comparing the 68% credible regions of the posterior estimates of $\nu_s \sin(i)$ and i against the 68% credible regions of their priors.

A comparison between prior and posterior is shown for 94 stars in $\nu_s \sin(i)$, i and P_{rot} in **Extended Data Figure 2**, arranged by age. In the Figure, results with means and error bars that closely resemble the prior distribution can be interpreted as prior-dominated (i.e. poorly informed by the data). The projected splitting, $\nu_s \sin(i)$, is overall well constrained, with only one star appearing prior dominated. This is expected, as the projected splitting is what we observe on the star before decoupling inclination and rotation. The angle of inclination i , sampled as $\cos(i)$, more closely follows the prior distribution in most cases. Combining the two, the rotation period P_{rot} has no stars directly corresponding to the effective prior on period, and globally follows a trend with increasing age. The three outliers with fast rotation at late ages (KICs 6603624, 8760414 and 8938364) are discussed in more detail in below.

The rotation rates as presented in this work are a product of our Bayesian sampling of both projected splitting and angle of inclination. As seen in the Figure there are instances where i or $\nu_s \sin(i)$ closely resemble the prior (and are therefore prior-dominated). There are no cases of this when looking at the resulting period measurements, as they will have been informed by at least one strongly data-driven parameter (judging from the Figure, commonly $\nu_s \sin(i)$). From this, we conclude that our ensemble of asteroseismic rotation is not strongly dominated by the priors imposed on projected splitting and inclination in our Bayesian analysis.

⁸This is a metallicity issue, as the oldest stars have metallicities outside the range of the rotational model grids.

3.2 Comparisons to previous studies

In order to validate our results, we compared our rotational parameters to those obtained in the literature, as well as those resulting from the work presented in LEGACY and Kages, which were unreported and received through private communication by the authors of the catalogue papers.

Comparisons with LEGACY and ‘Kages’ are shown in **Extended Data** Figure 3 for projected splitting, inclination angle, and rotation period. In all three cases we show the fractional difference between the values obtained in this work and those from LEGACY and Kages. On the right of the Figure, we show the distribution of the fractional differences for the three parameters. The projected splitting is in good agreement with both studies, however LEGACY finds slightly lower $\nu_s \sin(i)$ for the faster rotators, deviating from our work by over 1σ . Neither LEGACY nor ‘Kages’ used a spatially isotropic prior for the inclination angle in their analyses, instead opting for a uniform prior. As posterior estimates of inclination angle are only loosely data-driven, the introduction of an isotropic prior should result in our analysis reporting globally higher inclination angles. This effect is seen in the comparisons of both inclination angles and rotation rates for the LEGACY stars, where we find overall lower rotation rates compared to LEGACY for stars at very similar $\nu_s \sin(i)$.

A number of stars are excluded from **Extended Data** Figure 3 and compared individually as extreme outliers: KICs 5094751, 6196457, 8349582, 8494142, 8554498, 105114430 and 11133306 all have fast rotation rates (< 5 days) in Kages, but are found to have a broader spread of rotation rates in this work. At similar values for $\nu_s \sin(i)$, ‘Kages’ found much lower inclination angles with highly asymmetrical uncertainties. Based on a comparison between the summary statistics of these stars, we conclude that the results found in this work have better marginalised over inclination angle, improving our measure of rotation.

Conversely, KICs 6603624, 8760414 and 8938364 have extremely slow rotation periods in LEGACY, but extremely *fast* (< 3 days) in this work. KICs 8760414 and 8938364 are excluded from the gyrochronology analysis below based on checks for \hat{R} and the number of effective samples. Both stars have ages greater than 10 Gyr, making their fast rotation rates highly unlikely under any model of rotational evolution. The posterior estimate of P for KIC 6603624 is well-defined in our analysis, but with an age of 7.8 Gyr, its measured rotation of 1.2 days is also highly unlikely under any model of rotational evolution. The LEGACY estimate of rotation is similarly extreme at 378 days. These three stars have the lowest inclination angles in our sample ($< 10^\circ$), at which point the power in the split components of the seismic modes is so low that it becomes difficult to probe the measure of splitting. The split components for these stars will have a height roughly 3% of the central mode. For comparison, for the next lowest inclined star at 17° , this rises to 10%. In cases such as these with lower signal-to-noise, spurious peaks may be interpreted as split components.

We also compared our asteroseismic estimates of stellar rotation with similar studies in the literature, shown in **Extended Data** Figure 4. These included: a study of the binary solar analogues 16 Cyg A & B [12], ; a study of surface and seismic rotation which our catalogue shares 5 stars with [38], ; and an asteroseismic study of differential rotation with which our catalogue shares 40 targets [39]. For the latter, we used their reported splitting value a_1 , which is equivalent to ν_s .

Overall, **Extended Data** Figure 4 shows no strong disagreements between our asteroseismic measurements for stellar rotation and those from the literature. The scatter of the fractional differences lies cleanly around the zero line, with a mean and spread of $0.0^{+16.4}_{-15.6}$ %. The increase in uncertainty with period is due to more slowly rotating stars being more difficult to constrain using asteroseismology.

It is of note that 16 Cyg A was found to be rotating slightly faster in our analysis compared to the literature [12] (deviating within 2σ), despite the fit being performed on the same data. We found an inclination angle that is slightly lower for 16 Cyg A but at a similar projected splitting, which would explain finding a lower value of rotation.

There are three outliers at low period in **Extended Data** Figure 4: KICs 6603624, 8760414 and 8938364. These are the same targets found to be outliers in a comparison to the LEGACY and ‘Kages’ measurements (see above), with anomalously fast rotation rates and low inclination angles. As these stars represent the lowest inclination angles in our sample, and were found to disagree with two independent studies, we opted to exclude them from the gyrochronology analysis, and to flag the rotation measurements for these three stars presented in this work.

3.3 Seismic vs spectroscopic rotation

A distinct difference between rotation rates obtained through different techniques may hold information about differential rotation (both latitudinal and radial) of near-surface layers, such as those we observe in the Sun [40]. A previous comparison of spectroscopic and seismic rotation rates, performed on a sample of 22 stars, found no significant radial differential rotation [39]. In this work they not only considered rotation rates from spots, but also spectroscopic measures of the projected surface rotation $v \sin(i)$, which they found to be more reliable.

With our expanded sample of asteroseismic rotation we can perform a similar analysis, to both validate our sample and probe radial differential rotation. **Extended Data** Figure 5 shows a comparison between spectroscopic $v \sin(i)$ measurements as listed in LEGACY and ‘Kages’ (left) and Benomar et al. (2015) [41] (right). In these cases the asteroseismic $v \sin(i)$ has been calculated using our measure of asteroseismic $\nu_s \sin(i)$ and the known asteroseismic radii. Three stars (KICs 6603624, 8760414 and 8938364) have been excluded from this figure due to strong disagreements of measured rotation rates with the literature (see above).

For the LEGACY and ‘Kages’ sample, we find no strong deviation from the 1:1 line except at very low velocities, which is likely due to biases inherent to spectroscopic line broadening measurements [42, 43]. For the Benomar et al. (2015) sample stars lie a lot closer to the 1:1 line.

Overall, there appears to be a global offset where spectroscopic measurements of projected rotation appear faster than asteroseismic measures. Based on the LEGACY and ‘Kages’ $v \sin(i)$ values, this offset is roughly 18% (i.e. spectroscopic projected rotation rates are faster than asteroseismic rates). This offset is much smaller ($\sim 5\%$) for the [41] sample, albeit for far fewer stars. These offsets are within the typical disagreement between spectroscopic methods, based on comparisons of projected rotation measurements for red giant stars [43, see Figure 2], especially at $< 5 \text{ km s}^{-1}$.

It is worth addressing the impact this comparison would have on our conclusions for

gyrochronology, were we to take the spectroscopic rotation rates as if they were the truth, even at velocities of $< 5 \text{ km s}^{-1}$. For the majority of cases, the seismic velocities are slower than the spectroscopic velocities, which would bias our ensemble towards favouring a standard evolution over a weakened magnetic braking scenario (as stars would overall appear to be rotating slower at late ages). Based on the comparison to spot rotation presented in the main body of this paper, and given known issues comparing seismic and spectroscopic rotation at low velocities, we conclude that the divergence seen in **Extended Data** Figure 5 does not undermine our results.

3.4 Asteroseismic detection biases

Stars that are rotating very slowly will show small rotational splittings, which may be indistinguishable from a non-rotating case, whereas stars that are spinning very fast may find that their splitting is so wide that they cross over with split modes at different radial orders (n). Asteroseismic selection techniques may also introduce additional biases on a population level. We consider the limitations this places on our asteroseismic method.

- *Fast rotating stars:* In principle, a star could rotate fast enough that the highest-frequency peak in a split dipole ($\ell = 2$) mode would overlap with the radial ($\ell = 0$) mode. This would occur if $2\nu_s > \delta\nu_{02}$, the small separation. For the star in our sample with the smallest $\delta\nu_{02}$ (so the highest risk of this happening), this would occur at a rotation of 8.8 days or faster. For the star with the largest $\delta\nu_{02}$, this limit was roughly 2 days.

In practice, the model would still properly fit a spectrum where this kind of mode-overlapping occurred, as the $\ell = 0$ modes don't split, meaning that there are no degenerate solutions in this regime. A more serious issue may be when the $\ell = 1$ mode at lower frequencies overlapped with the $\ell = 2$ mode as well, which would only occur at a rotation rate of 2 days or faster for the star with the smallest $\delta\nu_{01}$.

- *Slow rotating stars:* Except for the most fast rotating cases, asteroseismic mode splitting is detected by a broadening of the mode frequencies as the split components begin to separate from the central mode. If there is to be a hard limit at which this measurement is not possible, it should be in the case that $\nu_s < 2/T$, where T is the observation time, and $1/T$ is the frequency bin width. For the star in our sample with the shortest observing time, this limit would be ~ 89 days, which quickly rises for longer observing baselines. While this constitutes a hard limit, slower rotation rates will become harder and harder to constrain as the rate of splitting becomes smaller. A benefit of the Bayesian approach used in this paper is that it should accurately reflect the increased uncertainty associated with the small rotational splitting.

While these two limits cause concern when studying the most extreme rotators, the results presented in this work are driven by stars with rotation rates in the regime of 10 to 40 days, and so we do not expect these limits to bias our conclusions.

Finally, we also consider bias in asteroseismic detections. The probability of an asteroseismic detection with *Kepler* scales with temperature and radius (i.e. scales inversely with

$\log(g)$, and therefore main sequence age) [33, 44]. More active, magnetic stars may also suppress their oscillation modes, making it more likely to detect seismic oscillations in older stars [45]. Through these two effects, asteroseismology is biased towards detecting older over younger stars. As it is older stars, undergoing magnetic braking, that are required to drive the distinction between stellar models, we do not expect this observing bias to effect our conclusions.

4 Verifying consequences for gyrochronology

4.1 Limits of our stellar models

The rotational stellar models used in this work [28] are constructed for metallicities of $-0.4 < [\text{Fe}/\text{H}] < 0.4$ dex, in steps of 0.1 dex. Our sample of 91 stars contained 4 stars with metallicities below -0.4 dex, which are shown as shaded symbols in Figure 3. Of these, 3 were included in our final sample of 73 stars used to evaluate our stellar models: KICs 7970740, 8684723 and KIC 9965715. All three are classed as MS stars, with metallicities of -0.54 ± 0.10 , -0.42 ± 0.10 and -0.44 ± 0.18 dex respectively, placing them within 3σ of the metallicity limits of our stellar models. KICs 8684723 and 9965715 strongly agree with the WMB model, whereas KIC 7970740 weakly prefers the standard model. Excluding these stars was not found to significantly alter the joint posterior distribution shown in Figure 4.

A recent study [46] compared different rotational evolution models [26, 47, of which we use the former in this work] while studying the effect metallicity has on rotation. They found that metal-rich stars spin down significantly more effectively than metal poor stars. The population of 73 stars used in our stellar model comparisons is roughly centered on a $[\text{Fe}/\text{H}]$ of 0 dex, with a spread of 0.16 dex, with no stars significantly above or below ± 0.4 dex (as discussed above). While differences in stellar rotational evolution as a function of metallicity in this region are still somewhat pronounced, they are much less so than for more metal-rich or poor stars [46, see Figure 2]. However, for stars with $[\text{Fe}/\text{H}] < 0$, the alternative model prescription [47] sees stars spin down more slowly than the models used in this work (i.e. they will rotate faster at later ages). This work only explores the presence of weakened magnetic braking for a single braking law, and a comparison to alternative braking model will be explored in a future paper.

When constructing KDEs from our stellar model samples (see Section 2), we selected fixed resolutions (or band-widths) for the KDEs. In mass, the band-width of $0.02 M_{\odot}$, was larger than the uncertainties of 24 of 73 stars used to construct our joint posterior. For the stars with the smallest uncertainties, this significantly limits the size of the KDE being evaluated (as subsections of the full stellar models are used to evaluate individual stars, for computational efficiency). In order to confirm that these stars do not significantly affect the ensemble’s preference towards the WMB model, we recalculated the joint posterior distribution for P_s , excluding stars with an uncertainty on mass smaller than the KDE band-width. While the 24 stars with small uncertainties do favour the WMB model, they do so very weakly, whereas the remaining 49 stars with larger uncertainties strongly favour the WMB model. Their removal from the total joint posterior probability does not significantly alter it from the distribution shown in Figure 4.

4.2 Systematic uncertainties from asteroseismology

In our model analysis, we used asteroseismic mass and age obtained using BASTA, as reported in LEGACY and Kages. Asteroseismic properties obtained through stellar models can be subject to systematic errors arising from differences in input physics and choice of stellar models not included in the reported statistical uncertainties. A quantification of these different systematic effects can be found in the ‘Kages’ catalogue paper [3]. Combining their reported median systematic uncertainties due to input physics results in median uncertainties of 20% (up from 14%) on age and 5% (up from 3%) on mass for the ‘Kages’ sample. For LEGACY, the median uncertainties are 18% (up from 10%) and 5.6% (up from 4%) for age and mass respectively.

We re-ran our model analysis after inflating uncertainties on mass and age. We increased uncertainties by the fractional difference between the BASTA statistical uncertainties and the median full statistical and systematic uncertainties described above. For example, a LEGACY star with a mass of $2.0 \pm 0.5 M_{\odot}$ would have its uncertainty inflated by 1.6% of its mass, to $0.53 M_{\odot}$.

To further test the limits of this analysis, we also reran our mixture model fit, this time only shifting the asteroseismic ages younger by the systematic uncertainty, and retaining the statistical uncertainty (i.e. the ages of LEGACY and ‘Kages’ stars were reduced by 8% and 6% respectively). In this scenario where all asteroseismic ages are overestimated, ‘true’ fast rotators at young ages would have been mistaken for fast rotators at old ages, suggesting the presence of weakened magnetic braking where none existed.

The results of these tests are discussed further in the main body of the paper.

4.3 Binaries and Planet Hosts

For stars to be good probes of existing gyrochronology relations, their rotational evolution must occur in isolation. If a star interacts with a close binary companion (through tides, or a merger) the natural angular momentum loss can be disturbed, causing gyrochronology to mispredict ages [48, 49]. Between LEGACY and Kages, we have 8 known binaries.

First, KIC 8379927, KIC 7510397, KIC 10454113 and KIC 9025370 are spectroscopic binaries. This does not affect the asteroseismic analysis, but may affect their rotational evolution. Of these, KICs 8379927, 7510397 and 9025370 were included in the gyrochronology analysis. None of them preferred one model strongly over the other, with all three finding flat posteriors for P_g .

Second, the binary pairs of KIC 9139151 & 9139163 and 16 Cyg A & B are individually observed binary components with wide orbital separations, so we do not expect their binarity to have affected their rotational evolution.

While we chose not to account for star-planet tidal interactions in this work, we note that this may also disturb the natural stellar rotational evolution when tidal forces are at play [50, 51, 52], although this has been disputed by observations of asteroseismic planet hosts [53].

References

- [1] Handberg, R. & Lund, M. N. Automated preparation of Kepler time series of planet hosts for asteroseismic analysis. *Monthly Notices of the Royal Astronomical Society* **445**, 2698–2709 (2014).
- [2] García, R. A. *et al.* Preparation of Kepler light curves for asteroseismic analyses. *Monthly Notices of the Royal Astronomical Society* **414**, L6–L10 (2011).
- [3] Silva Aguirre, V. *et al.* Ages and fundamental properties of Kepler exoplanet host stars from asteroseismology. *Monthly Notices of the Royal Astronomical Society* **452**, 2127 (2015).
- [4] Silva Aguirre, V. *et al.* Standing on the Shoulders of Dwarfs: The Kepler Asteroseismic LEGACY Sample. II. Radii, Masses, and Ages. *The Astrophysical Journal* **835**, 173 (2017).
- [5] Christensen-Dalsgaard, J. ASTEC—the Aarhus STellar Evolution Code. *Astrophysics and Space Science* **316**, 13 (2008).
- [6] Huber, D. *et al.* Stellar Spin-Orbit Misalignment in a Multiplanet System. *Science* **342**, 331 (2013).
- [7] Buchhave, L. A. & Latham, D. W. The Metallicities of Stars with and without Transiting Planets. *The Astrophysical Journal* **808**, 187 (2015).
- [8] Astropy Collaboration *et al.* Astropy: A community Python package for astronomy. *Astronomy and Astrophysics* **558**, A33 (2013).
- [9] Astropy Collaboration *et al.* The Astropy Project: Building an Open-science Project and Status of the v2.0 Core Package. *The Astronomical Journal* **156**, 123 (2018).
- [10] Ginsburg, A. *et al.* Astroquery: An Astronomical Web-querying Package in Python. *The Astronomical Journal* **157**, 98 (2019).
- [11] McKinney, W. Data Structures for Statistical Computing in Python. In *Proceedings of the 9th Python in Science Conference*, 51–56 (2010).
- [12] Davies, G. R. *et al.* Asteroseismic inference on rotation, gyrochronology and planetary system dynamics of 16 Cygni. *Monthly Notices of the Royal Astronomical Society* **446**, 2959 (2015).
- [13] Harvey, J. High-Resolution Helioseismology. *Future Missions in Solar, Heliospheric & Space Plasma Physics* **235**, 199 (1985).
- [14] Tassoul, M. Asymptotic approximations for stellar nonradial pulsations. *The Astrophysical Journal Supplement Series* **43**, 469 (1980).
- [15] Vrad, M., Mosser, B. & Samadi, R. Period spacings in red giants. II. Automated measurement. *Astronomy and Astrophysics* **588**, A87 (2016).

- [16] Hogg, D. W., Bovy, J. & Lang, D. Data analysis recipes: Fitting a model to data. *ArXiv e-prints* arXiv:1008.4686 (2010).
- [17] Hall, O. J. *et al.* Testing asteroseismology with Gaia DR2: Hierarchical models of the Red Clump. *Monthly Notices of the Royal Astronomical Society* **486**, 3569–3585 (2019). 1904.07919.
- [18] Mazumdar, A. *et al.* Measurement of acoustic glitches in solar-type stars from oscillation frequencies observed by Kepler. *The Astrophysical Journal* **782**, 18 (2014). 1312.4907.
- [19] Chaplin, W. J. & Basu. *Asteroseismic Data Analysis: Foundations and Techniques* (Princeton University Press, Princeton, New Jersey, 2017), 1st edn.
- [20] Van Hoey, S., van der Kwast, J., Nopens, I. & Seuntjens, P. Python package for model STructure ANalysis (pySTAN). *EGU General Assembly Conference Abstracts* **15**, EGU2013–10059 (2013).
- [21] Carpenter, B. *et al.* Stan: A Probabilistic Programming Language. *Journal of Statistical Software* **76**, 1–32 (2017).
- [22] van der Walt, S., Colbert, S. C. & Varoquaux, G. The NumPy Array: A Structure for Efficient Numerical Computation. *Computing in Science and Engineering* **13**, 22–30 (2011).
- [23] Salvatier, J., Wiecki, T. V. & Fonnesbeck, C. Probabilistic programming in Python using PyMC3. *PeerJ Computer Science* **2**, e55 (2016).
- [24] The Theano Development Team *et al.* Theano: A Python framework for fast computation of mathematical expressions. *arXiv:1605.02688 [cs]* (2016). 1605.02688.
- [25] Gelman, A. & Rubin, D. B. Inference from Iterative Simulation Using Multiple Sequences. *Statistical Science* **7**, 457–472 (1992).
- [26] van Saders, J. L. & Pinsonneault, M. H. Fast Star, Slow Star; Old Star, Young Star: Subgiant Rotation as a Population and Stellar Physics Diagnostic. *The Astrophysical Journal* **776**, 67 (2013).
- [27] van Saders, J. L. *et al.* Weakened magnetic braking as the origin of anomalously rapid rotation in old field stars. *Nature* **529**, 181 (2016).
- [28] van Saders, J. L., Pinsonneault, M. H. & Barbieri, M. Forward Modeling of the Kepler Stellar Rotation Period Distribution: Interpreting Periods from Mixed and Biased Stellar Populations. *The Astrophysical Journal* **872**, 128 (2019).
- [29] Girardi, L. *et al.* TRILEGAL, a TRIdimensional model of thE GALaxy: Status and Future. *Astrophysics and Space Science Proceedings* **26**, 165 (2012).
- [30] Berger, T. A. *et al.* The \$Gaia\$-\$Kepler\$ Stellar Properties Catalog I: Homogeneous Fundamental Properties for 186,000 \$Kepler\$ Stars. *arXiv:2001.07737 [astro-ph]* (2020). 2001.07737.

- [31] Gaia Collaboration *et al.* Gaia Data Release 2. Summary of the contents and survey properties. *ArXiv e-prints* arXiv:1804.09365 (2018).
- [32] Raghavan, D. *et al.* A Survey of Stellar Families: Multiplicity of Solar-type Stars. *The Astrophysical Journal Supplement Series* **190**, 1–42 (2010).
- [33] Chaplin, W. J. *et al.* Ensemble Asteroseismology of Solar-Type Stars with the NASA Kepler Mission. *Science* **332**, 213 (2011).
- [34] Seabold, S. & Perktold, J. Statsmodels: Econometric and Statistical Modeling with Python 5 (2010).
- [35] Betancourt, M. J. & Girolami, M. Hamiltonian Monte Carlo for Hierarchical Models. *arXiv:1312.0906 [stat]* (2013). 1312.0906.
- [36] Foreman-Mackey, D., Hogg, D. W., Lang, D. & Goodman, J. Emcee: The MCMC Hammer. *Publications of the Astronomical Society of the Pacific* **125**, 306–312 (2013). 1202.3665.
- [37] Foreman-Mackey, D. Corner.py: Scatterplot matrices in Python. *The Journal of Open Source Software* **1** (2016).
- [38] Nielsen, M. B., Schunker, H., Gizon, L. & Ball, W. H. Constraining differential rotation of Sun-like stars from asteroseismic and starspot rotation periods. *Astronomy and Astrophysics* **582**, A10 (2015).
- [39] Benomar, O. *et al.* Asteroseismic detection of latitudinal differential rotation in 13 Sun-like stars. *Science* **361**, 1231 (2018).
- [40] Beck, J. G. A comparison of differential rotation measurements - (Invited Review). *Solar Physics* **191**, 47–70 (2000).
- [41] Benomar, O., Takata, M., Shibahashi, H., Ceillier, T. & García, R. A. Nearly uniform internal rotation of solar-like main-sequence stars revealed by space-based asteroseismology and spectroscopic measurements. *Monthly Notices of the Royal Astronomical Society* **452**, 2654–2674 (2015).
- [42] Doyle, A. P., Davies, G. R., Smalley, B., Chaplin, W. J. & Elsworth, Y. Determining stellar macroturbulence using asteroseismic rotational velocities from Kepler. *Monthly Notices of the Royal Astronomical Society* **444**, 3592–3602 (2014).
- [43] Tayar, J. *et al.* Rapid Rotation of Low-mass Red Giants Using APOKASC: A Measure of Interaction Rates on the Post-main-sequence. *The Astrophysical Journal* **807**, 82 (2015).
- [44] Schofield, M. *et al.* The Asteroseismic Target List for Solar-like Oscillators Observed in 2 minute Cadence with the Transiting Exoplanet Survey Satellite. *The Astrophysical Journal Supplement Series* **241**, 12 (2019).

- [45] Mathur, S. *et al.* Revisiting the impact of stellar magnetic activity on the detection of solar-like oscillations by Kepler. *Frontiers in Astronomy and Space Sciences* **6**, 46 (2019). 1907.01415.
- [46] Amard, L. & Matt, S. P. The Impact of Metallicity on the Evolution of Rotation and Magnetic Activity of Sun-Like Stars. *arXiv:2001.10404 [astro-ph]* (2020). 2001.10404.
- [47] Matt, S. P., Brun, A. S., Baraffe, I., Bouvier, J. & Chabrier, G. The Mass-dependence of Angular Momentum Evolution in Sun-like Stars. *The Astrophysical Journal* **799**, L23 (2015).
- [48] Leiner, E., Mathieu, R. D., Vanderburg, A., Gosnell, N. M. & Smith, J. C. Blue Lurkers: Hidden Blue Stragglers on the M67 Main Sequence Identified from Their Kepler/K2 Rotation Periods. *The Astrophysical Journal* **881**, 47 (2019).
- [49] Fleming, D. P., Barnes, R., Davenport, J. R. A. & Luger, R. Rotation Period Evolution in Low-mass Binary Stars: The Impact of Tidal Torques and Magnetic Braking. *The Astrophysical Journal* **881**, 88 (2019).
- [50] Maxted, P. F. L., Serenelli, A. M. & Southworth, J. Comparison of gyrochronological and isochronal age estimates for transiting exoplanet host stars. *Astronomy and Astrophysics* **577**, A90 (2015).
- [51] Gallet, F. & Delorme, P. Star-planet tidal interaction and the limits of gyrochronology. *arXiv e-prints* **1905**, arXiv:1905.06070 (2019).
- [52] Benbakoura, M., Réville, V., Brun, A. S., Le Poncin-Lafitte, C. & Mathis, S. Evolution of star-planet systems under magnetic braking and tidal interaction. *Astronomy and Astrophysics* **621**, A124 (2019).
- [53] Ceillier, T. *et al.* Rotation periods and seismic ages of KOIs - comparison with stars without detected planets from Kepler observations. *Monthly Notices of the Royal Astronomical Society* **456**, 119–125 (2016).

Extended Data




Figure 1: A probabilistic graphical model (PGM) represented algebraically in Equation 1. The shaded circle indicates observed data, and solid black points represent other fixed information, such as the KDEs and observational uncertainties. The remaining circles represent parameters. The underline indicates that the symbol represents a set of parameters or data. Here, κ_s and κ_{WMB} represent the KDEs of standard and WMB model populations respectively. P_s is the mixture model weighting factor. The latent parameters $\underline{\theta}$, our observations $\underline{\mathcal{D}}$ and their uncertainties $\underline{\sigma_{\mathcal{D}}}$ include temperature (T_{eff}), mass (M), log-age ($\ln(t)$), metallicity ($[\text{Fe}/\text{H}]$) and log-rotation ($\ln(P)$). This model is *hierarchical*, as all the latent parameters are drawn from the common probability distribution set by P_s and described in Equation 2.




Figure 2: Comparisons between posterior estimates of rotational parameters (data points and error bars) and the priors on these parameters (shaded regions). Shown are projected splitting ($\nu_s \sin(i)$), inclination angle (i , sampled as $\cos(i)$) and rotation period (P). Both the prior and values for P are a transformation of the upper two parameters (see **Methods** and **Supplementary Information**) In all cases, the extent of the error bars and shaded regions indicate the 68% credible (1σ) interval of the posterior and prior distributions respectively. The solid lines indicate the median of the prior distributions. In this figure, results with means and errorbars that closely resemble the prior distribution can be interpreted as prior-dominated (i.e. poorly informed by the data). All stars are sorted and coloured and sorted by age. In the case of inclination angle i and rotation period P , the displayed priors are transformed from the priors imposed on the sampled parameters from which their posteriors were derived.




Figure 3: Comparisons between posterior estimates of rotational parameters from this work, LEGACY and ‘Kages’ [20, 21, private communication]. Shown are projected splitting ($\nu_s \sin(i)$), inclination angle (i , sampled as $\cos(i)$) and rotation period (P). Fractional differences are plotted against stellar rotation obtained in this work. The Δ indicates the fractional difference between this work and the literature (i.e. stars above the zero-line have higher values in this work). The right hand panels the distribution of the fractional differences around the zero line. The colour legend is consistent throughout all panels. The x-axis units on the right hand panels are equivalent to the y-axis of the left hand panels. 10 stars have been omitted from this plot and are discussed in more detail in **Methods**: KICs 5094751, 6196457, 8349582, 8494142, 8554498, 105114430 and 11133306 all have extremely low rotation periods in Kages, with high uncertainties. Conversely, KICs 6603624, 8760414 and 8938364 have extremely high rotation periods in LEGACY with low uncertainties. In cases where stars had asymmetric error bars, the larger of the two was used when propagating uncertainty for the purposes of this figure.




Figure 4: Fractional differences between posterior estimates of asteroseismic rotation period from this work. Literature sources are: Davies et al. (2015) [12] (16 Cyg A & B), Nielsen et al. (2015) [38] (5 stars) and Benomar et al. (2018) [39] (40 stars). We used the reported parameter a_1 from the latter, which represents the rotational splitting in the case of uniform latitudinal rotation in their model. The dashed line represents the median of the sample shown, with the dotted lines representing the 15.9th and 84.1st percentiles. In cases where stars had asymmetric error bars, the larger of the two was used when propagating uncertainty for the purposes of this figure.




Figure 5: Comparisons between asteroseismic and spectroscopic measures of projected surface rotation, $v \sin(i)$. All asteroseismic (x-axis) values are from this work, all spectroscopic (y-axis) values are from the literature. *Left*: comparisons to 81 stars values reported in LEGACY and Kages. *Right*: comparisons to 16 stars observed by Benomar et al. (2015) [41]. Asteroseismic values are transformed from projected splitting ($\nu_s \sin(i)$) using the asteroseismic radius measurements presented in LEGACY and ‘Kages’. The solid lines indicate the 1:1 line, while the dash-dotted lines represent the 2:1 and 1:2 lines. The dashed lines indicate the location of $v \sin(i) = 5 \text{ kms}^{-1}$, the point at which surface and seismic measures of projected rotation begin to differ strongly [43].

Table 1: Parameters for the 94 stars for which seismic rotation rates were obtained in this work. Temperature (T_{eff}), age, mass, metallicity ($[\text{Fe}/\text{H}]$) and surface gravity ($\log(g)$) are adopted from the LEGACY [21, 22, L] and Kages [8, 20, K] catalogues, as listed in the Source column. Projected splitting ($\nu_s \sin(i)$), inclination angle (i) and asteroseismic rotation (P) are from this work. Uncertainties were taken using the 15.9th and 84.1st percentiles of posterior distributions on the parameters, which are frequently asymmetrical in linear space. Reported values are the median of the posteriors. For parameters with no direct posterior samples (e.g. rotation) the full posterior samples were transformed before taking the summary statistics. The stellar type denotes whether a star is roughly classified as belonging to the main sequence (MS), Sub-Giants (SG) or ‘hot’ stars (H) (see text). The flags indicate the following: 0; no issues, used in the gyrochronology analysis. 1; has either a number of effective samples $n_{\text{eff}} < 1000$ for the asteroseismic splitting, or Gelman-Rubin convergence metric of $\hat{R} > 1.1$ [25], indicating that rotation measurements for these stars are less robust than those with a flag of 0. 2; was found to strongly disagree with multiple literature values, excluded from the gyrochronology analysis. 3; fell outside the model range of the stellar models, and were therefore not used in the gyrochronology analysis. Table is continued on the next page.

KIC	T_{eff} [K]	Age [Gyr]	Mass [M_{\odot}]	$[\text{Fe}/\text{H}]$ [dex]	$\log(g)$ [dex]	$\nu_s \sin(i)$ [μHz]	i [$^{\circ}$]	P_{rot} [days]	Flag	Type	Source
1435467	6326 \pm 77	3.02 $^{+0.50}_{-0.35}$	1.32 $^{+0.03}_{-0.05}$	0.01 \pm 0.10	4.100 $^{+0.009}_{-0.009}$	1.58 $^{+0.10}_{-0.09}$	63.4 $^{+10.2}_{-6.6}$	6.5 $^{+0.8}_{-0.6}$	0	H	L
2837475	6614 \pm 77	1.63 $^{+0.11}_{-0.18}$	1.43 $^{+0.02}_{-0.02}$	0.01 \pm 0.10	4.163 $^{+0.007}_{-0.007}$	3.12 $^{+0.08}_{-0.08}$	70.7 $^{+6.0}_{-4.4}$	3.5 $^{+0.2}_{-0.2}$	0	H	L
3425851	6343 \pm 85	3.32 $^{+0.85}_{-0.64}$	1.18 $^{+0.05}_{-0.05}$	-0.04 \pm 0.10	4.243 $^{+0.008}_{-0.008}$	1.17 $^{+0.48}_{-0.62}$	60.9 $^{+20.1}_{-22.7}$	8.1 $^{+8.6}_{-2.7}$	0	H	K
3427720	6045 \pm 77	2.23 $^{+0.24}_{-0.24}$	1.11 $^{+0.02}_{-0.01}$	-0.06 \pm 0.10	4.387 $^{+0.005}_{-0.004}$	0.30 $^{+0.06}_{-0.06}$	56.4 $^{+22.9}_{-23.4}$	31.6 $^{+10.2}_{-11.8}$	0	MS	L
3456181	6384 \pm 77	2.09 $^{+0.13}_{-0.13}$	1.50 $^{+0.03}_{-0.02}$	-0.15 \pm 0.10	3.949 $^{+0.008}_{-0.003}$	0.92 $^{+0.08}_{-0.04}$	58.2 $^{+20.4}_{-17.7}$	10.7 $^{+2.0}_{-2.8}$	0	H	L
3544595	5669 \pm 75	6.63 $^{+0.62}_{-0.57}$	0.90 $^{+0.01}_{-0.01}$	-0.18 \pm 0.10	4.468 $^{+0.003}_{-0.003}$	0.40 $^{+0.04}_{-0.03}$	66.0 $^{+15.6}_{-13.9}$	26.1 $^{+3.9}_{-4.7}$	0	MS	K
3632418	6193 \pm 77	2.63 $^{+0.18}_{-0.18}$	1.41 $^{+0.02}_{-0.02}$	-0.12 \pm 0.10	4.024 $^{+0.008}_{-0.008}$	0.98 $^{+0.03}_{-0.03}$	72.3 $^{+10.0}_{-7.4}$	11.2 $^{+0.6}_{-0.7}$	0	MS	L
3656476	5668 \pm 77	8.37 $^{+1.72}_{-1.57}$	1.04 $^{+0.05}_{-0.04}$	0.25 \pm 0.10	4.225 $^{+0.008}_{-0.010}$	0.21 $^{+0.02}_{-0.02}$	62.4 $^{+18.9}_{-20.5}$	48.0 $^{+8.1}_{-12.7}$	0	MS	L
3735871	6107 \pm 77	2.35 $^{+1.04}_{-0.85}$	1.09 $^{+0.04}_{-0.02}$	-0.04 \pm 0.10	4.396 $^{+0.007}_{-0.004}$	0.69 $^{+0.05}_{-0.13}$	70.4 $^{+13.4}_{-15.1}$	15.8 $^{+1.8}_{-2.5}$	0	MS	L
4141376	6134 \pm 91	3.27 $^{+0.59}_{-0.64}$	1.02 $^{+0.02}_{-0.02}$	-0.24 \pm 0.10	4.412 $^{+0.003}_{-0.003}$	0.76 $^{+0.13}_{-0.13}$	64.0 $^{+17.5}_{-30.6}$	13.4 $^{+3.4}_{-3.0}$	0	MS	K
4143755	5622 \pm 106	11.27 $^{+1.50}_{-1.35}$	0.92 $^{+0.02}_{-0.03}$	-0.40 \pm 0.11	4.102 $^{+0.002}_{-0.001}$	0.18 $^{+0.08}_{-0.05}$	45.8 $^{+30.6}_{-27.4}$	48.1 $^{+27.4}_{-32.7}$	1	MS	K

Continued on next page

Table 1: Continued from previous page.

KIC	T_{eff} [K]	Age [Gyr]	Mass [M_{\odot}]	[Fe/H] [dex]	$\log(g)$ [dex]	$\nu_s \sin(i)$ [μHz]	i [$^{\circ}$]	P_{rot} [days]	Flag	Type	Source
4349452	6270 \pm 79	3.45 $^{+0.81}_{-0.72}$	1.16 $^{+0.04}_{-0.05}$	-0.04 \pm 0.10	4.275 $^{+0.008}_{-0.007}$	1.50 $^{+0.09}_{-0.09}$	79.7 $^{+7.1}_{-10.0}$	7.5 $^{+0.5}_{-0.6}$	0	H	K
4914423	5845 \pm 88	6.67 $^{+0.69}_{-0.62}$	1.10 $^{+0.02}_{-0.03}$	0.07 \pm 0.11	4.155 $^{+0.004}_{-0.004}$	0.42 $^{+0.15}_{-0.14}$	61.3 $^{+19.8}_{-30.5}$	23.1 $^{+11.7}_{-9.6}$	0	MS	K
4914923	5805 \pm 77	7.57 $^{+1.66}_{-1.79}$	1.06 $^{+0.06}_{-0.05}$	0.08 \pm 0.10	4.197 $^{+0.008}_{-0.010}$	0.39 $^{+0.03}_{-0.03}$	46.6 $^{+13.3}_{-8.1}$	21.4 $^{+5.4}_{-3.5}$	0	MS	L
5094751	5952 \pm 75	6.35 $^{+1.05}_{-1.05}$	1.07 $^{+0.04}_{-0.04}$	-0.08 \pm 0.10	4.213 $^{+0.007}_{-0.008}$	0.39 $^{+0.27}_{-0.26}$	51.5 $^{+26.7}_{-31.1}$	22.9 $^{+19.3}_{-15.8}$	0	MS	K
5184732	5846 \pm 77	4.85 $^{+1.57}_{-0.88}$	1.15 $^{+0.04}_{-0.06}$	0.36 \pm 0.10	4.255 $^{+0.010}_{-0.008}$	0.55 $^{+0.02}_{-0.02}$	71.3 $^{+11.2}_{-10.8}$	19.9 $^{+1.3}_{-1.9}$	0	MS	L
5773345	6130 \pm 84	2.55 $^{+0.26}_{-0.24}$	1.47 $^{+0.03}_{-0.03}$	0.21 \pm 0.09	3.993 $^{+0.008}_{-0.007}$	1.08 $^{+0.08}_{-0.07}$	33.7 $^{+2.5}_{-2.8}$	5.9 $^{+0.7}_{-0.5}$	0	SG	L
5866724	6169 \pm 50	3.89 $^{+0.59}_{-0.48}$	1.20 $^{+0.03}_{-0.03}$	0.09 \pm 0.08	4.224 $^{+0.007}_{-0.007}$	1.34 $^{+0.07}_{-0.06}$	80.9 $^{+9.4}_{-9.4}$	8.4 $^{+0.5}_{-0.5}$	0	MS	K
5950854	5853 \pm 77	8.93 $^{+1.12}_{-1.12}$	0.97 $^{+0.03}_{-0.03}$	-0.23 \pm 0.10	4.238 $^{+0.007}_{-0.007}$	0.29 $^{+0.64}_{-0.12}$	27.6 $^{+46.0}_{-15.2}$	22.9 $^{+34.2}_{-19.7}$	1	MS	L
6106415	6037 \pm 77	5.03 $^{+1.28}_{-1.12}$	1.07 $^{+0.03}_{-0.04}$	-0.04 \pm 0.10	4.295 $^{+0.009}_{-0.009}$	0.69 $^{+0.02}_{-0.02}$	73.1 $^{+8.4}_{-6.3}$	16.0 $^{+0.7}_{-0.8}$	0	MS	L
6116048	6033 \pm 77	9.58 $^{+2.16}_{-1.90}$	0.94 $^{+0.05}_{-0.05}$	-0.23 \pm 0.10	4.254 $^{+0.009}_{-0.012}$	0.63 $^{+0.02}_{-0.02}$	76.3 $^{+9.0}_{-10.2}$	17.9 $^{+0.8}_{-1.2}$	0	MS	L
6196457	5871 \pm 94	5.52 $^{+0.51}_{-0.48}$	1.21 $^{+0.02}_{-0.03}$	0.17 \pm 0.11	4.049 $^{+0.005}_{-0.004}$	0.43 $^{+0.28}_{-0.19}$	52.5 $^{+26.0}_{-29.1}$	20.7 $^{+19.2}_{-13.7}$	0	MS	K
6225718	6313 \pm 76	2.41 $^{+0.53}_{-0.43}$	1.16 $^{+0.03}_{-0.03}$	-0.07 \pm 0.10	4.319 $^{+0.005}_{-0.007}$	0.81 $^{+0.03}_{-0.03}$	29.1 $^{+2.1}_{-1.8}$	6.9 $^{+0.6}_{-0.5}$	0	H	L
6278762	5046 \pm 74	11.54 $^{+0.99}_{-0.94}$	0.74 $^{+0.01}_{-0.01}$	-0.37 \pm 0.09	4.560 $^{+0.002}_{-0.003}$	0.30 $^{+0.09}_{-0.09}$	62.2 $^{+19.0}_{-29.5}$	33.0 $^{+13.6}_{-13.5}$	1, 3	MS	K
6508366	6331 \pm 77	2.06 $^{+0.13}_{-0.14}$	1.53 $^{+0.02}_{-0.02}$	-0.05 \pm 0.10	3.942 $^{+0.005}_{-0.007}$	2.28 $^{+0.04}_{-0.04}$	87.0 $^{+2.1}_{-3.2}$	5.1 $^{+0.1}_{-0.1}$	0	H	L
6521045	5825 \pm 75	6.50 $^{+0.46}_{-0.56}$	1.11 $^{+0.02}_{-0.02}$	0.02 \pm 0.10	4.125 $^{+0.004}_{-0.004}$	0.45 $^{+0.02}_{-0.02}$	75.7 $^{+9.7}_{-11.2}$	24.8 $^{+1.9}_{-2.0}$	0	MS	K
6603624	5674 \pm 77	7.82 $^{+0.94}_{-0.86}$	1.01 $^{+0.03}_{-0.02}$	0.28 \pm 0.10	4.320 $^{+0.004}_{-0.005}$	1.13 $^{+0.13}_{-0.13}$	6.9 $^{+0.8}_{-0.8}$	1.2 $^{+0.0}_{-0.0}$	2	MS	L
6679371	6479 \pm 77	1.95 $^{+0.18}_{-0.16}$	1.53 $^{+0.02}_{-0.02}$	0.01 \pm 0.10	3.934 $^{+0.007}_{-0.008}$	1.90 $^{+0.05}_{-0.06}$	82.1 $^{+5.5}_{-7.2}$	6.0 $^{+0.2}_{-0.2}$	0	H	L
6933899	5832 \pm 77	6.34 $^{+0.72}_{-0.62}$	1.13 $^{+0.03}_{-0.03}$	-0.01 \pm 0.10	4.087 $^{+0.008}_{-0.007}$	0.36 $^{+0.02}_{-0.02}$	64.3 $^{+16.1}_{-14.0}$	28.9 $^{+3.7}_{-4.8}$	0	MS	L
7103006	6344 \pm 77	2.47 $^{+0.22}_{-0.24}$	1.42 $^{+0.02}_{-0.02}$	0.02 \pm 0.10	4.015 $^{+0.007}_{-0.007}$	1.36 $^{+0.09}_{-0.16}$	56.8 $^{+14.8}_{-38.4}$	7.1 $^{+1.3}_{-1.0}$	0	H	L
7106245	6068 \pm 102	6.27 $^{+1.06}_{-1.06}$	0.92 $^{+0.02}_{-0.04}$	-0.99 \pm 0.19	4.325 $^{+0.007}_{-0.007}$	0.32 $^{+0.10}_{-0.12}$	33.4 $^{+3.2}_{-14.7}$	21.4 $^{+23.8}_{-13.2}$	1, 3	MS	L
7206837	6305 \pm 77	2.90 $^{+0.42}_{-0.30}$	1.30 $^{+0.03}_{-0.03}$	0.10 \pm 0.10	4.163 $^{+0.007}_{-0.007}$	1.53 $^{+0.12}_{-0.12}$	31.7 $^{+2.8}_{-2.8}$	4.0 $^{+0.4}_{-0.4}$	0	H	L
7296438	5775 \pm 77	7.23 $^{+1.49}_{-1.77}$	1.08 $^{+0.06}_{-0.05}$	0.19 \pm 0.10	4.201 $^{+0.009}_{-0.010}$	0.20 $^{+0.06}_{-0.06}$	50.5 $^{+28.2}_{-31.1}$	45.6 $^{+23.4}_{-29.0}$	1	MS	L
7510397	6171 \pm 77	2.82 $^{+0.17}_{-0.16}$	1.37 $^{+0.02}_{-0.02}$	-0.21 \pm 0.10	4.036 $^{+0.004}_{-0.004}$	0.64 $^{+0.06}_{-0.06}$	19.9 $^{+2.0}_{-2.0}$	6.1 $^{+0.7}_{-0.6}$	0	MS	L
7670943	6463 \pm 110	2.78 $^{+0.62}_{-0.45}$	1.24 $^{+0.04}_{-0.04}$	0.09 \pm 0.11	4.228 $^{+0.008}_{-0.008}$	1.83 $^{+0.14}_{-0.14}$	75.7 $^{+5.8}_{-11.5}$	6.0 $^{+0.6}_{-0.6}$	0	H	K
7680114	5811 \pm 77	7.68 $^{+1.45}_{-1.28}$	1.06 $^{+0.04}_{-0.05}$	0.05 \pm 0.10	4.172 $^{+0.008}_{-0.010}$	0.26 $^{+0.05}_{-0.04}$	37.0 $^{+44.3}_{-16.3}$	27.3 $^{+19.6}_{-13.5}$	0	MS	L
7771282	6248 \pm 77	3.24 $^{+0.35}_{-0.32}$	1.29 $^{+0.03}_{-0.03}$	-0.02 \pm 0.10	4.112 $^{+0.007}_{-0.007}$	1.01 $^{+0.14}_{-0.14}$	69.5 $^{+17.8}_{-17.8}$	10.4 $^{+2.3}_{-1.7}$	0	MS	L
7871531	5501 \pm 77	9.96 $^{+1.93}_{-1.77}$	0.83 $^{+0.03}_{-0.02}$	-0.26 \pm 0.10	4.478 $^{+0.007}_{-0.005}$	0.33 $^{+0.03}_{-0.03}$	71.3 $^{+12.0}_{-13.2}$	33.1 $^{+4.1}_{-4.1}$	0	MS	L
7940546	6235 \pm 77	2.33 $^{+0.08}_{-0.08}$	1.40 $^{+0.03}_{-0.01}$	-0.20 \pm 0.10	4.007 $^{+0.003}_{-0.003}$	1.14 $^{+0.03}_{-0.03}$	78.9 $^{+7.4}_{-7.9}$	9.9 $^{+0.3}_{-0.4}$	0	MS	L
7970740	5309 \pm 77	12.98 $^{+1.36}_{-2.00}$	0.73 $^{+0.01}_{-0.01}$	-0.54 \pm 0.10	4.539 $^{+0.004}_{-0.005}$	0.26 $^{+0.03}_{-0.02}$	60.7 $^{+17.4}_{-15.3}$	39.2 $^{+6.7}_{-9.0}$	0	MS	L
8006161	5488 \pm 77	3.59 $^{+1.53}_{-1.45}$	0.98 $^{+0.03}_{-0.03}$	0.34 \pm 0.10	4.494 $^{+0.007}_{-0.007}$	0.34 $^{+0.02}_{-0.02}$	37.0 $^{+4.1}_{-3.4}$	20.6 $^{+2.2}_{-1.8}$	0	MS	L
8077137	6072 \pm 75	6.23 $^{+0.56}_{-1.23}$	1.12 $^{+0.04}_{-0.05}$	-0.09 \pm 0.10	4.056 $^{+0.010}_{-0.013}$	0.84 $^{+0.06}_{-0.07}$	72.8 $^{+11.3}_{-12.6}$	13.0 $^{+1.3}_{-1.5}$	0	MS	K
8150065	6173 \pm 101	3.83 $^{+0.99}_{-0.67}$	1.19 $^{+0.04}_{-0.05}$	-0.13 \pm 0.15	4.220 $^{+0.008}_{-0.008}$	0.54 $^{+0.11}_{-0.13}$	64.0 $^{+18.0}_{-21.1}$	18.6 $^{+6.4}_{-4.9}$	0	MS	L
8179536	6343 \pm 77	3.54 $^{+1.04}_{-0.81}$	1.16 $^{+0.05}_{-0.06}$	-0.03 \pm 0.10	4.255 $^{+0.010}_{-0.010}$	1.46 $^{+0.10}_{-0.09}$	55.7 $^{+13.3}_{-7.4}$	6.5 $^{+1.2}_{-0.8}$	0	H	L

Continued on next page

Table 1: Continued from previous page.

KIC	T_{eff} [K]	Age [Gyr]	Mass [M_{\odot}]	[Fe/H] [dex]	$\log(g)$ [dex]	$\nu_s \sin(i)$ [μHz]	i [$^{\circ}$]	P_{rot} [days]	Flag	Type	Source
8228742	6122 \pm 77	2.89 $^{+0.16}_{-0.18}$	1.38 $^{+0.02}_{-0.02}$	-0.08 \pm 0.10	4.035 $^{+0.007}_{-0.005}$	0.64 $^{+0.04}_{-0.04}$	37.9 $^{+6.2}_{-4.0}$	11.0 $^{+2.0}_{-1.3}$	0	MS	L
8292840	6239 \pm 94	3.85 $^{+0.81}_{-0.75}$	1.15 $^{+0.05}_{-0.05}$	-0.14 \pm 0.10	4.240 $^{+0.008}_{-0.008}$	1.45 $^{+0.07}_{-0.07}$	76.2 $^{+8.9}_{-9.1}$	7.7 $^{+0.5}_{-0.5}$	0	MS	K
8349582	5699 \pm 74	8.03 $^{+0.80}_{-0.70}$	1.07 $^{+0.02}_{-0.02}$	0.30 \pm 0.10	4.163 $^{+0.004}_{-0.003}$	0.23 $^{+0.07}_{-0.06}$	59.6 $^{+20.7}_{-24.1}$	41.7 $^{+19.6}_{-14.9}$	1	MS	K
8379927	6067 \pm 120	1.99 $^{+0.85}_{-0.75}$	1.12 $^{+0.04}_{-0.04}$	-0.10 \pm 0.15	4.388 $^{+0.008}_{-0.007}$	1.12 $^{+0.02}_{-0.02}$	63.3 $^{+2.5}_{-2.3}$	9.2 $^{+0.3}_{-0.2}$	0	MS	L
8394589	6143 \pm 77	4.45 $^{+0.83}_{-0.83}$	1.04 $^{+0.04}_{-0.04}$	-0.29 \pm 0.10	4.322 $^{+0.008}_{-0.007}$	1.01 $^{+0.03}_{-0.03}$	71.1 $^{+7.9}_{-5.9}$	10.9 $^{+0.6}_{-0.6}$	0	MS	L
8424992	5719 \pm 77	9.61 $^{+1.92}_{-1.74}$	0.92 $^{+0.04}_{-0.04}$	-0.12 \pm 0.10	4.359 $^{+0.007}_{-0.007}$	0.22 $^{+0.06}_{-0.06}$	59.1 $^{+21.3}_{-20.3}$	42.3 $^{+19.4}_{-17.7}$	1	MS	L
8494142	6144 \pm 106	2.62 $^{+0.26}_{-0.24}$	1.42 $^{+0.03}_{-0.03}$	0.13 \pm 0.10	4.038 $^{+0.005}_{-0.005}$	0.67 $^{+0.22}_{-0.21}$	62.8 $^{+18.5}_{-18.8}$	14.5 $^{+9.9}_{-4.5}$	0	MS	K
8554498	5945 \pm 60	5.60 $^{+0.45}_{-0.42}$	1.20 $^{+0.02}_{-0.02}$	0.17 \pm 0.05	4.007 $^{+0.003}_{-0.003}$	0.25 $^{+0.09}_{-0.09}$	48.1 $^{+30.0}_{-36.6}$	35.7 $^{+25.6}_{-31.4}$	0	MS	K
8694723	6246 \pm 77	4.69 $^{+0.48}_{-0.51}$	1.14 $^{+0.02}_{-0.02}$	-0.42 \pm 0.10	4.113 $^{+0.007}_{-0.009}$	0.92 $^{+0.06}_{-0.05}$	34.7 $^{+3.4}_{-2.7}$	7.2 $^{+0.8}_{-0.9}$	0	MS	L
8760414	5873 \pm 77	11.66 $^{+1.28}_{-1.61}$	0.81 $^{+0.03}_{-0.02}$	-0.92 \pm 0.10	4.329 $^{+0.006}_{-0.005}$	0.69 $^{+0.26}_{-0.42}$	7.6 $^{+2.3}_{-1.7}$	2.0 $^{+4.5}_{-0.4}$	2, 3	MS	L
8866102	6325 \pm 75	2.60 $^{+0.56}_{-0.53}$	1.23 $^{+0.04}_{-0.04}$	0.01 \pm 0.10	4.262 $^{+0.008}_{-0.007}$	2.15 $^{+0.04}_{-0.04}$	78.2 $^{+6.6}_{-4.7}$	5.3 $^{+0.2}_{-0.2}$	0	H	K
8938364	5677 \pm 77	10.25 $^{+0.56}_{-0.65}$	0.99 $^{+0.01}_{-0.01}$	-0.13 \pm 0.10	4.173 $^{+0.007}_{-0.002}$	0.61 $^{+0.27}_{-0.50}$	8.7 $^{+57.5}_{-2.3}$	2.0 $^{+87.0}_{-0.2}$	2	MS	L
9025370	5270 \pm 180	6.55 $^{+1.26}_{-1.13}$	0.97 $^{+0.03}_{-0.03}$	-0.12 \pm 0.18	4.423 $^{+0.007}_{-0.004}$	0.43 $^{+0.04}_{-0.04}$	67.5 $^{+13.2}_{-19.1}$	24.7 $^{+3.7}_{-4.7}$	0	MS	L
9098294	5852 \pm 77	8.08 $^{+0.99}_{-0.73}$	0.97 $^{+0.02}_{-0.03}$	-0.18 \pm 0.10	4.308 $^{+0.005}_{-0.007}$	0.36 $^{+0.04}_{-0.04}$	58.2 $^{+21.0}_{-16.3}$	27.2 $^{+5.7}_{-7.0}$	0	MS	L
9139151	6302 \pm 77	1.32 $^{+0.94}_{-0.75}$	1.18 $^{+0.04}_{-0.05}$	0.10 \pm 0.10	4.382 $^{+0.008}_{-0.008}$	0.95 $^{+0.04}_{-0.04}$	73.5 $^{+11.0}_{-11.0}$	11.6 $^{+0.8}_{-1.1}$	0	H	L
9139163	6400 \pm 84	1.60 $^{+0.22}_{-0.21}$	1.40 $^{+0.02}_{-0.02}$	0.15 \pm 0.09	4.200 $^{+0.009}_{-0.008}$	1.59 $^{+0.07}_{-0.08}$	33.5 $^{+3.0}_{-3.0}$	4.0 $^{+0.3}_{-0.3}$	1	H	L
9206432	6538 \pm 77	1.53 $^{+0.21}_{-0.30}$	1.38 $^{+0.02}_{-0.02}$	0.16 \pm 0.10	4.220 $^{+0.005}_{-0.007}$	1.55 $^{+0.17}_{-0.20}$	34.3 $^{+5.7}_{-4.2}$	4.1 $^{+1.0}_{-0.5}$	0	H	L
9353712	6278 \pm 77	2.15 $^{+0.11}_{-0.13}$	1.51 $^{+0.03}_{-0.02}$	-0.05 \pm 0.10	3.943 $^{+0.007}_{-0.005}$	0.75 $^{+0.16}_{-0.17}$	37.6 $^{+29.1}_{-12.6}$	9.5 $^{+7.3}_{-3.9}$	0	H	L
9410862	6047 \pm 77	6.93 $^{+1.49}_{-1.33}$	0.97 $^{+0.05}_{-0.04}$	-0.31 \pm 0.10	4.300 $^{+0.009}_{-0.008}$	0.41 $^{+0.08}_{-0.08}$	46.4 $^{+28.3}_{-16.7}$	20.6 $^{+9.8}_{-8.1}$	1	MS	L
9414417	6253 \pm 75	2.65 $^{+0.16}_{-0.18}$	1.40 $^{+0.02}_{-0.03}$	-0.13 \pm 0.10	4.016 $^{+0.005}_{-0.005}$	1.09 $^{+0.05}_{-0.05}$	58.1 $^{+9.7}_{-6.9}$	9.0 $^{+1.1}_{-0.9}$	0	H	L
9592705	6174 \pm 92	2.33 $^{+0.46}_{-0.35}$	1.51 $^{+0.03}_{-0.04}$	0.22 \pm 0.10	3.961 $^{+0.003}_{-0.004}$	0.93 $^{+0.09}_{-0.09}$	59.3 $^{+17.8}_{-12.1}$	10.7 $^{+2.1}_{-1.9}$	0	SG	K
9812850	6321 \pm 77	2.71 $^{+0.35}_{-0.46}$	1.37 $^{+0.04}_{-0.04}$	-0.07 \pm 0.10	4.053 $^{+0.008}_{-0.009}$	1.54 $^{+0.08}_{-0.08}$	81.0 $^{+6.4}_{-10.0}$	7.4 $^{+0.4}_{-0.5}$	0	H	L
9955598	5457 \pm 77	6.29 $^{+1.84}_{-0.86}$	0.90 $^{+0.04}_{-0.04}$	0.05 \pm 0.10	4.497 $^{+0.007}_{-0.008}$	0.29 $^{+0.04}_{-0.04}$	53.4 $^{+20.8}_{-11.9}$	31.4 $^{+9.1}_{-6.4}$	0	MS	L
9965715	5860 \pm 180	2.92 $^{+0.75}_{-0.68}$	1.21 $^{+0.04}_{-0.04}$	-0.44 \pm 0.18	4.272 $^{+0.003}_{-0.004}$	1.75 $^{+0.05}_{-0.05}$	58.3 $^{+3.5}_{-3.2}$	5.6 $^{+0.3}_{-0.3}$	0	MS	L
10068307	6132 \pm 77	2.36 $^{+0.08}_{-0.10}$	1.47 $^{+0.01}_{-0.01}$	-0.23 \pm 0.10	3.967 $^{+0.004}_{-0.004}$	0.71 $^{+0.03}_{-0.03}$	41.7 $^{+4.1}_{-4.1}$	10.9 $^{+1.5}_{-1.5}$	0	SG	L
10079226	5949 \pm 77	3.06 $^{+0.70}_{-0.65}$	1.12 $^{+0.02}_{-0.03}$	0.11 \pm 0.10	4.366 $^{+0.005}_{-0.005}$	0.65 $^{+0.07}_{-0.08}$	75.1 $^{+10.6}_{-23.4}$	16.8 $^{+2.2}_{-3.0}$	1	MS	L
10162436	6146 \pm 77	2.46 $^{+0.10}_{-0.11}$	1.45 $^{+0.03}_{-0.01}$	-0.16 \pm 0.10	3.981 $^{+0.005}_{-0.005}$	0.84 $^{+0.05}_{-0.05}$	25.5 $^{+2.1}_{-1.8}$	5.9 $^{+0.6}_{-0.4}$	0	SG	L
10454113	6177 \pm 77	2.89 $^{+0.56}_{-0.53}$	1.17 $^{+0.03}_{-0.03}$	-0.07 \pm 0.10	4.314 $^{+0.005}_{-0.005}$	0.76 $^{+0.07}_{-0.07}$	40.9 $^{+4.8}_{-9.6}$	10.0 $^{+4.8}_{-2.5}$	0	MS	L
10514430	5784 \pm 98	7.84 $^{+0.40}_{-0.91}$	1.06 $^{+0.04}_{-0.02}$	-0.11 \pm 0.11	4.061 $^{+0.004}_{-0.004}$	0.18 $^{+0.05}_{-0.05}$	57.2 $^{+23.5}_{-27.2}$	53.6 $^{+23.6}_{-27.2}$	1	MS	K
10516096	5964 \pm 77	7.01 $^{+1.33}_{-1.45}$	1.06 $^{+0.05}_{-0.06}$	-0.11 \pm 0.10	4.169 $^{+0.010}_{-0.011}$	0.48 $^{+0.03}_{-0.03}$	71.8 $^{+12.5}_{-16.2}$	22.6 $^{+1.9}_{-3.1}$	0	MS	L
10586004	5770 \pm 83	6.43 $^{+0.64}_{-0.61}$	1.18 $^{+0.02}_{-0.03}$	0.29 \pm 0.10	4.071 $^{+0.005}_{-0.005}$	0.48 $^{+0.16}_{-0.17}$	59.3 $^{+20.6}_{-22.4}$	19.6 $^{+11.1}_{-6.5}$	1	MS	K
10644253	6045 \pm 77	2.39 $^{+1.12}_{-0.96}$	1.10 $^{+0.04}_{-0.04}$	0.06 \pm 0.10	4.396 $^{+0.007}_{-0.008}$	0.24 $^{+0.08}_{-0.08}$	55.6 $^{+24.0}_{-30.3}$	38.0 $^{+21.1}_{-19.1}$	0	MS	L
10666592	6350 \pm 80	2.11 $^{+0.29}_{-0.24}$	1.50 $^{+0.04}_{-0.04}$	0.26 \pm 0.08	4.017 $^{+0.009}_{-0.007}$	0.91 $^{+0.11}_{-0.11}$	47.2 $^{+27.2}_{-14.8}$	9.5 $^{+3.6}_{-3.2}$	0	H	K

Continued on next page

Table 1: *Continued from previous page.*

KIC	T_{eff} [K]	Age [Gyr]	Mass [M_{\odot}]	[Fe/H] [dex]	$\log(g)$ [dex]	$\nu_s \sin(i)$ [μHz]	i [$^{\circ}$]	P_{rot} [days]	Flag	Type	Source
10730618	6150 \pm 180	3.05 $^{+0.46}_{-0.29}$	1.34 $^{+0.04}_{-0.05}$	-0.11 \pm 0.18	4.062 $^{+0.008}_{-0.007}$	0.56 $^{+0.23}_{-0.25}$	55.6 $^{+23.7}_{-26.2}$	16.1 $^{+13.9}_{-7.2}$	0	MS	L
10963065	6140 \pm 77	7.15 $^{+1.92}_{-1.61}$	0.99 $^{+0.06}_{-0.06}$	-0.19 \pm 0.10	4.277 $^{+0.011}_{-0.011}$	0.67 $^{+0.03}_{-0.03}$	41.8 $^{+4.7}_{-3.6}$	11.5 $^{+1.3}_{-1.0}$	0	MS	L
11081729	6548 \pm 82	1.88 $^{+0.59}_{-0.42}$	1.30 $^{+0.04}_{-0.05}$	0.11 \pm 0.10	4.245 $^{+0.010}_{-0.009}$	3.35 $^{+0.10}_{-0.10}$	82.9 $^{+4.9}_{-5.9}$	3.4 $^{+0.1}_{-0.1}$	0	H	L
11133306	5982 \pm 82	5.14 $^{+0.86}_{-0.88}$	1.06 $^{+0.04}_{-0.03}$	-0.02 \pm 0.10	4.314 $^{+0.007}_{-0.005}$	0.39 $^{+0.14}_{-0.13}$	58.0 $^{+21.9}_{-24.1}$	24.6 $^{+14.3}_{-10.0}$	0	MS	K
11253226	6642 \pm 77	1.60 $^{+0.06}_{-0.13}$	1.41 $^{+0.02}_{-0.01}$	-0.08 \pm 0.10	4.173 $^{+0.004}_{-0.003}$	2.55 $^{+0.11}_{-0.12}$	49.3 $^{+6.3}_{-4.4}$	3.4 $^{+0.4}_{-0.3}$	0	H	L
11295426	5793 \pm 74	6.31 $^{+0.32}_{-0.34}$	1.07 $^{+0.01}_{-0.02}$	0.12 \pm 0.07	4.280 $^{+0.003}_{-0.003}$	0.22 $^{+0.03}_{-0.03}$	55.6 $^{+22.9}_{-20.9}$	42.6 $^{+11.5}_{-14.5}$	0	MS	K
11401755	5911 \pm 66	7.10 $^{+0.61}_{-0.59}$	1.06 $^{+0.03}_{-0.02}$	-0.20 \pm 0.06	4.039 $^{+0.004}_{-0.003}$	0.55 $^{+0.09}_{-0.10}$	64.6 $^{+17.4}_{-19.5}$	18.5 $^{+4.7}_{-5.3}$	0	MS	K
11772920	5180 \pm 180	10.67 $^{+2.73}_{-2.97}$	0.83 $^{+0.04}_{-0.04}$	-0.09 \pm 0.18	4.500 $^{+0.005}_{-0.008}$	0.31 $^{+0.03}_{-0.04}$	70.8 $^{+12.6}_{-14.3}$	35.1 $^{+5.3}_{-4.7}$	1	MS	L
11807274	6225 \pm 75	3.59 $^{+0.78}_{-0.45}$	1.24 $^{+0.04}_{-0.04}$	0.00 \pm 0.08	4.135 $^{+0.009}_{-0.007}$	1.42 $^{+0.07}_{-0.07}$	76.8 $^{+8.8}_{-9.4}$	7.9 $^{+0.5}_{-0.5}$	0	MS	K
11853905	5781 \pm 76	6.71 $^{+0.77}_{-0.67}$	1.12 $^{+0.02}_{-0.03}$	0.09 \pm 0.10	4.102 $^{+0.004}_{-0.005}$	0.27 $^{+0.09}_{-0.09}$	54.7 $^{+24.5}_{-28.8}$	34.0 $^{+18.9}_{-17.2}$	1	MS	K
11904151	5647 \pm 74	10.23 $^{+0.83}_{-0.67}$	0.92 $^{+0.01}_{-0.02}$	-0.15 \pm 0.10	4.344 $^{+0.003}_{-0.003}$	0.24 $^{+0.06}_{-0.07}$	64.9 $^{+18.1}_{-28.4}$	40.9 $^{+16.5}_{-12.8}$	1	MS	K
12009504	6179 \pm 77	3.97 $^{+0.57}_{-0.43}$	1.17 $^{+0.02}_{-0.04}$	-0.08 \pm 0.10	4.211 $^{+0.007}_{-0.005}$	1.16 $^{+0.03}_{-0.04}$	71.3 $^{+8.0}_{-5.7}$	9.4 $^{+0.5}_{-0.5}$	0	MS	L
12069127	6276 \pm 77	2.01 $^{+0.11}_{-0.13}$	1.57 $^{+0.03}_{-0.02}$	0.08 \pm 0.10	3.912 $^{+0.005}_{-0.004}$	0.54 $^{+1.26}_{-0.33}$	16.6 $^{+54.2}_{-6.3}$	5.2 $^{+39.9}_{-3.9}$	1	H	L
12069424	5825 \pm 50	6.67 $^{+0.81}_{-0.77}$	1.05 $^{+0.02}_{-0.02}$	0.10 \pm 0.03	4.287 $^{+0.007}_{-0.007}$	0.40 $^{+0.01}_{-0.01}$	44.7 $^{+6.2}_{-2.9}$	20.5 $^{+2.0}_{-1.1}$	1	MS	L
12069449	5750 \pm 50	7.39 $^{+0.89}_{-0.91}$	0.99 $^{+0.02}_{-0.02}$	0.05 \pm 0.02	4.353 $^{+0.007}_{-0.005}$	0.31 $^{+0.01}_{-0.01}$	34.0 $^{+3.0}_{-2.4}$	21.2 $^{+1.8}_{-1.5}$	0	MS	L
12258514	5964 \pm 77	4.05 $^{+0.18}_{-0.16}$	1.26 $^{+0.01}_{-0.01}$	0.00 \pm 0.10	4.126 $^{+0.004}_{-0.003}$	0.39 $^{+0.04}_{-0.03}$	35.0 $^{+22.9}_{-6.6}$	16.7 $^{+10.0}_{-3.6}$	1	MS	L
12317678	6580 \pm 77	2.46 $^{+0.22}_{-0.18}$	1.34 $^{+0.04}_{-0.01}$	-0.28 \pm 0.10	4.048 $^{+0.008}_{-0.009}$	1.27 $^{+0.13}_{-0.14}$	35.3 $^{+10.1}_{-5.2}$	5.2 $^{+1.8}_{-0.9}$	0	H	L