

Student Online Teaching Advice Notice

The materials and content presented within this session are intended solely for use in a context of teaching and learning at Trinity.

Any session recorded for subsequent review is made available solely for the purpose of enhancing student learning.

Students should not edit or modify the recording in any way, nor disseminate it for use outside of a context of teaching and learning at Trinity.

Please be mindful of your physical environment and conscious of what may be captured by the device camera and microphone during videoconferencing calls.

Recorded materials will be handled in compliance with Trinity's statutory duties under the Universities Act, 1997 and in accordance with the University's [policies and procedures](#).

Further information on data protection and best practice when using videoconferencing software is available at https://www.tcd.ie/info_compliance/data-protection/.

© Trinity College Dublin 2020



Trinity College Dublin
Coláiste na Tríonóide, Baile Átha Cliath
The University of Dublin

4 Relations

Task: Define subsets of Cartesian products with certain properties. Understand the predicates " $=$ " (equality) and other predicates in predicate logic in a more abstract light.

Start with $x = y$. The element x is some relation R to y (equality in this case).

We can also denote it as xRy or $(x, y) \in E$

Let $x, y \in \mathbb{R}$, then $E = \{(x, x) \mid x \in \mathbb{R}\} \subset \mathbb{R} \times \mathbb{R}$.

The "diagonal" in $\mathbb{R} \times \mathbb{R}$ gives exactly the elements equal to each other.

More generally:

Definition: Let A, B be sets. A subset of the Cartesian product $A \times B$ is called a relation between A and B . A subset of the Cartesian product $A \times A$ is called a relation on A .

Remark: Note how general this definition is. To make it useful for understanding predicates, we will need to introduce key properties relations can satisfy.

Example: $A = \{1, 3, 7\}$ $B = \{1, 2, 5\}$

We can define a relation S on $A \times B$ by $S = \{(1, 1), (1, 5), (3, 2)\}$. This means $1S1$, $1S5$ and $3S2$ and no other ordered pairs in $A \times B$ satisfy S .

Remark: The relations we defined involve 2 elements, so they are often called binary relations in the literature.

4.1 Equivalence Relations

Task: Define the most useful kind of relation.

Definition: A relation R on a set A is called

1. reflexive iff (if and only if) $\forall x \in A, xRx$


2. symmetric iff $\forall x, y \in A, xRy \rightarrow yRx$
3. transitive iff $\forall x, y, z \in A, xRy \wedge yRz \rightarrow xRz$

An equivalence relation on A is a relation that is reflexive, symmetric, and transitive.

Notation: Instead of xRy , an equivalence relation is often denoted by $x \equiv y$ or $x \sim y$.

Examples:

1. "=" equality is an equivalence relation.
 - (a) $x = x$ reflexive
 - (b) $x = y \Rightarrow y = x$ symmetric
 - (c) $x = y \wedge y = z \Rightarrow x = z$ transitive
2. $A = \mathbb{N}$
 $x \equiv y \pmod{3}$ is an equivalence relation. $x \equiv y \pmod{3}$ means $x - y = 3m$ for some $m \in \mathbb{Z}$, **i.e.** x and y have the same remainder when divided by 3. The set of all possible remainders is $\{0, 1, 2\}$
NB: In correct logic notation, $x \equiv y \pmod{3}$ if $\exists m \in \mathbb{Z}$ s.t. $x - y = 3m$
 - (a) $x \equiv x \pmod{3}$ since $x - x = 0 = 3 \times 0 \rightarrow$ reflexive
 - (b) $x \equiv y \pmod{3} \Rightarrow y \equiv x \pmod{3}$ because $x \equiv y \pmod{3}$ means $x - y = 3m$ for some $m \in \mathbb{Z} \Rightarrow y - x = -3m = 3 \times (-m) \Rightarrow y \equiv x \pmod{3} \rightarrow$ symmetric
 - (c) Assume $x \equiv y \pmod{3}$ and $y \equiv z \pmod{3}$
 $x \equiv y \pmod{3} \Rightarrow \exists m \in \mathbb{Z}$ s.t. $x - y = 3m \Rightarrow y = x - 3m$
 $y \equiv z \pmod{3} \Rightarrow \exists p \in \mathbb{Z}$ s.t. $y - z = 3p \Rightarrow y = z + 3p$
Therefore, $x - 3m = z + 3p \Leftrightarrow x - z = 3p + 3m = 3(p + m)$
Since $p, m \in \mathbb{Z}, p + m \in \mathbb{Z} \Rightarrow x \equiv z \pmod{3} \rightarrow$ transitive.
3. Let $f : A \rightarrow A$ be any function on a non-empty set A . We define the relation $R = \{(x, y) \mid f(x) = f(y)\}$
 - (a) $\forall x \in A, f(x) = f(x) \Rightarrow (x, x) \in R \rightarrow$ reflexive
 - (b) If $(x, y) \in R$, then $f(x) = f(y) \Rightarrow f(y) = f(x)$, **i.e.** $(y, x) \in R \rightarrow$ symmetric
 - (c) If $(x, y) \in R$ and $(y, z) \in R$, then $f(x) = f(y)$ and $f(y) = f(z)$, which by the transitivity of equality implies $f(x) = f(z)$, **i.e.** $(x, z) \in R$ as needed, so R is transitive as well.
 $f(x)$ can be $e^x, \sin x, |x|$, etc.



4. Let Γ be the set of all triangles in the plane. $ABC \sim A'B'C'$ if ABC and $A'B'C'$ are similar triangles, **i.e.** have equal angles.

(a) $\forall ABC \in \Gamma, ABC \sim ABC$ so \sim is reflexive

(b) $ABC \sim A'B'C' \Rightarrow A'B'C' \sim ABC$ so \sim is symmetric

(c) $ABC \sim A'B'C'$ and $A'B'C' \sim A''B''C'' \Rightarrow ABC \sim A''B''C''$,
so \sim is transitive

Clearly (a), (b), (c) use the fact that equality of angles is an equivalence relation.

Exercise: For various predicates you've encountered, check whether reflexive, symmetric or transitive. Examples of predicates include \neq , $<$, $>$, \leq , \geq , \subseteq , \rightarrow , \leftrightarrow