

Student Online Teaching Advice Notice

The materials and content presented within this session are intended solely for use in a context of teaching and learning at Trinity.

Any session recorded for subsequent review is made available solely for the purpose of enhancing student learning.

Students should not edit or modify the recording in any way, nor disseminate it for use outside of a context of teaching and learning at Trinity.

Please be mindful of your physical environment and conscious of what may be captured by the device camera and microphone during videoconferencing calls.

Recorded materials will be handled in compliance with Trinity's statutory duties under the Universities Act, 1997 and in accordance with the University's [policies and procedures](#).

Further information on data protection and best practice when using videoconferencing software is available at https://www.tcd.ie/info_compliance/data-protection/.

© Trinity College Dublin 2020



7 Abstract Algebra

Task: Understand binary operations, semigroups, monoids, and groups as well as their properties.

7.1 Binary Operations

Definition: Let A be a set. A binary operation $*$ on A is an operation applied to any two elements $x, y \in A$ that yields an element $x * y$ in A . In other words, $*$ is a binary operation on A if $\forall x, y \in A, x * y \in A$.

Examples:

1. $\mathbb{R}, +$ addition on $\mathbb{R} : \forall x, y \in \mathbb{R}, x + y \in \mathbb{R}$
2. $\mathbb{R}, -$ subtraction on $\mathbb{R} : \forall x, y \in \mathbb{R}, x - y \in \mathbb{R}$
3. \mathbb{R}, \times multiplication on $\mathbb{R} : \forall x, y \in \mathbb{R}, x \times y \in \mathbb{R}$
4. $\mathbb{R}, /$, division on \mathbb{R} is NOT a binary operation because $\forall x \in \mathbb{R} \exists 0 \in \mathbb{R}$ s.t. $\frac{x}{0}$ is undefined (not an element of \mathbb{R})
5. Let A be the set of all lists or strings. Concatenation is a binary operation.

Definition: A binary operation $*$ on a set A is called commutative if $\forall x, y \in A, x * y = y * x$

Examples:

1. $\mathbb{R}, +$ is commutative since $\forall x, y \in \mathbb{R}, x + y = y + x$
2. \mathbb{R}, \times is commutative since $\forall x, y \in \mathbb{R}, x \times y = y \times x$
3. $\mathbb{R}, -$ is not commutative since $\forall x, y \in \mathbb{R}, x - y \neq y - x$ in general. $x - y = y - x$ only if $x = y$
4. Let M_n be the set of n by n matrices with entries in \mathbb{R} , and let $*$ be matrix multiplication. $\forall A, B \in M_n, A * B \in M_n$, so $*$ is a binary operation, but $A * B \neq B * A$ in general. Therefore $*$ is not commutative.

Definition: A binary operation $*$ on a set A is called associative if $\forall x, y, z \in A$
 $(x * y) * z = x * (y * z)$

Examples:

1. $\mathbb{R}, +$ is associative since $\forall x, y, z \in \mathbb{R}, (x + y) + z = x + (y + z)$
2. \mathbb{R}, \times is associative since $\forall x, y, z \in \mathbb{R}, (x \times y) \times z = x \times (y \times z)$
3. Intersection \cap on sets is associative since $\forall A, B, C$ sets $(A \cap B) \cap C = A \cap (B \cap C)$.
4. Union \cup on sets is associative since $\forall A, B, C$ sets $(A \cup B) \cup C = A \cup (B \cup C)$
5. $\mathbb{R}, -$ is not associative since $(1 - 3) - 5 = -2 - 5 = -7$ but $1 - (3 - 5) = 1 - (-2) = 1 + 2 = 3$

Remark: When we are dealing with associative binary operations we can drop the parentheses, **i.e.** $(x * y) * z$ can be written $x * y * z$.

7.2 Semigroups

Definition: A semigroup is a set endowed with an associative binary operation. We denote the semigroup $(A, *)$

Examples:

1. $(\mathbb{R}, +)$ and (\mathbb{R}, \times) are semigroups.
2. Let A be a set and let $P(A)$ be its power set. $(P(A), \cap)$ and $(P(A), \cup)$ are both semigroups.
3. $(M_n, *)$, the set of $n \times n$ matrices with entries in \mathbb{R} with the operation of matrix multiplication (which is associative \rightarrow a bit gory to prove) forms a semigroup.

Since $*$ is associative on a semigroup, we can define a^n :

$$\begin{aligned} a^1 &= a \\ a^2 &= a * a \\ a^3 &= a * a * a \\ &\vdots \end{aligned}$$

Recursively, $a^1 = a$ and $a^n = a * a^{n-1}, \forall n > 1$

NB: In $(\mathbb{R}, \times), \forall a \in \mathbb{R}, a^n = \underbrace{a \times a \times \dots \times a}_{n \text{ times}}$, whereas in $(\mathbb{R}, +), \forall a \in \mathbb{R}, a^n = \underbrace{a + a + \dots + a}_{n \text{ times}} = na$. Be careful what $*$ stands for!

Theorem: Let $(A, *)$ be a semigroup. $\forall a \in A, a^m * a^n = a^{m+n}, \forall m, n \in \mathbb{N}^*$.

Proof: By induction on m .

Base Case: $m = 1 \quad a^1 * a^n = a * a^n = a^{1+n}$

Inductive Step: Assume the result is true for $m = p$, **i.e.** $a^p * a^n = a^{p+n}$
and seek to prove that $a^{p+1} * a^n = a^{p+1+n}$

$$a^{p+1} * a^n = (a * a^p) * a^n = a * (a^p * a^n) = a * a^{p+n} = a^{p+1+n}$$

Theorem: Let $(A, *)$ be a semigroup. $\forall a \in A, (a^m)^n = a^{mn}, \forall m, n \in \mathbb{N}^*$

Proof: By induction on n .

Base Case: $n = 1 \quad (a^m)^1 = a^m = a^{m \times 1}$

Inductive Step: Assume the result is true for $n = p$, **i.e.** $(a^m)^p = a^{mp}$
and seek to prove that $(a^m)^{p+1} = a^{m(p+1)}$

$$(a^m)^{p+1} = (a^m)^p * a^m = a^{mp} * a^m = a^{mp+m} = a^{m(p+1)} \text{ by the previous theorem.}$$

7.2.1 General Associative Law

Let $(A, *)$ be a semigroup. $\forall a_1, \dots, a_s \in A, a_1 * a_2 * \dots * a_s$ has the same value regardless of how the product is bracketed.

Proof Use associativity of $*$.

qed

NB: Unless $(A, *)$ has a commutative binary operation, $a_1 * a_2 * \dots * a_s$ does depend on the ORDER in which the a'_j s appear in $a_1 * a_2 * \dots * a_s$