




a
Clear
Cloud clearing
Complex survey: estimated $T_{\rm eff} > 3000$ K
$0.1071\,M_\odot$
BD interval
Reference stars
Effective temperature at 125 Myr (K)
Mass ($M_\odot$)
3500
3000
2500
2000
1500
0.03
0.05
0.08
0.10
0.15
0.20
0.25


b
Greater Pleiades Complex
UPK 303
GPC-1
AB Dor
Pleiades
GPC-2
HSC 1964
UPK 545
Orange: additional reference stars
from complex catalogue: 86.23
Position along Galactic rotation (pc)
Position toward Galactic center (pc)
300
200
100
0
−100
−200
−300
−400


c
Observed candidates
Hyades stars
Pleiades stars
Hyades BDs
Pleiades BDs
Fraction of catalogued Gaia counterparts
100%
80%
60%
40%
20%
0%
Gaia
counterparts
Quality
cuts
Distance +
search volume
Automatic
membership


d
Reference stars
Brown dwarfs
Faint-source quality losses not yet applied
Automatic recovery of simulated Gaia detections
60.3%
93.1%
97.0%
55.8%
96.0%
50.7%
Hyades
$R_{\rm birth} = 0.63$
Pleiades
$R_{\rm birth} = 0.63$
Pleiades
$R_{\rm birth} = 0.55$