

- 7.14 Suppose  $T \in \mathcal{L}(V)$  is self-adjoint,  $\lambda \in \mathbb{F}$ , and  $\epsilon > 0$ . Prove that if there exists  $v \in V$  such that  $\|v\| = 1$  and

$$\|Tv - \lambda v\| < \epsilon,$$

then  $T$  has an eigenvalue  $\lambda'$  such that  $|\lambda - \lambda'| < \epsilon$ .

**Proof.**

- 7.16 Give an example of an operator  $T$  on an inner product space such that  $T$  has an invariant subspace whose orthogonal complement is not invariant under  $T$ .

**Answer.**

- 7.17 Prove that the sum of any two positive operators on  $V$  is positive.

**Proof.** Suppose that  $S$  and  $T$  are positive operators on  $V$ . Since  $S$  and  $T$  are both self-adjoint, it follows immediately that  $S + T$  is self-adjoint because

$$(S + T)^* = S^* + T^* = S + T.$$

Now let  $v \in V$ . Thus

$$\begin{aligned}\langle (S + T)v, v \rangle &= \langle Sv + Tv, v \rangle \\ &= \langle Sv, v \rangle + \langle Tv, v \rangle \\ &\geq 0, \quad \quad \quad [\text{Since } Sv \geq 0, Tv \geq 0]\end{aligned}$$

so that  $S + T$  is a positive operator. □

- 7.19 Suppose that  $T$  is a positive operator on  $V$ . Prove that  $T$  is invertible if and only if

$$\langle Tv, v \rangle > 0$$

for every  $v \in V \setminus \{0\}$ .

**Proof.** Suppose first that  $T$  is invertible. Then it follows that  $T$  is injective, so that  $\text{null } T = \{0\}$ . Let  $v \in V$ . Now suppose that  $\langle Tv, v \rangle = 0$ . By the Spectral Theorem, there exists an orthonormal basis  $(e_1, \dots, e_n)$  of  $V$  consisting of eigenvectors of  $T$ . Let  $\lambda_1, \dots, \lambda_n$  denote the corresponding real ( $T$  is self-adjoint) and nonnegative (Theorem 7.27 (b)) eigenvalues. Since the eigenvectors are independent, they must be nonzero. Thus  $e_i \notin \text{null } T$ , so that  $0 \neq T(e_i) = \lambda_i e_i$ . That is, all the eigenvalues are positive. Now we have  $v = a_1 e_1 + \dots + a_n e_n$ , so that

$$\begin{aligned}0 &= \langle Tv, v \rangle \\ &= \langle T(a_1 e_1 + \dots + a_n e_n), a_1 e_1 + \dots + a_n e_n \rangle \\ &= \langle T(a_1 e_1) + \dots + T(a_n e_n), a_1 e_1 + \dots + a_n e_n \rangle \\ &= \langle a_1 \lambda_1 e_1 + \dots + a_n \lambda_n e_n, a_1 e_1 + \dots + a_n e_n \rangle \\ &= a_1 \overline{a_1} \lambda_1 \langle e_1, e_1 \rangle + \dots + a_n \overline{a_n} \lambda_n \langle e_n, e_n \rangle \\ &= |a_1|^2 \lambda_1 + \dots + |a_n|^2 \lambda_n.\end{aligned}$$

Since the eigenvalues are all positive, it must be the case  $a_1 = \cdots = a_n = 0$ , so that  $v = 0$ . So it follows that if  $v \in V$  is nonzero, we must have that  $\langle Tv, v \rangle > 0$ .

Conversely suppose that  $\langle Tv, v \rangle > 0$  for all nonzero  $v \in V$ . Let  $x \in \text{null } T$ . Then it follows that  $x$  is not nonzero because

$$0 = \langle 0, x \rangle = \langle Tx, x \rangle.$$

Thus  $x = 0$ ; that is  $\text{null } T = \{0\}$ . Thus  $T$  is injective (and surjective) and thus invertible.  $\square$

7.22 Prove that if  $S \in \mathcal{L}(\mathbb{R}^3)$  is an isometry, then there exists a nonzero vector  $x \in \mathbb{R}^3$  such that  $S^2x = x$ .

**Proof.** Assume that  $S \in \mathcal{L}(\mathbb{R}^3)$  is an isometry. Since  $\dim \mathbb{R}^3$  is odd, it follows by Theorem 7.38 that  $S$  has an eigenvalue of 1 or  $-1$ . Suppose first that  $S$  has an eigenvalue of 1. Then there exists a nonzero vector  $x$  such that  $Sx = x$ . Thus we have that

$$S^2x = S(Sx) = Sx = x.$$

Now suppose that  $S$  has an eigenvalue of  $-1$ . Then there exists a nonzero vector  $c$  such that  $Sc = -c$ . Let  $y = -c$ . Then we have that

$$S^2y = S(Sy) = S(S(-c)) = S(-S(c)) = S(c) = -c = y,$$

as desired.  $\square$