JOSEPH OKONOBOH
Mathematics
Cal State Long Beach

1. If Z is a standard normal variable, find

   (a) $P(Z^2 < 1)$ $\quad$ (b) $P(Z^2 > 3.84146)$.

   Solution.

   (a) We have that

   $$P(Z^2 < 1) = P(-1 < Z < 1) = 1 - 2 \cdot P(Z > 1) \approx 0.6826,$$

   and

   (b)

   $$P(Z^2 > 3.84146) = 2 \cdot P(Z > \sqrt{3.84146}) \approx 2 \cdot P(Z > 1.96) \approx 0.05.$$

2. If Y is a normal random variable with $\mu = 20$ and variance $\sigma^2 = 4$, i.e., $Y \sim N(20, 4)$, find

   (a) $P(16 \leq Y \leq 22)$ $\quad$ (b) $P(100 < 9Y - 80 < 145)$.

   Solution.

   (a) We have that

   $$\begin{aligned} P(16 \leq Y \leq 22) &= P\left(\frac{16-20}{2} \leq Z \leq \frac{22-20}{2}\right) \\ &= P(-2 \leq Z \leq 1) \\ &= 1 - [P(Z < -2) + P(Z > 1)] \\ &= 1 - [P(Z > 2) + P(Z > 1)] \\ &\approx 0.8185, \end{aligned}$$

   and

   (b)

   $$\begin{aligned} P(100 < 9Y - 80 < 145) &= P(20 < Y < 25) \\ &= P\left(\frac{20-20}{2} < Z < \frac{25-20}{2}\right) \\ &= P(0 < Z < 2.5) \\ &= P(Z > 0) - P(Z > 2.5) \\ &\approx 0.4938. \end{aligned}$$

3. The scores of a pre-employment test are normally distributed with mean $\mu = 70$ and standard deviation $\sigma = 5$. If only the top 1.5% of the applicants (based on their score on the pre-employment test) are to be considered, find the cut-off score (i.e., the value such that only 1.5% of the applicants score this value or higher).

   Solution. Let y be the cut-off score. Then we have that

   $$0.0015 = P(Y \geq y) = P\left(Z \geq \frac{y-70}{5}\right),$$

   so that $(y - 70)/5 \approx 2.97$; i.e., $y \approx 84.85$.

4. Using the fact that $\int_0^\infty e^{-y^2/2}dy = \sqrt{\frac{\pi}{2}}$, show that $\Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}$ by making the transformation $y = \frac{1}{2}x^2$.

   Proof. Using the substitution $y = \frac{1}{2}x^2$ we have that

$$\begin{aligned}\Gamma\left(\frac{1}{2}\right) &= \int_0^\infty y^{-\frac{1}{2}}e^{-y}dy \\ &= \int_0^\infty \frac{\sqrt{2}}{x}e^{-\frac{1}{2}x^2}x\,dx \\ &= \sqrt{2}\int_0^\infty e^{-\frac{1}{2}x^2}\,dx \\ &= \sqrt{2}\sqrt{\frac{\pi}{2}} = \sqrt{\pi}.\end{aligned}$$

5. If Y has an exponential distribution with $P(Y < 3) = 0.4512$, find
   (a) $E[Y]$ $\quad(b)$ $P(Y \geq 2)$.

   Solution.

   (a) We have that

$$\begin{aligned}0.4512 &= P(Y < 3) \\ &= P(Y \leq 3) \\ &= F(3) \\ &= \int_{-\infty}^{3} \frac{1}{\beta}e^{-\frac{y}{\beta}}dy \\ &= \int_0^3 \frac{1}{\beta}e^{-\frac{y}{\beta}}dy \\ &= -e^{-\frac{3}{\beta}} + 1,\end{aligned}$$

   so that $e^{-\frac{3}{\beta}} = 0.5488$; i.e., $\beta \approx 5$. Thus $E[Y] \approx 5$.

   (b)

$$\begin{aligned}P(Y \geq 2) &= 1 - P(Y < 2) \\ &= 1 - \int_0^2 \frac{1}{\beta}e^{-\frac{y}{\beta}}dy \\ &= e^{-\frac{2}{\beta}} \\ &\approx 0.6703.\end{aligned}$$

6. The length of time Y necessary to complete a key operation in the construction of houses has an exponential distribution with mean 10 hrs. The formula $C = 100 + 40Y + 3Y^2$ gives the cost C of completing the operation. Find the mean and variance of C.

Solution. First we want to find $E[Y^2]$. So

$$\begin{aligned}
E[Y^2] &= \frac{1}{10}\lim_{t\to\infty}\int_0^t y^2 e^{-\frac{y}{10}}\, dy \\
&= \lim_{t\to\infty}\left[-y^2 e^{-\frac{y}{10}}\Big|_0^t + 2\int_0^t y e^{-\frac{y}{10}}\, dy\right] && \text{[Integration by parts]} \\
&= 2\lim_{t\to\infty}\left[\int_0^t y e^{-\frac{y}{10}}\, dy\right] \\
&= 2\lim_{t\to\infty}\left[-10 y e^{-\frac{y}{10}}\Big|_0^t + 10\int_0^t e^{-\frac{y}{10}}\, dy\right] && \text{[Integration by parts]} \\
&= 200\lim_{t\to\infty}\left[\frac{1}{10}\int_0^t e^{-\frac{y}{10}}\, dy\right] \\
&= 200\cdot E[Y] = 2000.
\end{aligned}$$

Now the mean of C is given by $E[C]$ so that

$$\begin{aligned}
E[C] &= E[100 + 40Y + 3Y^2] \\
&= E[100] + 40E[Y] + 3E[Y^2] \\
&= 100 + 40\cdot 10 + 3\cdot 2000 \\
&= 6500,
\end{aligned}$$

and the variance of C, $V[Y]$, is $E[Y^2] - E[Y]^2 = 2000 - 100 = 1900$.

7. Suppose Y has density function $f(y) = ky^9 e^{-y/2}$, $y \geq 0$. Find
   (a) k.
   (b) $E[Y]$ and $V(Y)$.
   (c) $P(Y > 34.1696)$.
   (d) A value b such that $P(Y < b) = 0.10$.

Solution. By inspection we can see that f is the gamma distribution with $\alpha = 10$, $\beta = 2$.

(a) $k = \dfrac{1}{2^{10}\cdot\Gamma(10)} = \dfrac{1}{2^{10}\cdot 9!}$.

(b) $E[Y] = \alpha\beta = 20$ and $V(Y) = \alpha\beta^2 = 40$.

(c)

$$\begin{aligned}
P(Y > 34.1696) &= \frac{1}{2^{10}\cdot 9!}\int_{34.1696}^{\infty} y^9 e^{-y/2}dy \\
&= \frac{1}{2^{10}\cdot 9!}\int_{17.0848}^{\infty} 2^{10} z^9 e^{-z}dz \qquad \left[z = \frac{y}{2}\text{ substitution}\right] \\
&= \frac{1}{9!}\int_{17.0848}^{\infty} z^9 e^{-z}dz \\
&= \sum_{x=0}^{9} \frac{17.0848^x e^{-17.0848}}{x!} \\
&\approx 0.025.
\end{aligned}$$

(d) Suppose there exists b with $P(Y < b) = 0.10$, then we must have that

$$0.90 = P(Y \geq b) = P(Z \geq b/2),$$

and from Appendix 3, Table 3, we get $b/2 \approx 14$, so that $b \approx 28$.

8. The function $B(\alpha, \beta)$ is defined by $B(\alpha,\beta) = \displaystyle\int_0^1 y^{\alpha-1}(1-y)^{\beta-1}dy$.

   (a) Letting $y = \sin^2\theta$, show that $B(\alpha,\beta) = 2\displaystyle\int_0^{\pi/2} \sin^{2\alpha-1}\theta\cos^{2\beta-1}\theta\, d\theta$.

   (b) Write $\Gamma(\alpha)\Gamma(\beta)$ as a double integral, transform to polar coordinates, and then show that $B(\alpha,\beta) = \dfrac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}$.

9. Prove that the variance of a beta-distributed random variable with parameters α and β are given by

$$\sigma^2 = \frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)}.$$

10. Suppose Y has the density function $f(y) = k(y-2)^4(5-y)^6$, $2 \leq y \leq 5$. Find *(a)* k *(b)* $E[Y]$ and $V(Y)$.