

1. Consider the veracity or falsehood of each of the following statements. For bonus, argue for those that you believe are true while providing a counterexample for those that you believe are false.

- ① There is a field with 16 elements.
- ② In  $\mathbb{Z}[\sqrt{7}]$ ,  $\begin{pmatrix} 9 & 4 \\ 28 & 9 \end{pmatrix}$  is a prime.
- ③ The polynomial  $x^3 + ax^2 + bx + c \in \mathbb{Z}[x]$  where  $a$  is odd,  $b$  is even and  $c$  is odd, is always irreducible over  $\mathbb{Z}$ .
- ④  $\mathbb{Z}[\sqrt{-5}]$  is a UFD.
- ⑤ In  $\mathbb{Z}[\sqrt{7}]$ ,  $\begin{pmatrix} 8 & 3 \\ 21 & 8 \end{pmatrix}$  is a unit.

**Solution.**

- ① True.

**Example.** Let  $F = \mathbb{Z}_2[x]/(x^4 + x + 1)$ . That is,  $F$  consists of the polynomials in  $\mathbb{Z}_2[x]$  mod  $x^4 + x + 1$ . Thus  $F$  is the set of all polynomials of degree less than 4 with coefficients in  $\mathbb{Z}_2[x]$ , so that  $|F| = 16$ . Addition and multiplication in  $F$  are carried out mod  $(x^4 + x + 1)$ . It is clear that  $F$  is a commutative ring. Since

$$\begin{aligned} 1 \cdot 1 &= 1 \\ x(x^3 + 1) &= 1 \\ (x + 1)(x^3 + x^2 + x) &= 1 \\ x^2(x^3 + x^2 + 1) &= 1 \\ (x^2 + 1)(x^3 + x + 1) &= 1 \\ (x^2 + x)(x^2 + x + 1) &= 1 \\ x^3(x^3 + x^2 + x + 1) &= 1 \\ (x^3 + x^2)(x^3 + x) &= 1, \end{aligned}$$

it follows that every nonzero element of  $F$  has a multiplicative inverse, so that  $F$  is a field.

- ② True.

**Proof.** Let  $A = \begin{pmatrix} 9 & 4 \\ 28 & 9 \end{pmatrix}$ . Suppose  $A = BC$ . Then it follows that

$$-31 = \det(A) = \det(B) \det(C),$$

so that one of  $B$  and  $C$  has determinant  $\pm 1$ . Assume without loss of generality that  $\det(C) = \pm 1$ . It follows that  $C$  is invertible; thus  $AC^{-1} = B$ . That is  $A \mid B$ , so we can conclude that  $A$  is prime.  $\square$

③ True.

**Proof.** Let  $p(x) = x^3 + ax^2 + bx + c \in \mathbb{Z}[x]$ , where  $b$  is even and  $a$  and  $c$  are odd. Suppose to the contrary that  $p(x)$  is not irreducible. Then it follows that  $p(x)$  must have a root, say  $x_0 \in \mathbb{Z}$ . So  $0 = p(x_0) = x_0^3 + ax_0^2 + bx_0 + c$ . That is,  $x_0(-x_0^2 - ax_0 - b) = c$ . This says that  $x_0$  is a divisor of  $c$ . Then since  $c$  is odd, it must be the case that  $x_0$  is also odd. But then we must have that  $x_0^3$  is odd,  $ax_0^2$  is odd, and  $bx_0$  is even, so that  $x_0^3 + ax_0^2 + bx_0 + c$  is odd, a contradiction since  $x_0^3 + ax_0^2 + bx_0 + c = 0$  is even. Thus  $p(x)$  is irreducible.  $\square$

④ True.

**Proof.** First we want to show that the elements

$$A = \begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}, B = \begin{pmatrix} 3 & 0 \\ 0 & 3 \end{pmatrix}, C = \begin{pmatrix} 1 & 1 \\ -5 & 1 \end{pmatrix}, \text{ and } D = \begin{pmatrix} 1 & -1 \\ 5 & 1 \end{pmatrix}$$

are irreducible in  $\mathbb{Z}[\sqrt{-5}]$ . We shall only show that  $A$  and  $C$  are irreducible since the arguments for  $B$  and  $D$  are similar. Suppose  $A = A_1 A_2$ . Then it follows that

$$4 = \det(A) = \det(A_1) \det(A_2).$$

Observe that since we are in  $\mathbb{Z}[\sqrt{-5}]$ , it is impossible for the determinant of any matrix to be 2. Moreover, since the determinant of every matrix is nonnegative, we must have that either  $A_1$  or  $A_2$  has determinant of 1, so that one of  $A_1$  and  $A_2$  is a unit. Thus  $A$  is irreducible. Now suppose  $C = C_1 C_2$ . Then it follows that

$$6 = \det(C) = \det(C_1) \det(C_2).$$

No matrix has determinant 2 or 3 in  $\mathbb{Z}[\sqrt{-5}]$ . Thus one of  $C_1$  or  $C_2$  must be a unit, and it follows that  $C$  is irreducible. Since the units in  $\mathbb{Z}[\sqrt{-5}]$  are  $\pm 1$ , it follows that none of the irreducibles above are associates. It follows immediately that  $\mathbb{Z}[\sqrt{-5}]$  is not a UFD because we have the following two distinct factorizations into irreducibles:

$$\begin{pmatrix} 6 & 0 \\ 0 & 6 \end{pmatrix} = \begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix} \begin{pmatrix} 3 & 0 \\ 0 & 3 \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ -5 & 1 \end{pmatrix} \begin{pmatrix} 1 & -1 \\ 5 & 1 \end{pmatrix}.$$

$\square$

⑤ True. Since the determinant of the matrix in question is 1, it is a unit.

2. In a previous homework we encountered the integral domain  $R$  of  $2 \times 2$  matrices of the form  $A = \begin{pmatrix} a & b \\ -b & a-b \end{pmatrix}$  where  $a, b \in \mathbb{Z}$ . Do the following:

- ① Prove that no element of  $R$  can have a negative determinant.
- ② Find a nontrivial unit.
- ③ Find all units. Give an argument for your answer.

- ④ Find an element whose determinant is a prime.
- ⑤ Decide whether  $\begin{pmatrix} 3 & 0 \\ 0 & 3 \end{pmatrix}$  is irreducible or not. If not factor it into irreducibles.
- ⑥ Do the same for  $\begin{pmatrix} 5 & 0 \\ 0 & 5 \end{pmatrix}$ .
- ⑦ Do the same for  $\begin{pmatrix} 34 & 41 \\ -41 & -7 \end{pmatrix}$ .
- ⑧ Show that the element  $A = \begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}$  is a prime by showing that if  $MN \equiv 0 \pmod A$ , then either  $M \equiv 0 \pmod A$  or  $N \equiv 0 \pmod A$ .

**Solution.**

- ① **Proof.** Let  $A = \begin{pmatrix} a & b \\ -b & a-b \end{pmatrix} \in R$ . Since

$$\det(A) = a(a-b) + b^2 = \left(a - \frac{b}{2}\right)^2 + \frac{3}{4}b^2 \geq 0,$$

it follows that no element of  $R$  can have a negative determinant.  $\square$

- ② A nontrivial unit in  $R$  is  $\begin{pmatrix} 1 & 1 \\ -1 & 1-1 \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ -1 & 0 \end{pmatrix}$ .
- ③ Let  $A = \begin{pmatrix} a & b \\ -b & a-b \end{pmatrix} \in R$  be a unit. It follows by ① that

$$1 = \det(A) = a^2 + b^2 - ab;$$

thus we want integers  $a$  and  $b$  such that  $a^2 + b^2 - ab = 1$ . By completing the square we get that

$$a^2 + b^2 - ab = 1 \text{ iff } a = \frac{b}{2} \pm \sqrt{\frac{4-3b^2}{4}}.$$

For the discriminant to be positive, we must have that  $b = 0$  or  $|b| = 1$ . It follows that  $(a, b)$  is an integral solution of  $a^2 + b^2 - ab = 1$  iff

$$(a, b) \in \{(-1, 0), (1, 0), (0, 1), (1, 1), (0, -1), (-1, -1)\}.$$

Thus the group of units is

$$\left\{ I, -I, \begin{pmatrix} 0 & 1 \\ -1 & -1 \end{pmatrix}, \begin{pmatrix} 1 & 1 \\ -1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & -1 \\ 1 & 1 \end{pmatrix}, \begin{pmatrix} -1 & -1 \\ 1 & 0 \end{pmatrix} \right\}.$$

- ④ The element is  $\begin{pmatrix} 2 & 1 \\ -1 & 1 \end{pmatrix}$  has determinant 3.

- ⑤ The matrix  $\begin{pmatrix} 3 & 0 \\ 0 & 3 \end{pmatrix}$  is not irreducible since we have the following factorization into irreducibles:

$$\begin{pmatrix} 3 & 0 \\ 0 & 3 \end{pmatrix} = \begin{pmatrix} -1 & -2 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} 1 & 2 \\ -2 & -1 \end{pmatrix}.$$

The factors in the factorization above are irreducible since their determinants are prime.

- ⑥ Let  $B = \begin{pmatrix} 5 & 0 \\ 0 & 5 \end{pmatrix}$ . Claim that  $B$  is irreducible.

**Proof.** Suppose  $B = XY$ . Then it follows that

$$25 = \det(B) = \det(X) \det(Y).$$

Now suppose that  $\det(X) = 5$ . Then if we have that  $X = \begin{pmatrix} x & y \\ -y & x-y \end{pmatrix}$ , it follows that  $x^2 - xy + y^2 = 5$ . That is

$$x = \frac{y}{2} \pm \sqrt{\frac{20 - 3y^2}{4}}.$$

By observing the discriminant, we see that  $y$  can only take on values 0, 1, and 2. But  $x$  is not an integer for any of these values. Thus no matrix in  $R$  exists with determinant 5. It follows that one of  $X$  and  $Y$  must have determinant 1, so that this matrix is a unit; thus  $B$  is irreducible in  $R$ .  $\square$

- ⑦ The matrix  $\begin{pmatrix} 34 & 41 \\ -41 & -7 \end{pmatrix}$  is not irreducible since we have the following factorization into irreducibles:

$$\begin{pmatrix} 34 & 41 \\ -41 & -7 \end{pmatrix} = \begin{pmatrix} 2 & 1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} 4 & 1 \\ -1 & 3 \end{pmatrix} \begin{pmatrix} 7 & 3 \\ -3 & 4 \end{pmatrix}.$$

The factors in the factorization above are irreducible since their determinants are prime.

### 3. On Nilpotent Elements.

- ① Let  $R$  be a ring. An element  $m \in R$  is called nilpotent if  $m^k = 0$  for some positive integer  $k$ . Let  $r = 1 + m + m^2 + \cdots + m^{k-1}$ . Show  $r$  is invertible by finding its inverse.

- ② Exemplify ① by using the matrix  $\begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$ .

**Solution.**

- ① a

**BONUS.** Consider the integral domain  $R = \mathbb{Z}[\sqrt{3}]$ . Let  $A = \begin{pmatrix} 5 & 3 \\ 9 & 5 \end{pmatrix}$ . Decide if (1)-(4) are irreducible or not. Argue your case.

- (1)  $A$ .
- (2)  $\begin{pmatrix} 19 & 11 \\ 33 & 19 \end{pmatrix}$ .
- (3)  $\begin{pmatrix} 34 & 20 \\ 60 & 34 \end{pmatrix}$ .
- (4)  $\begin{pmatrix} 362 & 209 \\ 627 & 362 \end{pmatrix}$ .
- (5) Factor  $\begin{pmatrix} 69 & 0 \\ 0 & 69 \end{pmatrix}$  into irreducibles.
- (6) Show that  $A$  is prime by showing that  $R_A$  is a field. **Hint.** Show that if  $M \in R$  has even determinant, then  $A \mid M$ , and if  $M$  has odd determinant, then  $A \mid (M - I)$ .
- (7) Find a nontrivial common divisor of  $\begin{pmatrix} 89 & 53 \\ 159 & 89 \end{pmatrix}$  and  $\begin{pmatrix} 86 & 48 \\ 144 & 86 \end{pmatrix}$ , and show why it is a common divisor.
- (8) Find the greatest common divisor of  $\begin{pmatrix} 89 & 53 \\ 159 & 89 \end{pmatrix}$  and  $\begin{pmatrix} 86 & 48 \\ 144 & 86 \end{pmatrix}$ , and give reasons.
- (9) Find the lcm of  $\begin{pmatrix} 89 & 53 \\ 159 & 89 \end{pmatrix}$  and  $\begin{pmatrix} 86 & 48 \\ 144 & 86 \end{pmatrix}$ .

**More Bonus.** Argue every element is a product of irreducibles in  $R$ .

**Hard Bonus.** Argue every irreducible is prime.

**Solution.**

(1)