

1. A box contains 4 red balls, 3 white balls, 1 blue ball, and 2 green balls. Seven balls are selected with replacement. Find
- (a) the probability of selecting 3 red balls, 2 white balls, 1 blue ball, and 1 green ball.
 - (b) the expectation and variance of the number of red balls selected.
 - (c) the covariance of the number of red balls selected and the number of white balls selected

Solution.

- (a) $p(3r, 2w, 1b, 1g) = \frac{7!}{3!2!1!1!}(0.4)^3(0.3)^2(0.1)^1(0.2)^1 = 0.048384.$
 - (b) $E[\text{red}] = 7 \cdot 0.4 = 2.8, V(\text{red}) = 7 \cdot 0.4 \cdot 0.6 = 1.68.$
 - (c) $\text{Cov}(\text{red}, \text{white}) = -7 \cdot 0.4 \cdot 0.3 = -0.84.$
2. Suppose in a certain (large) community, 40% of the population is under 30 years of age, 30% of the population is between 30 and 50 years of age, 20% of the population is between 50 and 70 years of age, and 10% of the population is over 70. If eight people are randomly selected from the community, find the probability that
- (a) exactly four are under 30, exactly two are between 30 and 50, exactly one is between 50 and 70, and exactly one is over 70.
 - (b) exactly three are under 30 and exactly two are between 50 and 70.
 - (c) exactly five are under 50.

Solution.

- (a) $P = \frac{8!}{4!2!1!1!}(0.4)^4(0.3)^2(0.2)^1(0.1)^1 = 0.0387072.$
 - (b) $P = \frac{8!}{3!2!3!}(0.4)^3(0.2)^2(0.4)^3 = 0.0917504.$
 - (c) $P = \frac{8!}{5!3!}(0.7)^5(0.3)^3 = 0.25412.$
3. Suppose Y has a normal distribution with parameters $\mu = M$ and $\sigma^2 = 1$ where M has a gamma distribution with parameters $\alpha = 3$ and $\beta = 2$. Find (a) $E[Y]$ and (b) $V(Y)$.

Solution.

- (a) $E[Y] = E[E(Y|M)] = E[M] = 3 \cdot 2 = 6.$
 - (b) $V(Y) = E[V(Y|M)] + V[E(Y|M)] = E[\sigma^2] + V[M] = \sigma^2 E[1] + \alpha\beta = 13.$
4. Suppose Y is an exponential random variable with parameter Λ and Λ itself is a geometric random variable with parameter $p = 0.4$. Find $E[Y]$ and $V(Y)$.

Solution.

- (a) $E[Y] = E[E(Y|\Lambda)] = E[\Lambda] = 1/0.4 = 2.5.$

(b)

$$\begin{aligned} V(Y) &= E[V(Y|\Lambda)] + V(E[Y|\Lambda]) \\ &= E[\Lambda^2] + V(\Lambda) \\ &= 2V(\Lambda) + E[\Lambda]^2 \\ &= 2\frac{1-0.4}{0.4^2} + 2.5^2 = 13.75. \end{aligned}$$

5. Let Y be exponentially distributed with mean 3.

- (a) Use the method of distribution functions to find the density function of Y^2 .
- (b) Find a transformation G such that $G(Y)$ is uniformly distributed on $(1, 2)$.

Solution.

- (a) Let $U = Y^2$. We have that $F_U(u) = 0$ if $u \leq 0$. Now if $u > 0$, we have that

$$\begin{aligned} F_U(u) &= P(U \leq u) = P(Y^2 \leq u) \\ &= P(-\sqrt{u} \leq Y \leq \sqrt{u}) \\ &= \frac{1}{3} \int_{-\sqrt{u}}^{\sqrt{u}} e^{-y/3} dy \\ &= e^{\sqrt{u}/3} - e^{-\sqrt{u}/3}, \end{aligned}$$

so that

$$f_U(u) = \frac{dF_U(u)}{du} = \begin{cases} \frac{1}{6\sqrt{u}} (e^{\sqrt{u}/3} + e^{-\sqrt{u}/3}) & \text{if } u > 0 \\ 0 & \text{if otherwise} \end{cases}$$

- (b) The distribution of Y is given by

$$F_Y(y) = \begin{cases} 0 & \text{if } y < 0 \\ 1 - e^{-y/3} & \text{if } y \geq 0. \end{cases}$$

Let U be a uniformly distributed random variable on $(1, 2)$. Then the distribution of U is given by

$$F_U(u) = \begin{cases} 0 & \text{if } u < 1, \\ u - 1 & \text{if } 1 \leq u \leq 2, \\ 1 & \text{if } u > 2. \end{cases}$$

Let $1 \leq u < 2$. We want to find $H(u)$ such that $P(Y \leq H(u)) = u - 1$. Thus we have that

$$u - 1 = P(Y \leq H(u)) = 1 - e^{-H(u)/3}$$

so that $H(u) = -3\ln(2 - u)$. Now

$$\begin{aligned}u - 1 &= P(Y \leq H(u)) \\&= P(Y \leq -3\ln(2 - u)) \\&= P\left(-\frac{1}{3}Y \geq \ln(2 - u)\right) \\&= P\left(2 - e^{-1/3Y} \leq u\right)\end{aligned}$$

if $1 \leq u < 2$. So let $G(Y) = 2 - e^{-1/3Y}$.

6. Suppose Y has a probability density function

$$f(y) = \begin{cases} 3y^2 & \text{if } 0 \leq y \leq 1 \\ y^3 & \text{if otherwise} \end{cases}$$

Use the method of transformation to find the probability density functions of

a) $U_1 = 3Y - 1$ and b) $U_2 = Y^3$.

7. Let Y_1 and Y_2 have joint probability density function

$$f(y_1, y_2) = \begin{cases} 6(1 - y_2) & \text{if } 0 < y_1 \leq y_2 \leq 1 \\ 0 & \text{if otherwise} \end{cases}$$

Use the method of transformation to find the probability density functions of $U = Y_1/Y_2$.

8. Suppose Y_1 and Y_2 are two independent exponentially distributed random variables each with mean β . Use the method of moment generating functions to obtain the probability density function of $Y_1 + Y_2$.