

1. Consider the veracity or falsehood of each of the following statements. For bonus, argue for those that you believe are true while providing a counterexample for those that you believe are false.

- ① Every non-constant complex polynomial has a complex root.
- ② Conjugation of complex numbers is a field automorphism of the complex numbers.
- ③ Let $x, y \in R$, a finite ring. If $x * y = 1$, then $y * x = 1$ also.
- ④ There are exactly four quadratics in $\mathbb{Z}_2[x]$.
- ⑤ If $p(x)$ is a real polynomial, then it either has a real root or there is a quadratic polynomial with real coefficients that divides it.

Solution.

- ① True.

This follows from the Fundamental Theorem of Algebra.

- ② True.

Proof. Let \bar{a} denote the conjugate of the complex number a . We now want to show that

$$f : \mathbb{C} \rightarrow \mathbb{C}, c \mapsto \bar{c}$$

is an isomorphism. Let a_1 and a_2 be complex numbers. Since $\overline{a_1 a_2} = \bar{a}_1 \cdot \bar{a}_2$, and $\overline{a_1 + a_2} = \bar{a}_1 + \bar{a}_2$, it follows that

$$f(a_1 a_2) = f(a_1) f(a_2) \text{ and } f(a_1 + a_2) = f(a_1) + f(a_2),$$

so that f is an homomorphism. It now remains to show that f is a bijection. The map f must be surjective because $f(\bar{a}_1) = a_1$. Also if $f(a_1) = f(a_2)$, then the real parts of a_1 and a_2 must be equal. Similarly, their imaginary parts must be equal, so that $a_1 = a_2$. That is f is injective and we can conclude that it is a bijection. Thus f is a field automorphism. \square

- ③ True.

Proof. Let R be a finite ring, and consider $x, y \in R$ such that $x * y = 1$. The map $f : R \rightarrow R, r \mapsto r * x$ is bijective because for $r_1, r_2 \in R$ with $f(r_1) = f(r_2)$, we have that $r_1 * x = r_2 * x$. We then cancel x on both sides by multiplying each side on the right by y to get $r_1 = r_2$; thus f is injective, and since R is finite, we can conclude that f is also bijective. Thus there exists $r_3 \in R$ such that $r_3 * x = 1$. Multiply the preceding equality on the right by y to get $r_3 = y$. \square

- ④ False.

There are exactly 8 quadratics in $\mathbb{Z}_2[x]$, and they are

$$0, 1, x, x + 1, x^2, x^2 + 1, x^2 + x, x^2 + x + 1.$$

- ⑤ If $p(x)$ is 0, then it is trivially true. However, if $p(x)$ is a constant non-zero polynomial then it is not true. We shall now show that the statement is true if $p(x)$ is a non-constant real polynomial.

Proof. Consider the polynomial

$$p(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_0,$$

where each $a_i \in \mathbb{R}$, $a_n \neq 0$, and $n \geq 1$. By the Fundamental Theorem of Algebra, $p(x)$ has a root λ . If λ is real, then we are done. So assume that λ is a non-real complex number. Observe that the conjugate of λ , $\bar{\lambda}$, is also a root of $p(x)$ since

$$\begin{aligned} p(\bar{\lambda}) &= a_n \bar{\lambda}^n + a_{n-1} \bar{\lambda}^{n-1} + \cdots + a_0 \\ &= a_n \overline{\lambda^n} + a_{n-1} \overline{\lambda^{n-1}} + \cdots + a_0 \\ &= \overline{a_n \lambda^n} + \overline{a_{n-1} \lambda^{n-1}} + \cdots + \overline{a_0} & [\bar{a} = a \ \forall a \in \mathbb{R}] \\ &= \overline{a_n \lambda^n + a_{n-1} \lambda^{n-1} + \cdots + a_0} \\ &= \overline{p(\lambda)} = \overline{0} = 0. & [p(\lambda) = 0] \end{aligned}$$

Since λ is not real, we must have that $\lambda \neq \bar{\lambda}$. Thus the quadratic polynomial $(x - \lambda)(x - \bar{\lambda})$ divides $p(x)$. To complete the proof, we must show that this quadratic polynomial has real coefficients. Now we have that

$$(x - \lambda)(x - \bar{\lambda}) = x^2 - (\lambda + \bar{\lambda})x + \lambda\bar{\lambda} = x^2 - 2 \cdot \operatorname{Re}(\lambda)x + |\lambda|^2,$$

where $\operatorname{Re}(c)$ and $|c|$ denote the real part and magnitude of a complex number c . Thus the quadratic polynomial $(x - \lambda)(x - \bar{\lambda})$ has real coefficients. \square

2. On Complex & Real.

- ① Find a ring isomorphism (it has to be both additive and multiplicative) between \mathbb{C} and the subring $\mathcal{C} = \left\{ \begin{pmatrix} a & b \\ -b & a \end{pmatrix} : a, b \in \mathbb{R} \right\} \subseteq \mathcal{M}_2(\mathbb{R})$.
- ② In the notes we gave two descriptions of the quaternions:

$$\mathcal{Q} = \left\{ \begin{pmatrix} a & b & c & d \\ -b & a & -d & c \\ -c & d & a & -b \\ -d & -c & b & a \end{pmatrix} : a, b, c, d \in \mathbb{R} \right\} \text{ and } \mathcal{H} = \left\{ \begin{pmatrix} \alpha & \beta \\ -\bar{\beta} & \bar{\alpha} \end{pmatrix} : \alpha, \beta \in \mathbb{C} \right\}.$$

Find an isomorphism between these two rings (it has to be both additive and multiplicative).

Solution.

3. Let F be a field and consider the set R of all matrices of the form $\begin{pmatrix} a & b \\ -b & a-b \end{pmatrix}$ where $a, b \in F$. Do the following:

- ① Show R is closed under addition, subtraction and multiplication so it is a subring of $\mathcal{M}_2(F)$, the 2×2 matrices with entries in F .
- ② Find a positive integer n so that if we let the field $F = \mathbb{Z}_n$, then R will be an integral domain.
- ③ Find a positive integer n so that if we let the field $F = \mathbb{Z}_n$, then R will **NOT** be an integral domain.
- ④ Find a positive integer n so that if we let the field $F = \mathbb{Z}_n$, then R will be a field.
- ⑤ In any one of the situations ②, ③, or ④, find a unit of order bigger than 2. Just do one.
- ⑥ Suppose now that instead of F , we take $a, b \in \mathbb{Z}$, the integers. Prove it is an integral domain.

Bonus. Find $G(R)$, the group of units, in the case when the entries are integers (last situation), and find all elements of finite order in that group.

4. Consider the set R of matrices of the form $\frac{1}{2} \begin{pmatrix} a & b \\ 5b & a \end{pmatrix}$, $a, b \in \mathbb{Z}, a \equiv b \pmod{2}$.

- ① Show $I_2 \in R$.
- ② Show R is closed under addition, negation and multiplication so it is a subring of $\mathcal{M}_2(\mathbb{Q})$.
- ③ Compute the characteristic polynomial of any such matrix, and observe it is monic with integer coefficients.
- ④ Show there are infinitely many units in R .

Bonus. Find $\mathbb{I}(R)$, the group of units of R .