

ISyE6669 Deterministic Optimization
Homework 12
Fall 2021

Note: In all your answers, you need to give sufficient details and explanation how you derive or come to your solution.

1. In the lectures, we introduced the concept of a convex cone and gave three important examples: the nonnegative orthant cone \mathbb{R}_+^n , the second-order cone (SOC) \mathbb{L}^n , and the positive semidefinite cone (PSD) \mathbb{S}_+^n . Given a convex cone K , we can compare vectors using the cone K as $x \succeq_K y$ if and only if $x - y \in K$.

For example, if the cone is the nonnegative orthant $\mathbb{R}_+^n = \{(x_1, \dots, x_n) : x_i \geq 0, \forall i = 1, 2, \dots, n\}$. Then comparing vectors with respect to \mathbb{R}_+^n is the familiar componentwise comparison. That is, $x \succeq_{\mathbb{R}_+^n} y$ if and only if $x_i \geq y_i$ for each $i = 1, 2, \dots, n$. If the cone is \mathbb{L}^n or \mathbb{S}_+^n , the comparison is not componentwise comparison anymore.

As an exercise, let

$$x = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \quad y = \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix}.$$

Answer the following questions.

- (a) Using \mathbb{R}_+^3 , is it true that $x \succeq_{\mathbb{R}_+^3} y$? Explain why. Plot $x - y$ and \mathbb{R}_+^3 .

Solution

False.

$x \succeq_{\mathbb{R}_+^n} y \iff x_i \geq y_i$ for each $i = 1, 2, \dots, n$. However, $x_1 = 1 < 3 = y_1$.

$x - y = [-2, 0, 2]^T$ and \mathbb{R}_+^3 is plotted in Figure 1.

- (b) Using \mathbb{L}^3 , is it true that $x \succeq_{\mathbb{L}^3} y$? Explain why. Plot $x - y$ and \mathbb{L}^3 .

Solution

True.

$x \succeq_{\mathbb{L}^3} y \iff \sqrt{(x_1 - y_1)^2 + (x_2 - y_2)^2} \leq x_3 - y_3$.

$$\begin{aligned} \sqrt{(x_1 - y_1)^2 + (x_2 - y_2)^2} &= \sqrt{(1 - 3)^2 + (2 - 2)^2} \\ &= \sqrt{2^2 + 0^2} \\ &= \sqrt{4 + 0} \\ &= 2 \\ &= x_3 - y_3 \end{aligned}$$

$x - y$ and \mathbb{L}^3 is plotted in Figure 1.

- (c) Now define three matrices

$$A = \begin{bmatrix} -6 & 7 & 8 \\ 7 & -8 & 9 \\ 8 & 9 & -10 \end{bmatrix}, \quad B = \begin{bmatrix} 0 & 1 & 2 \\ 1 & 2 & 3 \\ 2 & 3 & 4 \end{bmatrix}, \quad C = \begin{bmatrix} -10 & 9 & 8 \\ 9 & -8 & 7 \\ 8 & 7 & -12 \end{bmatrix}.$$




Figure 1: $x - y$ and \mathbb{R}_+^3




Figure 2: $x - y$ and \mathbb{L}_+^3

Is it true that $A \preceq_{\mathbb{S}_+^3} B$? Is it true that $A \preceq_{\mathbb{S}_+^3} C$? Explain why. Remember A real symmetric matrix is positive semidefinite if and only if all of its eigenvalues are nonnegative.

Solution

i. $A \preceq_{\mathbb{S}_+^3} B$

False.

This is true if $B - A \succeq_{\mathbb{S}_+^3} 0$ i.e. $B - A$ is PSD and has nonnegative eigenvalues. Implemented the computation and tests in python and the resulting eigenvalues are $[-2.58, 14, 18.58]$. Since, not all the eigenvalues are nonnegative, the answer to the

question is false.

- ii. $A \succeq_{\mathbb{S}_+^3} C$
False.

Similarly, this is true if $A - C \succeq_{\mathbb{S}_+^3} 0$ i.e. $A - C$ is PSD and has nonnegative eigenvalues. The eigenvalues of $A - C$ are $[-1.76, 5.06, 2.69]$. Since, not all the eigenvalues are nonnegative, the answer to the question is false.

Notebook *HW12Q1.ipynb* is included in the Canvas submission.

2. There are n villages with known coordinates $v_1 = (x_1, y_1), v_2 = (x_2, y_2), \dots, v_n = (x_n, y_n)$, where x_i is the x-coordinate of the point v_i and y_i is the y-coordinate of the point v_i .

We want to locate a fire station $v = (x, y)$ such that the longest distance from the fire station to a possible fire in the villages is minimized. The distance is measured by the Euclidean distance. That is, the distance between two points $u = (a_1, b_1)$ and $w = (a_2, b_2)$ is given by $\|u - w\| = \sqrt{(a_1 - a_2)^2 + (b_1 - b_2)^2}$. Here, $\|u - w\|$ is also called the ℓ_2 norm of the vector $u - w$.

Formulate a second-order conic program (SOCP) for this problem. Note that the objective function of your SOCP must be a linear function of the variables and the constraints must be either linear constraints or second-order conic constraints.

Randomly generate the coordinates of $n = 20$ villages: generate the x-y coordinate of a village as a pair of random numbers uniformly distributed on the interval $[0, 10] \times [0, 10]$. Implement your SOCP model in CVXPY using the coordinates of the 20 villages that you generate. Solve the model and output the fire station's location and the minimum longest distance from the first station to a village.

Solution

The original problem

$$\min \max \|v - v_i\|$$

can be reformulated as a SOCP

$$\begin{aligned} \min \quad & z \\ \text{s.t.} \quad & \|v - v_i\| \leq z \end{aligned}$$

Figure 3 shows the implementation of the problem in python and the results. Notebook *HW12Q2.ipynb* is included in the Canvas submission.

```

import numpy as np
from cvxpy import *

z = Variable()
# fire station location
u = Variable(2)
# village locations
np.random.seed(0)
V = np.random.rand(20,2) * 10

max_dist = Minimize(z)
constraints = [norm(V - reshape(u, (1,2)), axis=1) <= z]
prob = Problem(max_dist, constraints)
optimal = prob.solve()
print('status:', prob.status)
print('fire station location:', u.value)
print('minimum longest distance:', optimal)
# print(V)

status: optimal
fire station location: [4.88041509 4.90032512]
minimum longest distance: 5.798487257610159

```

Figure 3: Q2 python implementation and results