

Discrete Mathematics

Relations and Functions

H. Turgut Uyar Ayşegül Gençata Yayimli Emre Harmancı

2001-2011

1 / 95

License



©2001-2011 T. Uyar, A. Yayimli, E. Harmancı

You are free:

- ▶ to Share — to copy, distribute and transmit the work
- ▶ to Remix — to adapt the work

Under the following conditions:

- ▶ Attribution — You must attribute the work in the manner specified by the author or licensor (but not in any way that suggests that they endorse you or your use of the work).
- ▶ Noncommercial — You may not use this work for commercial purposes.
- ▶ Share Alike — If you alter, transform, or build upon this work, you may distribute the resulting work only under the same or similar license to this one.

Legal code (the full license):

<http://creativecommons.org/licenses/by-nc-sa/3.0/>

2 / 95

Topics

Relations

Introduction
Relation Properties
Equivalence Relations

Functions

Introduction
Pigeonhole Principle
Recursion

3 / 95

Relation

Definition

relation: $\alpha \subseteq A \times B \times C \cdots \times N$

- ▶ **tuple:** an element of a relation
- ▶ $\alpha \subseteq A \times B$: *binary relation*
- ▶ $\alpha \subseteq A \times A$: *binary relation on A*
- ▶ **representations:**
 - ▶ by drawing
 - ▶ with a matrix

4 / 95

Relation Example

Example

$$A = \{a_1, a_2, a_3, a_4\}, B = \{b_1, b_2, b_3\}$$

$$\alpha = \{(a_1, b_1), (a_1, b_3), (a_2, b_2), (a_2, b_3), (a_3, b_1), (a_3, b_3), (a_4, b_1)\}$$



	b_1	b_2	b_3
a_1	1	0	1
a_2	0	1	1
a_3	1	0	1
a_4	1	0	0

$$M_\alpha = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}$$

5 / 95

Composite Relation

Definition

composite relation:

$$\alpha \subseteq A \times B \wedge \beta \subseteq B \times C$$

$$\Rightarrow \alpha\beta = \{(a, c) \mid a \in A, c \in C, \exists b \in B[a\alpha b \wedge b\beta c]\}$$

$$\triangleright M_{\alpha\beta} = M_\alpha \times M_\beta$$

6 / 95

Composite Relation Example

Example



7 / 95

Composite Relation Matrix Example

Example

$$M_\alpha = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix}$$

$$M_\beta = \begin{bmatrix} 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 1 & 1 & 0 \end{bmatrix}$$

$$M_{\alpha\beta} = \begin{bmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 \\ 1 & 1 & 1 & 0 \end{bmatrix}$$

8 / 95

Composite Relation Associativity

Theorem

$$(\alpha\beta)\gamma = \alpha(\beta\gamma) = \alpha\beta\gamma$$

9 / 95

Composite Relation Associativity

Proof.

$$\begin{aligned}(a, d) &\in (\alpha\beta)\gamma \\ \Leftrightarrow \exists c[(a, c) \in \alpha\beta \wedge (c, d) \in \gamma] \\ \Leftrightarrow \exists c[\exists b[(a, b) \in \alpha \wedge (b, c) \in \beta] \wedge (c, d) \in \gamma] \\ \Leftrightarrow \exists b[(a, b) \in \alpha \wedge \exists c[(b, c) \in \beta \wedge (c, d) \in \gamma]] \\ \Leftrightarrow \exists b[(a, b) \in \alpha \wedge (b, d) \in \beta\gamma] \\ \Leftrightarrow (a, d) &\in \alpha(\beta\gamma)\end{aligned}$$

□

10 / 95

Composite Relation Theorems

- ▶ $\alpha, \delta \subseteq A \times B \wedge \beta, \gamma \subseteq B \times C$
 - ▶ $\alpha(\beta \cup \gamma) = \alpha\beta \cup \alpha\gamma$
 - ▶ $\alpha(\beta \cap \gamma) \subseteq \alpha\beta \cap \alpha\gamma$
 - ▶ $(\alpha \cup \delta)\beta = \alpha\beta \cup \delta\beta$
 - ▶ $(\alpha \cap \delta)\beta \subseteq \alpha\beta \cap \delta\beta$
 - ▶ $(\alpha \subseteq \delta \wedge \beta \subseteq \gamma) \Rightarrow \alpha\beta \subseteq \delta\gamma$

11 / 95

Composite Relation Theorems

$$\alpha(\beta \cup \gamma) = \alpha\beta \cup \alpha\gamma.$$

$$\begin{aligned}(x, y) &\in \alpha(\beta \cup \gamma) \\ \Leftrightarrow \exists z[(x, z) \in \alpha \wedge (z, y) \in (\beta \cup \gamma)] \\ \Leftrightarrow \exists z[(x, z) \in \alpha \wedge ((z, y) \in \beta \vee (z, y) \in \gamma)] \\ \Leftrightarrow \exists z[[(x, z) \in \alpha \wedge (z, y) \in \beta] \\ \vee [(x, z) \in \alpha \wedge (z, y) \in \gamma]] \\ \Leftrightarrow (x, y) &\in \alpha\beta \vee (x, y) \in \alpha\gamma \\ \Leftrightarrow (x, y) &\in \alpha\beta \cup \alpha\gamma\end{aligned}$$

□

12 / 95

Converse Relation

Definition

$$\alpha^{-1} : \{(y, x) | (x, y) \in \alpha\}$$

$$\blacktriangleright M_{\alpha^{-1}} = M_{\alpha}^T$$

13 / 95

Converse Relation Theorems

- $\blacktriangleright (\alpha^{-1})^{-1} = \alpha$
- $\blacktriangleright (\alpha \cup \beta)^{-1} = \alpha^{-1} \cup \beta^{-1}$
- $\blacktriangleright (\alpha \cap \beta)^{-1} = \alpha^{-1} \cap \beta^{-1}$
- $\blacktriangleright \overline{\alpha^{-1}} = \overline{\alpha}^{-1}$
- $\blacktriangleright (\alpha - \beta)^{-1} = \alpha^{-1} - \beta^{-1}$
- $\blacktriangleright \alpha \subset \beta \Rightarrow \alpha^{-1} \subset \beta^{-1}$

14 / 95

Converse Relation Theorems

$$\overline{\alpha^{-1}} = \overline{\alpha}^{-1}.$$

$$\begin{aligned} & (x, y) \in \overline{\alpha^{-1}} \\ \Leftrightarrow & (y, x) \in \overline{\alpha} \\ \Leftrightarrow & (y, x) \notin \alpha \\ \Leftrightarrow & (x, y) \notin \alpha^{-1} \\ \Leftrightarrow & (x, y) \in \overline{\alpha^{-1}} \end{aligned}$$



15 / 95

Converse Relation Theorems

$$(\alpha \cap \beta)^{-1} = \alpha^{-1} \cap \beta^{-1}.$$

$$\begin{aligned} & (x, y) \in (\alpha \cap \beta)^{-1} \\ \Leftrightarrow & (y, x) \in (\alpha \cap \beta) \\ \Leftrightarrow & (y, x) \in \alpha \wedge (y, x) \in \beta \\ \Leftrightarrow & (x, y) \in \alpha^{-1} \wedge (x, y) \in \beta^{-1} \\ \Leftrightarrow & (x, y) \in \alpha^{-1} \cap \beta^{-1} \end{aligned}$$



16 / 95

Converse Relation Theorems

$$(\alpha - \beta)^{-1} = \alpha^{-1} - \beta^{-1}.$$

$$\begin{aligned} (\alpha - \beta)^{-1} &= (\alpha \cap \overline{\beta})^{-1} \\ &= \alpha^{-1} \cap \overline{\beta^{-1}} \\ &= \alpha^{-1} \cap \overline{\beta^{-1}} \\ &= \alpha^{-1} - \beta^{-1} \end{aligned}$$

□

17 / 95

Converse Composite Relation

Theorem

$$(\alpha\beta)^{-1} = \beta^{-1}\alpha^{-1}$$

Proof.

$$\begin{aligned} (c, a) &\in (\alpha\beta)^{-1} \\ \Leftrightarrow (a, c) &\in \alpha\beta \\ \Leftrightarrow \exists b[(a, b) &\in \alpha \wedge (b, c) \in \beta] \\ \Leftrightarrow \exists b[(c, b) &\in \beta^{-1} \wedge (b, a) \in \alpha^{-1}] \\ \Leftrightarrow (c, a) &\in \beta^{-1}\alpha^{-1} \end{aligned}$$

□

18 / 95

Converse Composite Relation Matrix

$$\begin{aligned} \blacktriangleright M_{(\alpha\beta)^{-1}} &= M_{\beta^{-1}} \times M_{\alpha^{-1}} \\ \blacktriangleright M_{\alpha\beta}^T &= M_{\beta}^T \times M_{\alpha}^T \end{aligned}$$

19 / 95

Converse Composite Relation Matrix Example

Example

$$\begin{aligned} M_{\alpha} &= \begin{vmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{vmatrix} & M_{\beta} &= \begin{vmatrix} 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 1 & 1 & 0 \end{vmatrix} \\ M_{\alpha\beta^{-1}} &= \begin{vmatrix} 1 & 0 & 0 & 0 & 1 \\ 1 & 1 & 1 & 0 & 1 \\ 0 & 1 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 & 0 \end{vmatrix} \end{aligned}$$

20 / 95

Binary Relation Properties

- ▶ $\alpha \subseteq A \times A$
- ▶ $\alpha\alpha : \alpha^2$
 - ▶ $\alpha\alpha \dots \alpha : \alpha^n$
- ▶ *identity relation*: $E = \{(x, x) | x \in A\}$
- ▶ properties: reflexivity, symmetry, transitivity

21 / 95

Reflexivity

reflexive

$$\alpha \subseteq A \times A$$

$$\forall a [a\alpha a]$$

▶ nonreflexive:

$$\exists a [\neg(a\alpha a)]$$

▶ irreflexive:

$$\forall a [\neg(a\alpha a)]$$

22 / 95

Reflexivity Examples

Example

$$\mathcal{R}_1 \subseteq \{1, 2\} \times \{1, 2\}$$

$$\mathcal{R}_1 = \{(1, 1), (2, 2)\}$$

- ▶ \mathcal{R}_1 is reflexive

Example

$$\mathcal{R}_2 \subseteq \{1, 2, 3\} \times \{1, 2, 3\}$$

$$\mathcal{R}_2 = \{(1, 1), (2, 2)\}$$

- ▶ \mathcal{R}_2 is nonreflexive

23 / 95

Reflexivity Examples

Example

$$\mathcal{R} \subseteq \{1, 2, 3\} \times \{1, 2, 3\}$$

$$\mathcal{R} = \{(1, 2), (2, 1), (2, 3)\}$$

- ▶ \mathcal{R} is irreflexive

24 / 95

Reflexivity Examples

Example

$$\mathcal{R} \subseteq \mathbb{Z} \times \mathbb{Z}$$

$$(a, b) \in \mathcal{R} \equiv ab \geq 0$$

- \mathcal{R} is reflexive

25 / 95

Symmetry

symmetric

$$\alpha \subseteq A \times A$$

$$\forall a, b [(a = b) \vee (a\alpha b \wedge b\alpha a) \vee (\neg(a\alpha b) \wedge \neg(b\alpha a))]$$

$$\forall a, b [(a = b) \vee (a\alpha b \leftrightarrow b\alpha a)]$$

- asymmetric:

$$\exists a, b [(a \neq b) \wedge (a\alpha b \wedge \neg(b\alpha a)) \vee (\neg(a\alpha b) \wedge b\alpha a)]$$

- antisymmetric:

$$\forall a, b [(a = b) \vee \neg(a\alpha b) \vee \neg(b\alpha a)]$$

$$\Leftrightarrow \forall a, b [\neg(a\alpha b \wedge b\alpha a) \vee (a = b)]$$

$$\Leftrightarrow \forall a, b [(a\alpha b \wedge b\alpha a) \rightarrow (a = b)]$$

26 / 95

Symmetry Examples

Example

$$\mathcal{R} \subseteq \{1, 2, 3\} \times \{1, 2, 3\}$$

$$\mathcal{R} = \{(1, 2), (2, 1), (2, 3)\}$$

- \mathcal{R} is asymmetric

27 / 95

Symmetry Examples

Example

$$\mathcal{R} \subseteq \mathbb{Z} \times \mathbb{Z}$$

$$(a, b) \in \mathcal{R} \equiv ab \geq 0$$

- \mathcal{R} is symmetric

28 / 95

Symmetry Examples

Example

$$\mathcal{R} \subseteq \{1, 2, 3\} \times \{1, 2, 3\}$$

$$\mathcal{R} = \{(1, 1), (2, 2)\}$$

- \mathcal{R} is both symmetric and antisymmetric

Transitivity

transitive

$$\alpha \subseteq A \times A$$

$$\forall a, b, c [(a\alpha b \wedge b\alpha c) \rightarrow (a\alpha c)]$$

► nontransitive:

$$\exists a, b, c [(a\alpha b \wedge b\alpha c) \wedge \neg(a\alpha c)]$$

► antitransitive:

$$\forall a, b, c [(a\alpha b \wedge b\alpha c) \rightarrow \neg(a\alpha c)]$$

Transitivity Examples

Example

$$\mathcal{R} \subseteq \{1, 2, 3\} \times \{1, 2, 3\}$$

$$\mathcal{R} = \{(1, 2), (2, 1), (2, 3)\}$$

- \mathcal{R} is antitransitive

Transitivity Examples

Example

$$\mathcal{R} \subseteq \mathbb{Z} \times \mathbb{Z}$$

$$(a, b) \in \mathcal{R} \equiv ab \geq 0$$

- \mathcal{R} is nontransitive

Converse Relation Properties

Theorem

The reflexivity, symmetry and transitivity properties are preserved in the converse relation.

Closure

- ▶ reflexive closure:

$$r_\alpha = \alpha \cup E$$

- ▶ symmetric closure:

$$s_\alpha = \alpha \cup \alpha^{-1}$$

- ▶ transitive closure:

$$t_\alpha = \bigcup_{j=1 \dots n} \alpha^j = \alpha \cup \alpha^2 \cup \alpha^3 \cup \dots \cup \alpha^n$$

Special Relations

predecessor - successor

$$\mathcal{R} \subseteq \mathbb{Z} \times \mathbb{Z}$$

$$(a, b) \in \mathcal{R} \equiv a - b = 1$$

- ▶ irreflexive
- ▶ antisymmetric
- ▶ antitransitive

Special Relations

adjacency

$$\mathcal{R} \subseteq \mathbb{Z} \times \mathbb{Z}$$

$$(a, b) \in \mathcal{R} \equiv |a - b| = 1$$

- ▶ irreflexive
- ▶ symmetric
- ▶ antitransitive

Special Relations

strict order

$$\mathcal{R} \subseteq \mathbb{Z} \times \mathbb{Z}$$

$$(a, b) \in \mathcal{R} \equiv a < b$$

- ▶ irreflexive
- ▶ antisymmetric
- ▶ transitive

37 / 95

Special Relations

partial order

$$\mathcal{R} \subseteq \mathbb{Z} \times \mathbb{Z}$$

$$(a, b) \in \mathcal{R} \equiv a \leq b$$

- ▶ reflexive
- ▶ antisymmetric
- ▶ transitive

38 / 95

Special Relations

preorder

$$\mathcal{R} \subseteq \mathbb{Z} \times \mathbb{Z}$$

$$(a, b) \in \mathcal{R} \equiv |a| \leq |b|$$

- ▶ reflexive
- ▶ asymmetric
- ▶ transitive

39 / 95

Special Relations

limited difference

$$\mathcal{R} \subseteq \mathbb{Z} \times \mathbb{Z}$$

$$(a, b) \in \mathcal{R} \equiv |a - b| \leq m$$

- ▶ reflexive
- ▶ symmetric
- ▶ nontransitive

40 / 95

Special Relations

comparability

$$\mathcal{R} \subseteq \mathbb{U} \times \mathbb{U}$$

$$(a, b) \in \mathcal{R} \equiv (a \subseteq b) \vee (b \subseteq a)$$

- ▶ reflexive
- ▶ symmetric
- ▶ nontransitive

41 / 95

Special Relations

brotherhood

- ▶ irreflexive
- ▶ symmetric
- ▶ transitive

- ▶ can a relation be symmetric, transitive and irreflexive?

42 / 95

Compatible Relations

Definition

compatible relation: γ

- ▶ reflexive
- ▶ symmetric

- ▶ undirected graph
- ▶ matrix representation as a triangle matrix

- ▶ $\alpha\alpha^{-1}$ is a compatible relation

43 / 95

Compatible Relation Example

Example

$$A = \{a_1, a_2, a_3, a_4\}$$

$$\mathcal{R} = \{$$

$$(a_1, a_1), (a_2, a_2),$$

$$(a_3, a_3), (a_4, a_4),$$

$$(a_1, a_2), (a_2, a_1),$$

$$(a_2, a_4), (a_4, a_2),$$

$$(a_3, a_4), (a_4, a_3)$$

$$\}$$



$$\begin{vmatrix} 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 1 & 1 & 1 \end{vmatrix}$$

$$\begin{vmatrix} 1 & & & \\ 0 & 0 & & \\ 0 & 1 & 1 & \end{vmatrix}$$

44 / 95

Compatible Relation Example

Example ($\alpha\alpha^{-1}$)

A: persons, B: languages

$A = \{a_1, a_2, a_3, a_4, a_5, a_6\}$

$B = \{b_1, b_2, b_3, b_4, b_5\}$

$\alpha \subseteq A \times B$

$$M_\alpha = \begin{bmatrix} 1 & 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 & 0 \end{bmatrix}$$

$$M_{\alpha^{-1}} = \begin{bmatrix} 1 & 1 & 0 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \end{bmatrix}$$

45 / 95

Compatible Relation Example

Example ($\alpha\alpha^{-1}$)

$\alpha\alpha^{-1} \subseteq A \times A$

$$M_{\alpha\alpha^{-1}} = \begin{bmatrix} 1 & 1 & 0 & 1 & 0 & 1 \\ 1 & 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 & 0 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 & 1 & 0 \\ 1 & 1 & 1 & 1 & 0 & 1 \end{bmatrix}$$



46 / 95

Compatible Class

Definition

compatibility block: $C \subseteq A$

$\forall a, b [a \in C \wedge b \in C \rightarrow a\gamma b]$

- ▶ **maximal compatibility block:**
not a subset of another compatibility block
- ▶ an element can be a member of more than one MCB
- ▶ **complete cover:** C_γ
set of all MCBs

47 / 95

Compatible Block Examples

Example ($\alpha\alpha^{-1}$)

- ▶ $C_1 = \{a_4, a_6\}$
- ▶ $C_2 = \{a_2, a_4, a_6\}$
- ▶ $C_3 = \{a_1, a_2, a_4, a_6\}$ (MCC)



$$C_\gamma(A) = \left\{ \begin{aligned} &\{a_1, a_2, a_4, a_6\}, \\ &\{a_3, a_4, a_6\}, \\ &\{a_4, a_5\} \end{aligned} \right\}$$

48 / 95

Equivalence Relations

Definition

equivalence relation: ϵ

- ▶ reflexive
- ▶ symmetric
- ▶ transitive
- ▶ equivalence classes
- ▶ every element is a member of exactly one equivalence class
- ▶ complete cover: C_ϵ

49 / 95

Partitioning

- ▶ every equivalence relation *partitions* a set into equivalence classes
- ▶ every *partitioning* corresponds to an equivalence relation

50 / 95

Equivalence Relation Example

Example

$$\mathcal{R} \subseteq \mathbb{Z} \times \mathbb{Z}$$

$$(a, b) \in \mathcal{R} \equiv 5 \mid |a - b|$$

- ▶ $x \bmod 5$ partitions \mathbb{Z} into 5 equivalence classes

51 / 95

References

Required Text: Grimaldi

- ▶ Chapter 5: Relations and Functions
 - ▶ 5.1. Cartesian Products and Relations
- ▶ Chapter 7: Relations: The Second Time Around
 - ▶ 7.1. Relations Revisited: Properties of Relations
 - ▶ 7.4. Equivalence Relations and Partitions

Supplementary Text: O'Donnell, Hall, Page

- ▶ Chapter 10: Relations

52 / 95

Function

Definition

function: $f : X \rightarrow Y$

$$\forall x \in X \forall y_1, y_2 \in Y (x, y_1), (x, y_2) \in f \Rightarrow y_1 = y_2$$

- ▶ X : domain, Y : codomain (or range)
- ▶ $(x, y) \in f \equiv y = f(x)$
- ▶ y is image of x under f

Subset Image

Definition

subset image:

$$f : X \rightarrow Y \wedge X_1 \subseteq X$$

$$f(X_1) = \{y | y \in Y, x \in X_1 \wedge y = f(x)\}$$

Subset Image Example

Example

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x^2$$

- ▶ $f(\mathbb{Z}) = \{0, 1, 4, 9, 16, \dots\}$
- ▶ $A = \{-2, 1\}$
 $f(A) = \{1, 4\}$

One-to-one Function

Definition

$f : X \rightarrow Y$ is **one-to-one**:

$$\forall x_1, x_2 \in X f(x_1) = f(x_2) \Rightarrow x_1 = x_2$$

One-to-one Function Examples

Example

$$f: \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = 3x + 7$$

$$\begin{aligned} f(x_1) &= f(x_2) \\ \Rightarrow 3x_1 + 7 &= 3x_2 + 7 \\ \Rightarrow 3x_1 &= 3x_2 \\ \Rightarrow x_1 &= x_2 \end{aligned}$$

Counterexample

$$g: \mathbb{Z} \rightarrow \mathbb{Z}$$

$$g(x) = x^4 - x$$

$$\begin{aligned} g(0) &= 0^4 - 0 = 0 \\ g(1) &= 1^4 - 1 = 0 \end{aligned}$$

Onto Function

Definition

$f: X \rightarrow Y$ is **onto**:

$$\forall y \in Y \exists x \in X f(x) = y$$

$$\triangleright f(X) = Y$$

Onto Function Examples

Example

$$f: \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x^3$$

Counterexample

$$f: \mathbb{Z} \rightarrow \mathbb{Z}$$

$$f(x) = 3x + 1$$

Bijjective Function

Definition

$f: X \rightarrow Y$ is **bijjective**:

f is both one-to-one and onto

Subset Image Properties

- ▶ $f : A \rightarrow B \wedge A_1, A_2 \subseteq A$:
 - ▶ $f(A_1 \cup A_2) = f(A_1) \cup f(A_2)$
 - ▶ $f(A_1 \cap A_2) \subseteq f(A_1) \cap f(A_2)$
 - ▶ if f is one-to-one:
 $f(A_1 \cap A_2) = f(A_1) \cap f(A_2)$

61 / 95

Composite Function

Definition

$$f : X \rightarrow Y, g : Y \rightarrow Z$$

$$g \circ f : X \rightarrow Z$$

$$(g \circ f)(x) = g(f(x))$$

- ▶ is not commutative
- ▶ is associative:
 $f \circ (g \circ h) = (f \circ g) \circ h$

62 / 95

Composite Function Examples

Example (commutativity)

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x^2$$

$$g : \mathbb{R} \rightarrow \mathbb{R}$$

$$g(x) = x + 5$$

$$g \circ f : \mathbb{R} \rightarrow \mathbb{R}$$

$$(g \circ f)(x) = x^2 + 5$$

$$f \circ g : \mathbb{R} \rightarrow \mathbb{R}$$

$$(f \circ g)(x) = (x + 5)^2$$

63 / 95

Composite Function Theorems

Theorem

$$f : X \rightarrow Y, g : Y \rightarrow Z:$$

f is one-to-one \wedge g is one-to-one $\Rightarrow g \circ f$ is one-to-one

Proof.

$$\begin{aligned} (g \circ f)(a_1) &= (g \circ f)(a_2) \\ \Rightarrow g(f(a_1)) &= g(f(a_2)) \\ \Rightarrow f(a_1) &= f(a_2) \\ \Rightarrow a_1 &= a_2 \end{aligned}$$

□

64 / 95

Composite Function Theorems

Theorem

$$f : X \rightarrow Y, g : Y \rightarrow Z:$$

f is onto $\wedge g$ is onto $\Rightarrow g \circ f$ is onto

Proof.

$$\forall z \in Z \exists y \in Y g(y) = z$$

$$\forall y \in Y \exists x \in X f(x) = y$$

$$\Rightarrow \forall z \in Z \exists x \in X g(f(x)) = z$$

□

65 / 95

Identity Function

Definition

identity function: 1_X

$$1_X : X \rightarrow X$$

$$1_X(x) = x$$

66 / 95

Inverse Function

Definition

$f : X \rightarrow Y$ is **invertible**:

$$\exists f^{-1} : Y \rightarrow X \ f^{-1} \circ f = 1_X \wedge f \circ f^{-1} = 1_Y$$

► f^{-1} : **inverse** of function f

67 / 95

Inverse Function Examples

Example

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = 2x + 5$$

$$f^{-1} : \mathbb{R} \rightarrow \mathbb{R}$$

$$f^{-1}(x) = \frac{x-5}{2}$$

$$(f^{-1} \circ f)(x) = f^{-1}(f(x)) = f^{-1}(2x + 5) = \frac{(2x+5)-5}{2} = \frac{2x}{2} = x$$

$$(f \circ f^{-1})(x) = f(f^{-1}(x)) = f\left(\frac{x-5}{2}\right) = 2\frac{x-5}{2} + 5 = (x-5) + 5 = x$$

68 / 95

Inverse Function

Theorem

If a function is invertible, its inverse is unique.

Proof.

$$f : X \rightarrow Y$$

$$g, h : Y \rightarrow X$$

$$g \circ f = 1_X \wedge f \circ g = 1_Y$$

$$h \circ f = 1_X \wedge f \circ h = 1_Y$$

$$h = h \circ 1_Y = h \circ (f \circ g) = (h \circ f) \circ g = 1_X \circ g = g \quad \square$$

69 / 95

Invertible Function

Theorem

A function is invertible if and only if it is one-to-one and onto.

70 / 95

Invertible Function

If invertible then one-to-one.

$$f : A \rightarrow B$$

$$\begin{aligned} f(a_1) &= f(a_2) \\ \Rightarrow f^{-1}(f(a_1)) &= f^{-1}(f(a_2)) \\ \Rightarrow (f^{-1} \circ f)(a_1) &= (f^{-1} \circ f)(a_2) \\ \Rightarrow 1_A(a_1) &= 1_A(a_2) \\ \Rightarrow a_1 &= a_2 \end{aligned}$$

\square

If invertible then onto.

$$f : A \rightarrow B$$

$$\begin{aligned} &b \\ &= 1_B(b) \\ &= (f \circ f^{-1})(b) \\ &= f(f^{-1}(b)) \end{aligned}$$

\square

71 / 95

Invertible Function

If bijective then invertible.

$$f : A \rightarrow B$$

- ▶ f is onto $\Rightarrow \forall b \in B \exists a \in A f(a) = b$
- ▶ let $g : B \rightarrow A$ be defined by $a = g(b)$
- ▶ is it possible that $g(b) = a_1 \neq a_2 = g(b)$?
- ▶ this would mean: $f(a_1) = b = f(a_2)$
- ▶ but: f is one-to-one

\square

72 / 95

Permutations

- ▶ permutation: a bijective function on a set

$$\begin{pmatrix} a_1 & a_2 & \dots & a_n \\ p(a_1) & p(a_2) & \dots & p(a_n) \end{pmatrix}$$

- ▶ $n!$ permutations can be defined in a set of n elements

73 / 95

Permutation Examples

Example

$$A = \{1, 2, 3\}$$

$$p_1 = \begin{pmatrix} 1 & 2 & 3 \\ 1 & 2 & 3 \end{pmatrix} \quad p_2 = \begin{pmatrix} 1 & 2 & 3 \\ 1 & 3 & 2 \end{pmatrix}$$

$$p_3 = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 1 & 3 \end{pmatrix} \quad p_4 = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \end{pmatrix}$$

$$p_5 = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix} \quad p_6 = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{pmatrix}$$

74 / 95

Cyclic Permutation

- ▶ *cyclic permutation*:
 - ▶ a subset of elements form a cycle
 - ▶ the remaining elements do not change

$$(a_i, a_j, a_k) = \begin{pmatrix} \dots & a_i & \dots & a_n & \dots & a_j & \dots & a_k & \dots \\ \dots & a_j & \dots & a_n & \dots & a_k & \dots & a_i & \dots \end{pmatrix}$$

- ▶ *transposition*: a cyclic permutation of length 2

75 / 95

Cyclic Permutation Examples

Example

$$A = \{1, 2, 3, 4, 5\}$$

$$(1, 3, 5) = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 3 & 2 & 5 & 4 & 1 \end{pmatrix}$$

76 / 95

Permutation Composition

- ▶ permutation composition is not commutative

Example

$$A = \{1, 2, 3, 4, 5\}$$

$$\begin{aligned}(4, 1, 3, 5) \circ (5, 2, 3) &= \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 3 & 2 & 5 & 1 & 4 \end{pmatrix} \circ \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 1 & 3 & 5 & 4 & 2 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 5 & 3 & 2 & 1 & 4 \end{pmatrix} \\ (5, 2, 3) \circ (4, 1, 3, 5) &= \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 1 & 3 & 5 & 4 & 2 \end{pmatrix} \circ \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 3 & 2 & 5 & 1 & 4 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 3 & 5 & 4 & 1 & 2 \end{pmatrix}\end{aligned}$$

77 / 95

Cyclic Permutation Composition

- ▶ all permutations that are not cyclic can be written as a composition of disjoint cyclic permutations

Example

$$A = \{1, 2, 3, 4, 5, 6, 7, 8\}$$

$$\begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ 3 & 4 & 6 & 5 & 2 & 1 & 8 & 7 \end{pmatrix} = (1, 3, 6) \circ (2, 4, 5) \circ (7, 8)$$

78 / 95

Transposition Composition

- ▶ all cyclic permutations can be written as a composition of transpositions

Example

$$A = \{1, 2, 3, 4, 5\}$$

$$(1, 2, 3, 4, 5) = (1, 2) \circ (1, 3) \circ (1, 4) \circ (1, 5)$$

79 / 95

Pigeonhole Principle

Definition

pigeonhole principle (Dirichlet drawers):

if m pigeons go into n holes and $m > n$

at least one hole contains more than one pigeon

- ▶ if $f : X \rightarrow Y \wedge |X| > |Y|$ then f cannot be one-to-one
- ▶ $\exists x_1, x_2 \in X \ x_1 \neq x_2 \wedge f(x_1) = f(x_2)$

80 / 95

Pigeonhole Principle Examples

Example

- ▶ In a room where there are 367 people, at least two persons have the same birthday.
- ▶ How many students should take an exam where the grades are between 0 and 100 so that two students have the same grade?

81 / 95

Generalized Pigeonhole Principle

Definition

generalized pigeonhole principle:

if m objects are distributed to n drawers
at least one of the drawers contains $\lceil m/n \rceil$ objects

Example

In a room where there are 100 people at least $\lceil 100/12 \rceil = 9$ persons were born in the same month.

82 / 95

Pigeonhole Principle Example

Theorem

There are two elements which total 10
in any subset of cardinality 6 of the set $S = \{1, 2, 3, \dots, 9\}$.

83 / 95

Pigeonhole Principle Example

Theorem

Let S be a set of positive integers smaller than or equal to 14,
with cardinality 6. The sums of the elements
in all nonempty subsets of S cannot be all different.

Proof Trial

$A \subseteq S$

s_A : sum of the elements of A

- ▶ holes:
 $1 \leq s_A \leq 9 + \dots + 14 = 69$
- ▶ pigeons: $2^6 - 1 = 63$

Proof.

look at the subsets for which
 $|A| \leq 5$

- ▶ holes:
 $1 \leq s_A \leq 10 + \dots + 14 = 60$
- ▶ pigeons: $2^6 - 2 = 62$

□

84 / 95

Pigeonhole Principle Example

Theorem

There is at least one pair of elements among 101 elements chosen from set $S = \{1, 2, 3, \dots, 200\}$ so that one of the elements of the pair divides the other.

Proof Method

- we first show that
 $\forall n \exists! p (n = 2^r p \wedge r \geq 0 \wedge \exists t \in \mathbb{Z} p = 2t + 1)$
- then, by using this theorem we prove the main theorem

85 / 95

Pigeonhole Principle Example

Theorem

$$\forall n \exists! p (n = 2^r p \wedge r \geq 0 \wedge \exists t \in \mathbb{Z} p = 2t + 1)$$

Proof of Existence.

$$n = 1: r = 0, p = 1$$

$$n = 2: r = 1, p = 1$$

$$n \leq k: n = 2^r p$$

$$n = k + 1:$$

$$n \text{ prime} : r = 0, p = n$$

$$\neg(n \text{ prime}) : n = n_1 n_2$$

$$n = 2^{r_1} p_1 \cdot 2^{r_2} p_2$$

$$n = 2^{r_1+r_2} \cdot p_1 p_2$$

Proof of Uniqueness.

if not unique:

$$n = 2^{r_1} p_1 = 2^{r_2} p_2$$

$$\Rightarrow 2^{r_1-r_2} p_1 = p_2$$

$$\Rightarrow 2 \mid p_2$$

□

□

86 / 95

Pigeonhole Principle Example

Theorem

There is at least one pair of elements among 101 elements chosen from set $S = \{1, 2, 3, \dots, 200\}$ so that one of the elements of the pair divides the other.

Proof.

- $T \subseteq S$, Assume that T is a subset of S that contains all odd elements of S : $|T| = 100$
- $f: S \rightarrow T, (s, t) \in f \Leftrightarrow s = 2^r t \wedge r \geq 0$
 - if 101 elements are chosen from S , at least two of them will have the same image in T : $f(s_1) = f(s_2) \Rightarrow 2^{m_1} t = 2^{m_2} t$

$$\frac{s_1}{s_2} = \frac{2^{m_1} t}{2^{m_2} t} = 2^{m_1 - m_2}$$

□

87 / 95

Recursive Functions

Definition

recursive function:

a function defined in terms of itself

$$f(n) = h(f(m))$$

- inductively defined function:**

the size reduced at every step of the recursion

$$f(n) = \begin{cases} k & n = 0 \\ h(f(n-1)) & n > 0 \end{cases}$$

88 / 95

Recursive Function Examples

Example

$$f_{91}(n) = \begin{cases} n - 10 & n > 100 \\ f_{91}(f_{91}(n + 11)) & n \leq 100 \end{cases}$$

89 / 95

Inductively Defined Function Examples

Example (factorial)

$$f(n) = \begin{cases} 1 & n = 0 \\ n \cdot f(n - 1) & n > 0 \end{cases}$$

Example (function power)

$$f^n = \begin{cases} f & n = 1 \\ f \circ f^{n-1} & n > 1 \end{cases}$$

90 / 95

Euclid Algorithm

Example (greatest common divisor)

$$333 = 3 \cdot 84 + 81$$

$$84 = 1 \cdot 81 + 3$$

$$81 = 27 \cdot 3 + 0$$

$$\gcd(333, 84) = 3$$

$$\gcd(a, b) = \begin{cases} b & b|a \\ \gcd(b, a \bmod b) & b \nmid a \end{cases}$$

91 / 95

Fibonacci Series

Fibonacci series

$$F_n = \text{fib}(n) = \begin{cases} 1 & n = 1 \\ 1 & n = 2 \\ \text{fib}(n - 1) + \text{fib}(n - 2) & n > 2 \end{cases}$$

$$\begin{array}{ccccccccccc} F_1 & F_2 & F_3 & F_4 & F_5 & F_6 & F_7 & F_8 & \dots \\ 1 & 1 & 2 & 3 & 5 & 8 & 13 & 21 & \dots \end{array}$$

92 / 95

Fibonacci Series

Theorem

$$\sum_{i=1}^n F_i^2 = F_n \cdot F_{n+1}$$

Proof.

$$n = 2 : \quad \sum_{i=1}^2 F_i^2 = F_1^2 + F_2^2 = 1 + 1 = 1 \cdot 2 = F_2 \cdot F_3$$

$$n = k : \quad \sum_{i=1}^k F_i^2 = F_k \cdot F_{k+1}$$

$$\begin{aligned} n = k + 1 : \quad \sum_{i=1}^{k+1} F_i^2 &= \sum_{i=1}^k F_i^2 + F_{k+1}^2 \\ &= F_k \cdot F_{k+1} + F_{k+1}^2 \\ &= F_{k+1} \cdot (F_k + F_{k+1}) \\ &= F_{k+1} \cdot F_{k+2} \end{aligned}$$

□

93 / 95

Ackermann Function

Ackermann function

$$ack(x, y) = \begin{cases} y + 1 & x = 0 \\ ack(x - 1, 1) & y = 0 \\ ack(x - 1, ack(x, y - 1)) & x > 0 \wedge y > 0 \end{cases}$$

94 / 95

References

Required Text: Grimaldi

- ▶ Chapter 5: Relations and Functions
 - ▶ 5.2. Functions: Plain and One-to-One
 - ▶ 5.3. Onto Functions: Stirling Numbers of the Second Kind
 - ▶ 5.5. The Pigeonhole Principle
 - ▶ 5.6. Function Composition and Inverse Functions

Supplementary Text: O'Donnell, Hall, Page

- ▶ Chapter 11: Functions

95 / 95