

Discrete Mathematics

Propositions

H. Turgut Uyar Ayşegül Gençata Yayımlı Emre Harmancı

2001-2012

1 / 67

Licence



©2001-2012 T. Uyar, A. Yayımlı, E. Harmancı

You are free:

- ▶ to Share — to copy, distribute and transmit the work
- ▶ to Remix — to adapt the work

Under the following conditions:

- ▶ Attribution — You must attribute the work in the manner specified by the author or licensor (but not in any way that suggests that they endorse you or your use of the work).
- ▶ Noncommercial — You may not use this work for commercial purposes.
- ▶ Share Alike — If you alter, transform, or build upon this work, you may distribute the resulting work only under the same or similar license to this one.

Legal code (the full license):

<http://creativecommons.org/licenses/by-nc-sa/3.0/>

2 / 67

Topics

Propositions

Introduction
Compound Propositions
Well-Formed Formulas
Metalanguage

Propositional Calculus

Introduction
Laws of Logic
Rules of Inference

3 / 67

Proposition

Definition

proposition (or **statement**):

a declarative sentence that is either true or false

- ▶ **law of the excluded middle**:
a proposition cannot be partially true or partially false
- ▶ **law of contradiction**:
a proposition cannot be both true and false

4 / 67

Proposition Examples

Example (proposition)

- ▶ The Moon revolves around the Earth.
- ▶ Elephants can fly.
- ▶ $3 + 8 = 11$

Example (not proposition)

- ▶ What time is it?
- ▶ Ali, throw the ball!
- ▶ $x < 43$

5 / 67

Proposition Variable

Definition

proposition variable:

a name that represents the proposition

- ▶ can take on the values *True* (*T*) or *False* (*F*)

Example

- ▶ p_1 : The Moon revolves around the Earth. (*T*)
- ▶ p_2 : Elephants can fly. (*F*)
- ▶ p_3 : $3 + 8 = 11$ (*T*)

6 / 67

Compound Propositions

- ▶ **compound propositions** are obtained by
 - ▶ negating a proposition, or
 - ▶ combining two or more propositions using **logical connectives**
- ▶ **primitive propositions** can not be decomposed into smaller units
- ▶ **truth table**:
a table that lists the truth value of the compound proposition for all possible values of its primitive propositions

7 / 67

Negation (NOT)

Table: $\neg p$

p	$\neg p$
T	F
F	T

Example

- ▶ $\neg p_1$: The Moon does not revolve around the Earth.
 $\neg T$: *False*
- ▶ $\neg p_2$: Elephants cannot fly.
 $\neg F$: *True*

8 / 67

Conjunction (AND)

Table: $p \wedge q$

p	q	$p \wedge q$
T	T	T
T	F	F
F	T	F
F	F	F

Example

- ▶ $p_1 \wedge p_2$: The Moon revolves around the Earth and elephants can fly.
 $T \wedge F$: *False*

9 / 67

Disjunction (OR)

Table: $p \vee q$

p	q	$p \vee q$
T	T	T
T	F	T
F	T	T
F	F	F

Example

- ▶ $p_1 \vee p_2$: The Moon revolves around the Earth or elephants can fly.
 $T \vee F$: *True*

10 / 67

Exclusive Disjunction (XOR)

Table: $p \underline{\vee} q$

p	q	$p \underline{\vee} q$
T	T	F
T	F	T
F	T	T
F	F	F

Example

- ▶ $p_1 \underline{\vee} p_2$: Either the Moon revolves around the Earth or elephants can fly.
 $T \underline{\vee} F$: *True*

11 / 67

Implication (IF)

Table: $p \rightarrow q$

p	q	$p \rightarrow q$
T	F	F
F	T	T
F	F	T
T	T	T

- ▶ p : **hypothesis**
- ▶ q : **conclusion**
- ▶ read:
 - ▶ if p then q
 - ▶ p is sufficient for q
 - ▶ q is necessary for p
- ▶ $\neg p \vee q$

12 / 67

Implication Examples

Example

- ▶ p_4 : $3 < 8$, p_5 : $3 < 14$, p_6 : $3 < 2$
- ▶ p_7 : The Sun revolves around the Earth.
- ▶ $p_4 \rightarrow p_5$: If 3 is less than 8, then 3 is less than 14.
 $T \rightarrow T$: *True*
- ▶ $p_4 \rightarrow p_6$: If 3 is less than 8, then 3 is less than 2.
 $T \rightarrow F$: *False*
- ▶ $p_2 \rightarrow p_1$: If elephants can fly then the Moon revolves around the Earth.
 $F \rightarrow T$: *True*
- ▶ $p_2 \rightarrow p_7$: If elephants can fly then the Sun revolves around the Earth.
 $F \rightarrow F$: *True*

13 / 67

Implication Examples

Example

- ▶ "If I weigh over 70 kg, then I will exercise."

Table: $p \rightarrow q$

p	q	$p \rightarrow q$
T	T	T
T	F	F
F	T	T
F	F	T

- ▶ p : I weigh over 70 kg.
- ▶ q : I exercise.
- ▶ when is this claim false?

14 / 67

Biconditional (IFF)

Table: $p \leftrightarrow q$

p	q	$p \leftrightarrow q$
T	F	F
T	T	T
F	T	F
F	F	T

- ▶ read:
 - ▶ p if and only if q
 - ▶ p is necessary and sufficient for q
- ▶ $(p \rightarrow q) \wedge (q \rightarrow p)$
- ▶ $\neg(p \vee q)$

15 / 67

Example

Example

- ▶ The parent tells the child:
"If you do your homework, you can play computer games."
- ▶ s : The child does her homework.
- ▶ t : The child plays computer games.
- ▶ which one does the parent mean?
 - ▶ $s \rightarrow t$
 - ▶ $\neg s \rightarrow \neg t$
 - ▶ $s \leftrightarrow t$

16 / 67

Well-Formed Formula

syntax

- ▶ which rules will be used to form compound propositions?
- ▶ formula that obeys these rules: **well-formed formula** (WFF)

semantics

- ▶ *interpretation*: calculating the value of a compound proposition by assigning values to its primitive propositions
- ▶ truth table: all interpretations of a proposition

17 / 67

Formula Examples

Example (not well-formed)

- ▶ $\vee p$
- ▶ $p \wedge \neg$
- ▶ $p \neg \wedge q$

18 / 67

Precedence

1. \neg
2. \wedge
3. \vee
4. \rightarrow
5. \leftrightarrow

► parentheses are used to change precedence

19 / 67

Precedence Examples

Example

- s : Phyllis goes out for a walk.
- t : The Moon is out.
- u : It is snowing.
- what do the following WFFs mean?
 - $t \wedge \neg u \rightarrow s$
 - $t \rightarrow (\neg u \rightarrow s)$
 - $\neg(s \leftrightarrow (u \vee t))$
 - $\neg s \leftrightarrow u \vee t$

20 / 67

Formula Attributes

1. *tautology*: True for all interpretations
2. *contradiction*: False for all interpretations
3. *valid*: True for some interpretations

21 / 67

Tautology Example

Example

Table: $p \wedge (p \rightarrow q) \rightarrow q$

p	q	$p \rightarrow q$ (A)	$p \wedge A$ (B)	$B \rightarrow q$
T	T	T	T	T
T	F	F	F	T
F	T	T	F	T
F	F	T	F	T

22 / 67

Contradiction Example

Example

Table: $p \wedge (\neg p \wedge q)$

p	q	$\neg p$	$\neg p \wedge q$ (A)	$p \wedge A$
T	T	F	F	F
T	F	F	F	F
F	T	T	T	F
F	F	T	F	F

23 / 67

Metalanguage

Definition

target language:
the language being worked on

Definition

metalanguage:
the language used when talking about the properties of the target language

- validity, contradiction and tautology are defined in the metalanguage

24 / 67

Metalanguage Examples

Example (for a Turk who is learning English)

- ▶ target language: English
- ▶ metalanguage: Turkish

Example (in an introductory programming course)

- ▶ target language: C, Python, Java, ...
- ▶ metalanguage: English, Turkish, ...

25 / 67

Metalogic

- ▶ $P_1, P_2, \dots, P_n \vdash Q$
There is a proof which infers the conclusion Q from the assumptions P_1, P_2, \dots, P_n .
- ▶ $P_1, P_2, \dots, P_n \models Q$
 Q must be true if P_1, P_2, \dots, P_n are all true.

26 / 67

Formal Systems

Definition

consistent: for all well-formed formulas P and Q
if $P \vdash Q$ then $P \models Q$

- ▶ each provable proposition is actually true

Definition

complete: for all well-formed formulas P and Q
if $P \models Q$ then $P \vdash Q$

- ▶ every true proposition can be proven

27 / 67

Gödel's Theorem

- ▶ Propositional logic is consistent and complete.

Gödel's Theorem

- ▶ Any logical system that is powerful enough to express ordinary arithmetic must be either inconsistent or incomplete.

28 / 67

Approaches in Propositional Calculus

1. semantic approach: *truth tables*
 - ▶ too complicated when the number of primitive statements grow
2. syntactic approach: *rules of inference*
 - ▶ obtaining new propositions from existing propositions using logical implications
3. axiomatic approach: *Boolean algebra*
 - ▶ substituting equivalent formulas in equations

29 / 67

Truth Table Example

Example ($p \rightarrow q$)

p	q	$p \rightarrow q$	$\neg q \rightarrow \neg p$	$q \rightarrow p$	$\neg p \rightarrow \neg q$
T	T	T	T	T	T
T	F	F	F	T	T
F	T	T	T	F	F
F	F	T	T	T	T

- ▶ *contrapositive:* $\neg q \rightarrow \neg p$
- ▶ *converse:* $q \rightarrow p$
- ▶ *inverse:* $\neg p \rightarrow \neg q$

30 / 67

Logical Equivalence

Definition

if $P \leftrightarrow Q$ is a tautology, then P and Q are **logically equivalent**:
 $P \Leftrightarrow Q$

31 / 67

Logical Equivalence Example

Example

$$\blacktriangleright \neg p \Leftrightarrow p \rightarrow F$$

Table: $\neg p \Leftrightarrow p \rightarrow F$

p	$\neg p$	$p \rightarrow F$ (A)	$\neg p \Leftrightarrow A$
T	F	F	T
F	T	T	T

32 / 67

Logical Equivalence Example

Example

$$\blacktriangleright p \rightarrow q \Leftrightarrow \neg p \vee q$$

Table: $(p \rightarrow q) \Leftrightarrow (\neg p \vee q)$

p	q	$p \rightarrow q$ (A)	$\neg p$	$\neg p \vee q$ (B)	$A \leftrightarrow B$
T	T	T	F	T	T
T	F	F	F	F	T
F	T	T	T	T	T
F	F	T	T	T	T

33 / 67

Laws of Logic

Double Negation (DN)

$$\neg(\neg p) \Leftrightarrow p$$

Commutativity (Co)

$$p \wedge q \Leftrightarrow q \wedge p$$

$$p \vee q \Leftrightarrow q \vee p$$

Associativity (As)

$$(p \wedge q) \wedge r \Leftrightarrow p \wedge (q \wedge r) \quad (p \vee q) \vee r \Leftrightarrow p \vee (q \vee r)$$

Idempotence (Ip)

$$p \wedge p \Leftrightarrow p$$

$$p \vee p \Leftrightarrow p$$

Inverse (In)

$$p \wedge \neg p \Leftrightarrow F$$

$$p \vee \neg p \Leftrightarrow T$$

34 / 67

Laws of Logic

Identity (Id)

$$p \wedge T \Leftrightarrow p$$

$$p \vee F \Leftrightarrow p$$

Domination (Do)

$$p \wedge F \Leftrightarrow F$$

$$p \vee T \Leftrightarrow T$$

Distributivity (Di)

$$p \wedge (q \vee r) \Leftrightarrow (p \wedge q) \vee (p \wedge r) \quad p \vee (q \wedge r) \Leftrightarrow (p \vee q) \wedge (p \vee r)$$

Absorption (Ab)

$$p \wedge (p \vee q) \Leftrightarrow p$$

$$p \vee (p \wedge q) \Leftrightarrow p$$

DeMorgan's Laws (DM)

$$\neg(p \wedge q) \Leftrightarrow \neg p \vee \neg q$$

$$\neg(p \vee q) \Leftrightarrow \neg p \wedge \neg q$$

35 / 67

Equivalence Example

Example

$$\begin{aligned} & p \rightarrow q \\ \Leftrightarrow & \neg p \vee q \\ \Leftrightarrow & q \vee \neg p && \text{Co} \\ \Leftrightarrow & \neg\neg q \vee \neg p && \text{DN} \\ \Leftrightarrow & \neg q \rightarrow \neg p \end{aligned}$$

36 / 67

Equivalence Example

Example

$$\begin{aligned}
 & \neg(\neg((p \vee q) \wedge r) \vee \neg q) \\
 \Leftrightarrow & \neg\neg((p \vee q) \wedge r) \wedge \neg\neg q && DM \\
 \Leftrightarrow & ((p \vee q) \wedge r) \wedge q && DN \\
 \Leftrightarrow & (p \vee q) \wedge (r \wedge q) && As \\
 \Leftrightarrow & (p \vee q) \wedge (q \wedge r) && Co \\
 \Leftrightarrow & ((p \vee q) \wedge q) \wedge r && As \\
 \Leftrightarrow & q \wedge r && Ab
 \end{aligned}$$

37 / 67

Duality

Definition

If s contains no logical connectives other than \wedge and \vee , then the **dual** of s , denoted s^d , is the statement obtained from s by replacing each occurrence of \wedge by \vee , \vee by \wedge , T by F , and F by T .

Example (dual proposition)

$$\begin{aligned}
 s &: (p \wedge \neg q) \vee (r \wedge T) \\
 s^d &: (p \vee \neg q) \wedge (r \vee F)
 \end{aligned}$$

38 / 67

Principle of Duality

principle of duality

Let s and t be statements that contain no logical connectives other than \wedge and \vee .

If $s \Leftrightarrow t$ then $s^d \Leftrightarrow t^d$.

39 / 67

Rules of Inference

Definition

if $P \rightarrow Q$ is a tautology, then P **logically implies** Q :
 $P \Rightarrow Q$

40 / 67

Logical Implication Example

Example

$$p \wedge (p \rightarrow q) \Rightarrow q$$

Table: $p \wedge (p \rightarrow q) \rightarrow q$

p	q	$p \rightarrow q$ (A)	$p \wedge A$ (B)	$B \rightarrow q$
T	T	T	T	T
T	F	F	F	T
F	T	T	F	T
F	F	T	F	T

41 / 67

Inference

- establishing the validity of an argument, starting from a set of propositions which are assumed or proven to be true

notation

$$\begin{array}{c}
 p_1 \\
 p_2 \\
 \dots \\
 p_n \\
 \hline
 \therefore q
 \end{array}
 \qquad
 p_1 \wedge p_2 \wedge \dots \wedge p_n \Rightarrow q$$

42 / 67

Basic Rules

Identity (ID)

$$\frac{p}{\therefore p}$$

Contradiction (CTR)

$$\frac{F}{\therefore p}$$

43 / 67

Basic Rules

Implication Introduction (Impl)

$$\frac{p \vdash q}{\therefore \vdash p \rightarrow q}$$

- ▶ if it can be shown that q is true assuming p is true, then $p \rightarrow q$ is true *without assuming p is true*
- ▶ p is a **provisional assumption** (PA)
- ▶ provisional assumptions have to be **discharged** at some point

44 / 67

Basic Rules

AND Introduction (AndI)

$$\frac{\begin{array}{c} p \\ q \end{array}}{\therefore p \wedge q}$$

AND Elimination (AndE)

$$\frac{p \wedge q}{\therefore p}$$

45 / 67

Basic Rules

OR Introduction (OrI)

$$\frac{p}{\therefore p \vee q}$$

OR Elimination (OrE)

$$\frac{\begin{array}{c} p \vee q \\ p \vdash r \\ q \vdash r \end{array}}{\therefore \vdash r}$$

46 / 67

Basic Rules

Modus Ponens (Implication Elimination - ImpE)

$$\frac{\begin{array}{c} p \rightarrow q \\ p \end{array}}{\therefore q}$$

Modus Tollens (MT)

$$\frac{\begin{array}{c} p \rightarrow q \\ \neg q \end{array}}{\therefore \neg p}$$

47 / 67

Modus Tollens

Example

$$\frac{\begin{array}{c} p \rightarrow q \\ \neg q \end{array}}{\therefore \neg p}$$

1. $p \rightarrow q$ A
2. $\neg q \rightarrow \neg p$ 1
3. $\neg q$ A
4. $\neg p$ $ImpE : 2, 3$

48 / 67

Modus Ponens Example

Example

- ▶ If Ali wins the lottery, he will buy a car.
- ▶ Ali has won the lottery.
- ▶ Therefore, Ali will buy a car.

49 / 67

Modus Tollens Example

Example

- ▶ If Ali wins the lottery, he will buy a car.
- ▶ Ali did not buy a car.
- ▶ Therefore, Ali did not win the lottery.

50 / 67

Fallacies

fallacy of affirming the conclusion

$$\frac{p \rightarrow q \quad q}{\therefore p}$$

- ▶ $(p \rightarrow q) \wedge q \rightarrow p$ is not a tautology:
if $p = F, q = T: (F \rightarrow T) \wedge T \rightarrow F$

51 / 67

Example of Affirming the Conclusion

Example

- ▶ If Ali wins the lottery, he will buy a car.
- ▶ Ali has bought a car.
- ▶ Therefore, Ali has won the lottery.

52 / 67

Fallacies

fallacy of denying the hypothesis

$$\frac{p \rightarrow q \quad \neg p}{\therefore \neg q}$$

- ▶ $(p \rightarrow q) \wedge \neg p \rightarrow \neg q$ is not a tautology:
if $p = F, q = T: (F \rightarrow T) \wedge T \rightarrow F$

53 / 67

Example of Denying the Hypothesis

Example

- ▶ If Ali wins the lottery, he will buy a car.
- ▶ Ali has not won the lottery.
- ▶ Therefore, Ali will not buy a car.

54 / 67

Disjunctive Syllogism

Disjunctive Syllogism (DS)

$$\frac{p \vee q \quad \neg p}{\therefore q}$$

1. $p \vee q$ A
2. $\neg p$ A
3. $p \rightarrow F$ 2
- 4a1. p PA
- 4a2. F $ImpE : 3, 4a1$
- 4a. q $CTR : 4a2$
- 4b1. q PA
- 4b. q $ID : 4b1$
5. q $OrE : 1, 4a, 4b$

55 / 67

Disjunctive Syllogism Example

Example

- Ali's wallet is either in his pocket or on his desk.
- Ali's wallet is not in his pocket.
- Therefore, Ali's wallet is on his desk.

56 / 67

Hypothetical Syllogism

Hypothetical Syllogism (HS)

$$\frac{p \rightarrow q \quad q \rightarrow r}{\therefore p \rightarrow r}$$

1. p PA
2. $p \rightarrow q$ A
3. q $ImpE : 2, 1$
4. $q \rightarrow r$ A
5. r $ImpE : 4, 3$
6. $p \rightarrow r$ $Impl : 1, 5$

57 / 67

Hypothetical Syllogism Example

Example (Star Trek)

Spock to Lieutenant Decker:

*It would be a suicide to attack the enemy ship now.
Someone who attempts suicide is not psychologically fit
to command the Enterprise.
Therefore, I am obliged to relieve you from duty.*

58 / 67

Hypothetical Syllogism Example

Example (Star Trek)

- p : Decker attacks the enemy ship.
- q : Decker attempts suicide.
- r : Decker is not psychologically fit to command the Enterprise.
- s : Spock relieves Decker from duty.

59 / 67

Hypothetical Syllogism Example

Example

$$\frac{p \quad p \rightarrow q \quad q \rightarrow r \quad r \rightarrow s}{\therefore s}$$

1. $p \rightarrow q$ A
2. $q \rightarrow r$ A
3. $p \rightarrow r$ $HS : 1, 2$
4. $r \rightarrow s$ A
5. $p \rightarrow s$ $HS : 3, 4$
6. p A
7. s $ImpE : 5, 6$

60 / 67

Inference Examples

Example

$p \rightarrow r$	1. $u \vee \neg x$ <i>A</i>	6. $r \rightarrow s$ <i>A</i>
$r \rightarrow s$	2. $\neg u$ <i>A</i>	7. $\neg r$ <i>MT : 6, 5</i>
$x \vee \neg s$	3. $\neg x$ <i>DS : 1, 2</i>	8. $p \rightarrow r$ <i>A</i>
$u \vee \neg x$	4. $x \vee \neg s$ <i>A</i>	9. $\neg p$ <i>MT : 8, 7</i>
$\neg u$	5. $\neg s$ <i>DS : 4, 3</i>	
$\therefore \neg p$		

61 / 67

Inference Examples

Example

$$\frac{\begin{array}{l} (\neg p \vee \neg q) \rightarrow (r \wedge s) \\ r \rightarrow x \\ \neg x \end{array}}{\therefore p}$$

1. $r \rightarrow x$ <i>A</i>	6. $(\neg p \vee \neg q) \rightarrow (r \wedge s)$ <i>A</i>
2. $\neg x$ <i>A</i>	7. $\neg(\neg p \vee \neg q)$ <i>MT : 6, 5</i>
3. $\neg r$ <i>MT : 1, 2</i>	8. $p \wedge q$ <i>DM : 7</i>
4. $\neg r \vee \neg s$ <i>Orl : 3</i>	9. p <i>AndE : 8</i>
5. $\neg(r \wedge s)$ <i>DM : 4</i>	

62 / 67

Inference Examples

Example

$p \rightarrow (q \vee r)$	1. $q \rightarrow \neg p$ <i>A</i>
$s \rightarrow \neg r$	2. p <i>A</i>
$q \rightarrow \neg p$	3. $\neg q$ <i>MT : 1, 2</i>
p	4. s <i>A</i>
s	5. $s \rightarrow \neg r$ <i>A</i>
$\therefore F$	6. $\neg r$ <i>ImpE : 5, 4</i>
	7. $p \rightarrow (q \vee r)$ <i>A</i>
	8. $q \vee r$ <i>ImpE : 7, 2</i>
	9. q <i>DS : 8, 6</i>
	10. $q \wedge \neg q : F$ <i>AndI : 9, 3</i>

63 / 67

Inference Examples

Example

If there is a chance of rain or her red headband is missing, then Lois will not mow her lawn. Whenever the temperature is over 20°C, there is no chance for rain. Today the temperature is 22°C and Lois is wearing her red headband. Therefore, Lois will mow her lawn.

64 / 67

Inference Examples

Example

- p : There is a chance of rain.
- q : Lois' red headband is lost.
- r : Lois mows her lawn.
- s : The temperature is over 20°C.

65 / 67

Inference Examples

Example

$(p \vee q) \rightarrow \neg r$	1. $s \wedge \neg q$ <i>A</i>
$s \rightarrow \neg p$	2. s <i>AndE : 1</i>
$s \wedge \neg q$	3. $s \rightarrow \neg p$ <i>A</i>
$\therefore r$	4. $\neg p$ <i>ImpE : 3, 2</i>
	5. $\neg q$ <i>AndE : 1</i>
	6. $\neg p \wedge \neg q$ <i>AndI : 4, 5</i>
	7. $\neg(p \vee q)$ <i>DM : 6</i>
	8. $(p \vee q) \rightarrow \neg r$ <i>A</i>
	9. $?$ <i>7, 8</i>

66 / 67

References

Required Text: Grimaldi

- ▶ Chapter 2: Fundamentals of Logic
 - ▶ 2.1. Basic Connectives and Truth Tables
 - ▶ 2.2. Logical Equivalence: The Laws of Logic
 - ▶ 2.3. Logical Implication: Rules of Inference

Supplementary Text: O'Donnell, Hall, Page

- ▶ Chapter 6: Propositional Logic