

# Discrete Mathematics

## Theorem Proving

H. Turgut Uyar   Ayşegül Gençata Yayımlı   Emre Harmancı

2001-2012

1 / 42

## License



©2001-2012 T. Uyar, A. Yayımlı, E. Harmancı

You are free:

- ▶ to Share — to copy, distribute and transmit the work
- ▶ to Remix — to adapt the work

Under the following conditions:

- ▶ Attribution — You must attribute the work in the manner specified by the author or licensor (but not in any way that suggests that they endorse you or your use of the work).
- ▶ Noncommercial — You may not use this work for commercial purposes.
- ▶ Share Alike — If you alter, transform, or build upon this work, you may distribute the resulting work only under the same or similar license to this one.

Legal code (the full license):

<http://creativecommons.org/licenses/by-nc-sa/3.0/>

2 / 42

## Topics

### Basic Techniques

Introduction  
Direct Proof  
Proof by Contradiction  
Equivalence Proofs

### Induction

Introduction  
Strong Induction

3 / 42

## Brute Force Method

- ▶ examining all possible cases one by one

### Theorem

*Every number from the set  $\{2, 4, 6, \dots, 26\}$  can be written as the sum of at most 3 square numbers.*

### Proof.

$2 = 1+1$	$10 = 9+1$	$20 = 16+4$
$4 = 4$	$12 = 4+4+4$	$22 = 9+9+4$
$6 = 4+1+1$	$14 = 9+4+1$	$24 = 16+4+4$
$8 = 4+4$	$16 = 16$	$26 = 25+1$
	$18 = 9+9$	

□

4 / 42

## Basic Rules

### Universal Specification (US)

$\forall x \, p(x) \Rightarrow p(a)$

### Universal Generalization (UG)

$p(a)$  for an arbitrarily chosen  $a \Rightarrow \forall x \, p(x)$

5 / 42

## Universal Specification Example

### Example

*All humans are mortal. Socrates is human.  
Therefore, Socrates is mortal.*

- ▶  $\mathcal{U}$ : all humans
- ▶  $p(x)$ :  $x$  is mortal
- ▶  $\forall x \, p(x)$ : All humans are mortal.
- ▶  $a$ : Socrates,  $a \in \mathcal{U}$ : Socrates is human.
- ▶ therefore,  $p(a)$ : Socrates is mortal.

6 / 42

## Universal Specification Example

### Example

$\forall x [j(x) \vee s(x) \rightarrow \neg p(x)]$	1. $\forall x [j(x) \vee s(x) \rightarrow \neg p(x)]$	$A$
$p(m)$	2. $p(m)$	$A$
$\therefore \neg s(m)$	3. $j(m) \vee s(m) \rightarrow \neg p(m)$	$US : 1$
	4. $\neg(j(m) \vee s(m))$	$MT : 3, 2$
	5. $\neg j(m) \wedge \neg s(m)$	$DM : 4$
	6. $\neg s(m)$	$AndE : 5$

7 / 42

## Universal Generalization Example

### Example

$\forall x [p(x) \rightarrow q(x)]$	1. $\forall x [p(x) \rightarrow q(x)]$	$A$
$\forall x [q(x) \rightarrow r(x)]$	2. $p(c) \rightarrow q(c)$	$US : 1$
$\therefore \forall x [p(x) \rightarrow r(x)]$	3. $\forall x [q(x) \rightarrow r(x)]$	$A$
	4. $q(c) \rightarrow r(c)$	$US : 3$
	5. $p(c) \rightarrow r(c)$	$HS : 2, 4$
	6. $\forall x [p(x) \rightarrow r(x)]$	$UG : 5$

8 / 42

## Vacuous Proof

### vacuous proof

to prove  $P \Rightarrow Q$ , show that  $P$  is false

9 / 42

## Vacuous Proof Example

### Theorem

$\forall S [\emptyset \subseteq S]$

### Proof.

$\emptyset \subseteq S \Leftrightarrow \forall x [x \in \emptyset \rightarrow x \in S]$

$\forall x [x \notin \emptyset]$

□

10 / 42

## Trivial Proof

### trivial proof

to prove  $P \Rightarrow Q$ , show that  $Q$  is true

11 / 42

## Trivial Proof Example

### Theorem

$\forall x \in \mathbb{R} [x \geq 0 \Rightarrow x^2 \geq 0]$

### Proof.

$\forall x \in \mathbb{R} [x^2 \geq 0]$

□

12 / 42

## Direct Proof

### direct proof

to prove  $P \Rightarrow Q$ , show that  $P \vdash Q$

13 / 42

## Direct Proof Example

### Theorem

$$\forall a \in \mathbb{Z} [3|(a-2) \Rightarrow 3|(a^2-1)]$$

### Proof.

$$\begin{aligned} 3|(a-2) &\Rightarrow a-2=3k \\ &\Rightarrow a+1=a-2+3=3k+3=3(k+1) \\ &\Rightarrow a^2-1=(a+1)(a-1)=3(k+1)(a-1) \end{aligned}$$

□

14 / 42

## Indirect Proof

### indirect proof

to prove  $P \Rightarrow Q$ , show that  $\neg Q \vdash \neg P$

15 / 42

## Indirect Proof Example

### Theorem

$$\forall x, y \in \mathbb{N} [x \cdot y > 25 \Rightarrow (x > 5) \vee (y > 5)]$$

### Proof.

- ▶  $\neg Q \Leftrightarrow (0 \leq x \leq 5) \wedge (0 \leq y \leq 5)$
- ▶  $0 = 0 \cdot 0 \leq x \cdot y \leq 5 \cdot 5 = 25$

□

16 / 42

## Indirect Proof Example

### Theorem

$$(\exists k, a, b, k \in \mathbb{N} [ab = 2k]) \Rightarrow (\exists i \in \mathbb{N} [a = 2i]) \vee (\exists j \in \mathbb{N} [b = 2j])$$

### Proof.

$$\blacktriangleright \neg Q \Leftrightarrow (\neg \exists i \in \mathbb{N} [a = 2i]) \wedge (\neg \exists j \in \mathbb{N} [b = 2j])$$

$$\begin{aligned} &\Rightarrow (\exists x \in \mathbb{N} [a = 2x + 1]) \wedge (\exists y \in \mathbb{N} [b = 2y + 1]) \\ &\Rightarrow ab = (2x + 1)(2y + 1) \\ &\Rightarrow ab = 4xy + 2(x + y) + 1 \\ &\Rightarrow \neg(\exists a, b, k \in \mathbb{N} [ab = 2k]) \end{aligned}$$

□

17 / 42

## Proof by Contradiction

### proof by contradiction

to prove  $P$ , show that  $\neg P \vdash Q \wedge \neg Q$

18 / 42

## Proof by Contradiction Example

### Theorem

There is no largest prime number.

### Proof.

- ▶  $\neg P$ : There is a largest prime number.
- ▶  $Q$ : The largest prime number is  $S$ .
- ▶ prime numbers:  $2, 3, 5, 7, 11, \dots, S$
- ▶  $2 \cdot 3 \cdot 5 \cdot 7 \cdot 11 \cdots S + 1$  is not divisible by a prime number between  $2..S$ 
  1. either it is prime itself:  $\neg Q$
  2. or it is divisible by a prime number greater than  $S$ :  $\neg Q$

□

19 / 42

## Proof by Contradiction Example

### Theorem

$\neg \exists a, b \in \mathbb{Z}^+ [\sqrt{2} = \frac{a}{b}]$

### Proof.

- ▶  $\neg P$ :  $\exists a, b \in \mathbb{Z}^+ [\sqrt{2} = \frac{a}{b}]$
- ▶  $Q$ :  $\gcd(a, b) = 1$

$$\Rightarrow 2 = \frac{a^2}{b^2}$$

$$\Rightarrow a^2 = 2b^2$$

$$\Rightarrow \exists i \in \mathbb{Z}^+ [a^2 = 2i]$$

$$\Rightarrow \exists j \in \mathbb{Z}^+ [a = 2j]$$

$$\Rightarrow 4j^2 = 2b^2$$

$$\Rightarrow b^2 = 2j^2$$

$$\Rightarrow \exists k \in \mathbb{Z}^+ [b^2 = 2k]$$

$$\Rightarrow \exists l \in \mathbb{Z}^+ [b = 2l]$$

$$\Rightarrow \gcd(a, b) \geq 2 : \neg Q$$

□

20 / 42

## Equivalence Proofs

- ▶ to prove  $P \Leftrightarrow Q$ , both  $P \Rightarrow Q$  and  $Q \Rightarrow P$  must be proven
- ▶ a method to prove  $P_1 \Leftrightarrow P_2 \Leftrightarrow \dots \Leftrightarrow P_n$ :  
 $P_1 \Rightarrow P_2 \Rightarrow \dots \Rightarrow P_n \Rightarrow P_1$

21 / 42

## Equivalence Proof Example

### Theorem

$a, b, n, q_1, r_1, q_2, r_2 \in \mathbb{Z}^+$

$a = q_1 \cdot n + r_1$

$b = q_2 \cdot n + r_2$

$r_1 = r_2 \Leftrightarrow n|(a - b)$

22 / 42

## Equivalence Proof Example

$$r_1 = r_2 \Rightarrow n|(a - b).$$

$$n|(a - b) \Rightarrow r_1 = r_2.$$

$$\begin{aligned} a - b &= (q_1 \cdot n + r_1) \\ &\quad - (q_2 \cdot n + r_2) \\ &= (q_1 - q_2) \cdot n \\ &\quad + (r_1 - r_2) \end{aligned}$$

$$\begin{aligned} r_1 = r_2 &\Rightarrow r_1 - r_2 = 0 \\ &\Rightarrow a - b = (q_1 - q_2) \cdot n \end{aligned}$$

□

$$\begin{aligned} a - b &= (q_1 \cdot n + r_1) \\ &\quad - (q_2 \cdot n + r_2) \\ &= (q_1 - q_2) \cdot n \\ &\quad + (r_1 - r_2) \end{aligned}$$

$$\begin{aligned} n|(a - b) &\Rightarrow r_1 - r_2 = 0 \\ &\Rightarrow r_1 = r_2 \end{aligned}$$

□

23 / 42

## Equivalence Proof Example

### Theorem

$$\begin{aligned} A &\subseteq B \\ \Leftrightarrow A \cup B &= B \\ \Leftrightarrow A \cap B &= A \\ \Leftrightarrow \overline{B} &\subseteq \overline{A} \end{aligned}$$

24 / 42

## Equivalence Proof Example

$$A \subseteq B \Rightarrow A \cup B = B.$$

$$A \cup B = B \Leftrightarrow A \cup B \subseteq B \wedge B \subseteq A \cup B$$

$$B \subseteq A \cup B$$

$$x \in A \cup B \Rightarrow x \in A \vee x \in B$$

$$A \subseteq B \Rightarrow x \in B$$

$$\Rightarrow A \cup B \subseteq B \quad \square$$

25 / 42

## Equivalence Proof Example

$$A \cup B = B \Rightarrow A \cap B = A.$$

$$A \cap B = A \Leftrightarrow A \cap B \subseteq A \wedge A \subseteq A \cap B$$

$$A \cap B \subseteq A$$

$$y \in A \Rightarrow y \in A \cup B$$

$$A \cup B = B \Rightarrow y \in B$$

$$\Rightarrow y \in A \cap B$$

$$\Rightarrow A \subseteq A \cap B \quad \square$$

26 / 42

## Equivalence Proof Example

$$A \cap B = A \Rightarrow \bar{B} \subseteq \bar{A}.$$

$$z \in \bar{B} \Rightarrow z \notin B$$

$$\Rightarrow z \notin A \cap B$$

$$A \cap B = A \Rightarrow z \notin A$$

$$\Rightarrow z \in \bar{A}$$

$$\Rightarrow \bar{B} \subseteq \bar{A} \quad \square$$

27 / 42

## Equivalence Proof Example

$$\bar{B} \subseteq \bar{A} \Rightarrow A \subseteq B.$$

$$\neg(A \subseteq B) \Rightarrow \exists w [w \in A \wedge w \notin B]$$

$$\Rightarrow \exists w [w \notin \bar{A} \wedge w \in \bar{B}]$$

$$\Rightarrow \neg(\bar{B} \subseteq \bar{A}) \quad \square$$

28 / 42

## Induction

### Definition

$S(n)$ : a predicate defined on  $n \in \mathbb{Z}^+$


$$S(n_0) \wedge (\forall k \geq n_0 [S(k) \Rightarrow S(k+1)]) \Rightarrow \forall n \geq n_0 S(n)$$

►  $S(n_0)$ : base step

►  $\forall k \geq n_0 [S(k) \Rightarrow S(k+1)]$ : induction step

29 / 42

## Induction



30 / 42

## Induction Example

### Theorem

$$\forall n \in \mathbb{Z}^+ [1 + 3 + 5 + \dots + (2n - 1) = n^2]$$

### Proof.

- ▶  $n = 1$ :  $1 = 1^2$
- ▶  $n = k$ : assume  $1 + 3 + 5 + \dots + (2k - 1) = k^2$
- ▶  $n = k + 1$ :

$$\begin{aligned} & 1 + 3 + 5 + \dots + (2k - 1) + (2k + 1) \\ &= k^2 + 2k + 1 \\ &= (k + 1)^2 \end{aligned}$$

□

31 / 42

## Induction Example

### Theorem

$$\forall n \in \mathbb{Z}^+, n \geq 4 [2^n < n!]$$

### Proof.

- ▶  $n = 4$ :  $2^4 = 16 < 24 = 4!$
- ▶  $n = k$ : assume  $2^k < k!$
- ▶  $n = k + 1$ :  
 $2^{k+1} = 2 \cdot 2^k < 2 \cdot k! < (k + 1) \cdot k! = (k + 1)!$

□

32 / 42

## Induction Example

### Theorem

$$\forall n \in \mathbb{Z}^+, n \geq 14 \exists i, j \in \mathbb{N} [n = 3i + 8j]$$

### Proof.

- ▶  $n = 14$ :  $14 = 3 \cdot 2 + 8 \cdot 1$
- ▶  $n = k$ : assume  $k = 3i + 8j$
- ▶  $n = k + 1$ :
  - ▶  $k = 3i + 8j, j > 0 \Rightarrow k + 1 = k - 8 + 3 \cdot 3$   
 $\Rightarrow k + 1 = 3(i + 3) + 8(j - 1)$
  - ▶  $k = 3i + 8j, j = 0, i \geq 5 \Rightarrow k + 1 = k - 5 \cdot 3 + 2 \cdot 8$   
 $\Rightarrow k + 1 = 3(i - 5) + 8(j + 2)$

□

33 / 42

## Strong Induction

### Definition

$$S(n_0) \wedge (\forall k \geq n_0 [(\forall i \leq k S(i)) \Rightarrow S(k + 1)]) \Rightarrow \forall n \geq n_0 S(n)$$

34 / 42

## Strong Induction Example

### Theorem

$$\forall n \in \mathbb{Z}^+, n \geq 2$$

*n can be written as the product of prime numbers*

### Proof.

- ▶  $n = 2$ :  $2 = 2$
- ▶ assume that the theorem is true for  $\forall i \leq k$
- ▶  $n = k + 1$ :
  1. if prime:  $n = n$
  2. if not prime:  $n = u \cdot v$   
 $u < k \wedge v < k \Rightarrow$  both  $u$  and  $v$  can be written as the product of prime numbers

□

35 / 42

## Strong Induction Example

### Theorem

$$\forall n \in \mathbb{Z}^+, n \geq 14 \exists i, j \in \mathbb{N} [n = 3i + 8j]$$

### Proof.

- ▶  $n = 14$ :  $14 = 3 \cdot 2 + 8 \cdot 1$
- ▶  $n = 15$ :  $15 = 3 \cdot 5 + 8 \cdot 0$
- ▶  $n = 16$ :  $16 = 3 \cdot 0 + 8 \cdot 2$
- ▶  $n \leq k$ : assume  $k = 3i + 8j$
- ▶  $n = k + 1$ :  $k + 1 = (k - 2) + 3$

□

36 / 42

## Flawed Induction Examples

### Theorem

$$\forall n \in \mathbb{Z}^+ [1 + 2 + 3 + \dots + n = \frac{n^2 + n + 2}{2}]$$

### invalid base step


- ▶  $n = k$ : assume  $1 + 2 + 3 + \dots + k = \frac{k^2 + k + 2}{2}$
- ▶  $n = k + 1$ :

$$\begin{aligned} & 1 + 2 + 3 + \dots + k + (k + 1) \\ &= \frac{k^2 + k + 2}{2} + k + 1 = \frac{k^2 + k + 2}{2} + \frac{2k + 2}{2} \\ &= \frac{k^2 + 3k + 4}{2} = \frac{(k + 1)^2 + (k + 1) + 2}{2} \end{aligned}$$

- ▶  $n = 1$ :  $1 \neq \frac{1^2 + 1 + 2}{2} = 2$

37 / 42

## Flawed Induction Examples



38 / 42

## Flawed Induction Examples

### Theorem

*All horses are of the same color.*

$A(n)$ : *All horses in sets of  $n$  cardinality are of the same color.*

$$\forall n \in \mathbb{N}^+ A(n)$$

39 / 42


## Flawed Induction Examples

### Flawed induction over $n$

- ▶  $n = 1$ :  $A(1)$   
All horses in sets of 1 horse are of the same color.
- ▶  $n = k$ : assume  $A(k)$  is true  
All horses in sets of  $k$  horses are of the same color.
- ▶  $A(k + 1) = \{a_1, a_2, \dots, a_k\} \cup \{a_2, a_3, \dots, a_{k+1}\}$ 
  - ▶ All horses in set  $\{a_1, a_2, \dots, a_k\}$  are of the same color ( $a_2$ )
  - ▶ All horses in set  $\{a_2, a_3, \dots, a_{k+1}\}$  are of the same color ( $a_2$ )

40 / 42

## Flawed Induction Examples



41 / 42

## References

### Required Text: Grimaldi

- ▶ Chapter 2: Fundamentals of Logic
  - ▶ 2.5. Quantifiers, Definitions, and the Proofs of Theorems
- ▶ Chapter 4: Properties of Integers: Mathematical Induction
  - ▶ 4.1. The Well-Ordering Principle: Mathematical Induction

### Supplementary Text: O'Donnell, Hall, Page

- ▶ Chapter 4: Induction

42 / 42