

Research proposal: Neighborhood environment and obesity

Kenta Okuyama

2019-01-26

Contents

1	Introduction	1
2	Litterature review	3
2.1	Physical environments	3
2.2	Social capital	3
3	Aims and Significance	4
4	Methods	4
4.1	Aim 1: longitudinal changes of neighborhood physical envion- ment and obesity	4
4.2	Aim 2: interaction effect of physial environment and social envi- ronment on obesity	4

1 Introduction

Obesity can lead to serious negative consequences, such as type 2 diabetes and cardiovascular disease, and its prevalence has been increasing worldwide [13, 3]. Environmental and policy interventions are now recognized as an urgent approach to combat the global obesity epidemic [5]. Although numerous research have been done on the influence of environmental features on obesity, there is no consistent evidence [15, 10]. One of the major challenges is that majority of the studies were conducted cross-sectionally, thereby causal relationship has not been elucidated yet. A few recent studies found a significant association between neighborhood physical environment features and obesity among mid-life to older adults [11, 1]. For example, Mason et al. found that both density of physical activity facilities and proximity to fastfood outlets were associated with adiposity in the expected direction among nation-wide adults cohort in UK. Barrientos et al. found that improvement of neighborhood environment in promoting both healthy eating and physical activity was related to BMI reduction among obese persons [1]. This was the first preliminary finding for the association between longitudinal changes of physical environment and BMI. However, despite these recent findings, the association between neighborhood physical environment and obesity are still unclear [14]. This is mainly because complex mechanism for the incidence of obesity due to numerous factors that can affect energy imbalance for diet and physical activity, such as socio-economic status, neighborhood deprivation, and genetic risk factors (Figure 1). Lacking or inappropriate control for any of these factors would lead biased findings for the influence of neighborhood environment on obesity. Those recent studies were either lacking of control for any of the factors, such as neighborhood deprivation, or conducted cross-sectionally.

As another reason why the association between neighborhood environment and obesity is unclear, there are often unmeasured factors that could influence obesity within neighborhood studies. For example, there is a growing body of evidences that not only physical environment, but also social environments might affect on physical activity and obesity through different pathways [2, 19]. Social environments are often measured by social capital which refers to mutual trust, supports, and participation of social activity among neighborhoods [7]. Several studies have examined the association between social capital and obesity, yet evidence for this association has been inconsistent [2]. A few studies have investigated whether neighborhood social capital and physical environment was associated to each other, but findings are inconsistent [9, 6]. Although both physical environment and social environment are both factors in the upstream for the obesity related behaviors, few studies have examined which factor affect on obesity more than the other. From these background, this study consists of two aims.

Aim 1 Examine the longitudinal association between changes in neighborhood physical environment (food and physical activity) and obesity.

Aim 2 Examine the association between neighborhood physical environment and its interaction with social environment on obesity.




Figure 1: Directed acyclic graph for neighborhood environments and obesity

2 Literature review

2.1 Physical environments

Originally, physical environments have been identified as one of the important determinants on human behaviors based on ecological and socio-ecological model which had evolved in the late 1900s [12, 16]. Ecological model specifies levels of influence on behavior, from individual, institutional, community, environmental, and policy factors [12]. Although there were growing recognitions for the potential effect of environment and policy interventions in the population level, few interventions have been applied to the control of chronic diseases as of late 1900s [18]. It is often challenging to implement environment and policy level interventions as it requires substantial investments and collaborations in various sectors (e.g. health, transportation, and urban planning).

Numerous efforts have been made to put it forward by [16] specific to physical activity to stimulate more research in this field in order to accumulate sufficient evidences for environment and policy changes.

Since early 2000s, neighborhood research have been accelerated, especially because technologies made it possible to assess physical environment, i.e. geographic information system (GIS). GIS, which used to be utilized in the field of geographics and urban planning to evaluate geography and structure of cities by numerical data, had been incorporated into health field [4]. Collaboration with urban planning and health fields put emphasis on the need of creating "walkable" neighborhoods by focussing on human-made built environments, which had been already recognized for safety perspective, such as New Urbanism, or Smart Growth America. Numerous research have been conducted worldwide to examine association between physical activity, chronic conditions and physical environments. Several places such as XXX implemented environment intervention based on these research evidences.

2.2 Social capital

Social capital has been defined "features of a social organization, such as trusts, norms, and networks that facilitated cooperation for mutual benefits" [8]. Originally, social capital has been recognized as an important determinant for government performance as well as crime prevention and safety [17]. It has been found to be an determinant for population health later by [7], whose study first identified the association between social capital level and mortality. Since these first studies on social capital, it has been distinguished into "contextual" and "compositional" effects. Compositional effect refers to individual level effect such as having contacts with friends, relatives and family members, or belong to social groups. Contextial effect refers to group/area level effect such as mutual trusts, supports, and norms within communities. These measures are usually collected via survey or constructed as a proxy by using census data.

An important thing to consider is the size of area when measuring social capital. Size of the area could be from minimum of individual to neighborhood, city, state, and countries. When measuring contextual effect of social capital, neighborhood is known to be optimal area size because more rapid diffusion of information occurs within smaller areas compared to larger areas. Numerous studies have examined the effect of individual and neighborhood social capital on health since early 2000.

3 Aims and Significance

Aim 1 Examining the casual relationship between physical environment and obesity would reinforce the evidence that emphasize the needs of environmental and policy interventions.

Aim 2 Examining the effect of physical environment and social environment could infer efficient and cost-effective approach to reduce the risk of obesity.

4 Methods

4.1 Aim 1: longitudinal changes of neighborhood physical environment and obesity


Study sample Nationwide sample of men and women from a national Swedish research database.

Analysis Temporal neighborhood walkability index and proximity to fast food outlets calculated by GIS would be used as primary exposure for obesity. Each exposure would be examined by separate models to predict the effect of each exposure on obesity (Figure 2).

4.2 Aim 2: interaction effect of physical environment and social environment on obesity

Study sample Nationwide sample of men and women from a national Swedish research database.

Analysis Using neighborhood proportion of voting as a proxy for social capital, examine its effect on adults' obesity and its interaction with physical environment.



(a) DAG for physical activity environment and obesity. (b) DAG for food environment and obesity

Figure 2: Two figures in line

References

- [1] Tonatiuh Barrientos-Gutierrez et al. “Neighborhood physical environment and changes in body mass index: results from the Multi-Ethnic Study of Atherosclerosis”. In: *American journal of epidemiology* 186.11 (2017), pp. 1237–1245.
- [2] E Carrillo-Álvarez, I Kawachi, and J Riera-Romaní. “Neighbourhood social capital and obesity: a systematic review of the literature”. In: *Obesity Reviews* 20.1 (2019), pp. 119–141.
- [3] Katherine M Flegal et al. “Prevalence and trends in obesity among US adults, 1999–2000”. In: *Jama* 288.14 (2002), pp. 1723–1727.
- [4] Lawrence D Frank et al. “Linking objectively measured physical activity with objectively measured urban form: findings from SMARTRAQ”. In: *American journal of preventive medicine* 28.2 (2005), pp. 117–125.
- [5] Steven L Gortmaker et al. “Changing the future of obesity: science, policy, and action”. In: *The Lancet* 378.9793 (2011), pp. 838–847.
- [6] Tomoya Hanibuchi et al. “Does walkable mean sociable? Neighborhood determinants of social capital among older adults in Japan”. In: *Health & place* 18.2 (2012), pp. 229–239.
- [7] Ichiro Kawachi et al. “Social capital, income inequality, and mortality.” In: *American journal of public health* 87.9 (1997), pp. 1491–1498.
- [8] APM Krouwel. “Making Democracy Work. Civic Traditions in Modern Italy [Review of: R. Putnam (1994) Making Democracy Work. Civic Traditions in Modern Italy]”. In: *Acta Politica* 30 (1995).

- [9] Kevin M Leyden. “Social capital and the built environment: the importance of walkable neighborhoods”. In: *American journal of public health* 93.9 (2003), pp. 1546–1551.
- [10] Joreintje D Mackenbach et al. “Obesogenic environments: a systematic review of the association between the physical environment and adult weight status, the SPOTLIGHT project”. In: *BMC public health* 14.1 (2014), p. 233.
- [11] Kate E Mason, Neil Pearce, and Steven Cummins. “Associations between fast food and physical activity environments and adiposity in mid-life: cross-sectional, observational evidence from UK Biobank”. In: *The Lancet Public Health* 3.1 (2018), e24–e33.
- [12] Kenneth R McLeroy et al. “An ecological perspective on health promotion programs”. In: *Health education quarterly* 15.4 (1988), pp. 351–377.
- [13] Marie Ng et al. “Global, regional, and national prevalence of overweight and obesity in children and adults during 1980–2013: a systematic analysis for the Global Burden of Disease Study 2013”. In: *The lancet* 384.9945 (2014), pp. 766–781.
- [14] Mark J Nieuwenhuijsen. “Influence of urban and transport planning and the city environment on cardiovascular disease”. In: *Nature Reviews Cardiology* (2018), p. 1.
- [15] Mia A Papas et al. “The built environment and obesity”. In: *Epidemiologic reviews* 29.1 (2007), pp. 129–143.
- [16] JamesF Sallis, Adrian Bauman, and Michael Pratt. “Environmental and policy interventions to promote physical activitya”. In: *American journal of preventive medicine* 15.4 (1998), pp. 379–397.
- [17] Robert J Sampson and W Byron Groves. “Community structure and crime: Testing social-disorganization theory”. In: *American journal of sociology* 94.4 (1989), pp. 774–802.
- [18] Thomas L Schmid, Michael Pratt, and Elizabeth Howze. “Policy as intervention: environmental and policy approaches to the prevention of cardiovascular disease.” In: *American Journal of Public Health* 85.9 (1995), pp. 1207–1211.
- [19] Shakira F Suglia et al. “Why the neighborhood social environment is critical in obesity prevention”. In: *Journal of Urban Health* 93.1 (2016), pp. 206–212.