

Fertilization Schedule

	Early Development	Late Development/Early Refinement				
	All Species	Junipers, Spruce, Fir, Cypress, Cryptomeria	Single-Flush Pines	Double-Flush Pines	Broadleaf Species (Can be Defoliated)	Broadleaf Species (Should not be Defoliated)
Spring	Synthetic or Organic Higher N-value (10 to 20) Applied as new buds swell Foliar feed/Root drench with Root Ruckus (or equivalent micronutrient/humic acid liquid organic)	Organic Higher N-Value (8 to 12) Applied as new buds swell Foliar feed/Root drench with Root Ruckus (or equivalent micronutrient/humic acid liquid organic)	No cake fertilizer applied Foliar feed/Root drench with Root Ruckus (or equivalent micronutrient/humic acid liquid organic)	Organic Higher N-Value (8 to 12) Applied as new buds swell Foliar feed/Root drench with Root Ruckus (or equivalent micronutrient/humic acid liquid organic)	No cake fertilizer applied Foliar feed/Root drench with Root Ruckus (or equivalent micronutrient/humic acid liquid organic)	No cake fertilizer applied Foliar feed/Root drench with Root Ruckus (or equivalent micronutrient/humic acid liquid organic)
Summer	Synthetic or Organic Higher N-value (10 to 20) Applied monthly or bi-monthly as recommended on packaging Foliar feed monthly with liquid kelp	Organic Higher N-Value (8 to 12) Applied monthly or bi-monthly as recommended on packaging Foliar feed monthly with liquid kelp	Organic Higher N-Value (8 to 12) Applied after sheaths drop and new candles harden off Applied monthly or bi-monthly as recommended on packaging Foliar feed monthly with liquid kelp	No cake fertilizer applied after decandling in early to mid Summer Foliar feed monthly with liquid kelp	Organic Higher N-Value (8 to 12) Applied after <i>second</i> flush of growth hardens off Applied monthly or bi-monthly as recommended on packaging Foliar feed monthly with liquid kelp	Organic Higher N-Value (8 to 12) Applied after <i>first</i> flush of growth hardens off Applied monthly or bi-monthly as recommended on packaging Foliar feed monthly with liquid kelp
Fall	Synthetic or Organic Higher N-value (10 to 20) Last application one month before average start of dormancy Last foliar feed one month before average start of dormancy	Organic Higher N-Value (8 to 12) Last application one month before average start of dormancy Last foliar feed one month before average start of dormancy	Organic Higher N-Value (8 to 12) Last application one month before average start of dormancy Last foliar feed one month before average start of dormancy	Organic Higher N-Value (8 to 12) Applied after <i>second</i> flush of growth hardens off Last application one month before average start of dormancy Last foliar feed one month before average start of dormancy	Organic Higher N-Value (8 to 12) Last application one month before average start of dormancy Last foliar feed one month before average start of dormancy	Organic Higher N-Value (8 to 12) Last application one month before average start of dormancy Last foliar feed one month before average start of dormancy
Winter	No fertilizer applied	No fertilizer applied	No fertilizer applied	No fertilizer applied	No fertilizer applied	No fertilizer applied

	Refinement				
	Junipers, Spruce, Fir, Cypress, Cryptomeria	Single-Flush Pines	Double-Flush Pines	Broadleaf Species (Can be Defoliated)	Broadleaf Species (Should not be Defoliated)
Spring	Organic Milder N-Value (5 to 9) Applied as new buds swell Foliar feed/Root drench with Root Ruckus (or equivalent micronutrient/humic acid liquid organic)	No cake fertilizer applied Foliar feed/Root drench with Root Ruckus (or equivalent micronutrient/humic acid liquid organic)	Organic Higher N-Value (8 to 12) Applied as new buds swell Foliar feed/Root drench with Root Ruckus (or equivalent micronutrient/humic acid liquid organic)	No cake fertilizer applied Foliar feed/Root drench with Root Ruckus (or equivalent micronutrient/humic acid liquid organic)	No cake fertilizer applied Foliar feed/Root drench with Root Ruckus (or equivalent micronutrient/humic acid liquid organic)
Summer	Organic Milder N-Value (5 to 9) Applied monthly or bi-monthly as recommended on packaging Foliar feed monthly with liquid kelp	Organic Milder N-Value (5 to 9) Applied after sheaths drop and new candles harden off Applied monthly or bi-monthly as recommended on packaging Foliar feed monthly with liquid kelp	No cake fertilizer applied after decandling in early to mid Summer Foliar feed monthly with liquid kelp	Organic Milder N-Value (5 to 9) Applied after second flush of growth hardens off Applied monthly or bi-monthly as recommended on packaging Foliar feed monthly with liquid kelp	Organic Milder N-Value (5 to 9) Applied after first flush of growth hardens off Applied monthly or bi-monthly as recommended on packaging Foliar feed monthly with liquid kelp
Fall	Organic Milder N-Value (5 to 9) Last application one month before average start of dormancy Last foliar feed one month before average start of dormancy	Organic Milder N-Value (5 to 9) Last application one month before average start of dormancy Last foliar feed one month before average start of dormancy	Organic Milder N-Value (5 to 9) Applied after second flush of growth hardens off Last application one month before average start of dormancy Last foliar feed one month before average start of dormancy	Organic Milder N-Value (5 to 9) Last application one month before average start of dormancy Last foliar feed one month before average start of dormancy	Organic Milder N-Value (5 to 9) Last application one month before average start of dormancy Last foliar feed one month before average start of dormancy
Winter	No fertilizer applied	No fertilizer applied	No fertilizer applied	No fertilizer applied	No fertilizer applied