

1 Introductory notation

1.1 Markov and Lagrange values

Symbol \mathcal{M} denotes double-infinite sequences from $\mathbb{N}^{\mathbb{Z}}$:

$$\mathcal{M} = \dots a_{-2} a_{-1} a_0 a_1 a_2 \dots$$

I will use $\lambda(\mathcal{M})$, $\mu(\mathcal{M})$ and $f(\mathcal{M})$ for Lagrange, Markov values and height function. Symbols γ and δ denote the lhs and rhs of sequence \mathcal{M} :

$$\begin{aligned}\gamma(\mathcal{M}) &= [0; a_{-1}, a_{-2}, \dots], \\ \delta(\mathcal{M}) &= [0; a_1, a_2, \dots], \\ f(\mathcal{M}) &= a_0 + \gamma(\mathcal{M}) + \delta(\mathcal{M}).\end{aligned}$$

At last, symbols M and L denote the Markov and Lagrange spectra.

1.2 Centered sequence

Definition. A sequence \mathcal{M} is called **centered**, if

$$\mu(\mathcal{M}) = f(\mathcal{M}). \tag{1}$$

Proposition. *Markov spectrum can be defined with only centered sequences:*

$$\{\mu(\mathcal{M}) \mid \mathcal{M} \in \mathbb{N}^{\mathbb{Z}}\} = M = \{\mu(\mathcal{M}) \mid \mathcal{M} \text{ is centered}\}.$$

1.3 Rectangle

Designation. Denote by

$$\{a_{i_1}, a_{i_1+1}, \dots, a_{i_2}\} \quad (i_1 \leq 0 \leq i_2)$$

the set of double-infinite sequences \mathcal{M} with fixed terms $a_{i_1}, a_{i_1+1}, \dots, a_{i_2}$ on the corresponding positions.

Terms a_s for $s < i_1$ and $s > i_2$ are arbitrary integers, chosen such that \mathcal{M} is centered and, maybe, satisfies some conditions.

Segments Δ_1 , Δ_2 and Δ are defined by the following equations:

$$\begin{aligned}\Delta_1 &= [\Delta'_1; \Delta''_1] = [\min \gamma(\mathcal{M}); \max \gamma(\mathcal{M})], \\ \Delta_2 &= [\Delta'_2; \Delta''_2] = [\min \delta(\mathcal{M}); \max \delta(\mathcal{M})], \\ \Delta &= [\Delta'; \Delta''] = a_0 + \Delta_1 + \Delta_2,\end{aligned} \tag{2}$$

where \mathcal{M} belongs to the set.

Note that we will use ' for the lower bound, and '' for the upper bound.

Definition. **Rectangle** is the segment Δ with the set of sequences, defining it.

1.4 Horizontal rectangle

Definition. Call a rectangle Δ **horizontal**, if

$$|\Delta_1| \geq |\Delta_2|. \quad (3)$$

In (3) we allow terms a_s for $s < i_1$ and $s > i_2$ to be integers $\{1, 2, 3\}$, regardless of the requirement that sequences $\mathcal{M} \in \Delta$ are centered.

In other words, Δ is horizontal, if and only if

$$|[0; a_{-1}, \dots, a_{i_1}, \overline{3}, \overline{1}] - [0; a_{-1}, \dots, a_{i_1}, \overline{1}, \overline{3}]| \geq |[0; a_1, \dots, a_{i_2}, \overline{3}, \overline{1}] - [0; a_1, \dots, a_{i_2}, \overline{1}, \overline{3}]|.$$

Clearly, we can always obtain a horizontal rectangle out of the vertical one, as we can reindex the sequence in the opposite direction.

1.5 Subrectangle

Consider a rectangle Δ , set by the sequence center

$$\{a_{i_1}, a_{i_1+1}, \dots, a_{i_2}\}.$$

We will use a **shorter notation** for subrectangles, produced by setting integers a_i for $i < i_1$ or $i > i_2$:

$$\{b_\ell \dots b_1, c_1 \dots c_r\} := \{b_\ell \dots b_1 a_{i_1} \dots a_{i_2} c_1 \dots c_r\}.$$

For example:

$$\{213, 3\} := \{213 a_{i_1} \dots a_{i_2} 3\}, \quad (\text{ex.1})$$

$$\{2, 0\} := \{2 a_{i_1} \dots a_{i_2}\}. \quad (\text{ex.2})$$

We will also shorter the notation (2): lhs and rhs are $\Delta_1(312)$ and $\Delta_2(3)$ for subrectangle (ex.1) and $\Delta_1(2)$ and a_2 for (ex.2).

1.6 Geometrical interpretation

Consider the mapping

$$\begin{aligned} \tilde{h} : \mathbb{N}^{\mathbb{Z}} &\rightarrow \mathbb{R}^2, \\ \tilde{h}(\mathcal{M}) &= (\gamma(\mathcal{M}); \delta(\mathcal{M})). \end{aligned}$$

In these terms, the Markov spectrum M is the projection of some subset $\mathcal{S} \subset C_4 \times C_4$ onto the diagonal.

Then **rectangle** Δ is indeed a rectangle $\Delta_1 \times \Delta_2$ and **subrectangles** are its subrectangles.

We will consider a family of rectangles whose projections cover the beginning of Hall's Ray.

Then we will present the algorithm to split rectangle into subrectangles so that their projections cover the projection of initial rectangle.

When we say that rectangles intersect, we, however, mean that their projections intersect.

The more «squarish» the rectangle, the easier the step.

That's why we will bound the aspect ratio of rectangles (see **good** rectangle).

Formulas to evaluate side lengths and aspect ratio are given in the section 2.

2 Calculations

2.1 Length of Δ_1 or Δ_2

Let's fix some terms of continued fraction $[0; q_1, q_2, q_3, \dots, q_n]$.
We will often need to measure difference

$$[0; q_1, q_2, q_3, \dots, q_n, \frac{1}{\Theta_R}] - [0; q_1, q_2, q_3, \dots, q_n, \frac{1}{\Theta_L}],$$

where Θ 's are some continuations of the continued fraction.

They generally look like $\Theta = [0; 12\overline{13}]$ or something¹. We will set Θ 's explicitly.

For the general proof, Θ 's will be taken from table 5.

Designation. For given continuation Θ_i denote by ε_i the resulting continued fraction:

$$\varepsilon_i = [0; q_1, q_2, \dots, q_n, \frac{1}{\Theta_i}]. \quad (3.4)$$

Then the following equality takes place:

$$|\varepsilon_i - \varepsilon_j| = \frac{|\Theta_i - \Theta_j|}{Q_n^2 (1 + pQ_i) (1 + pQ_j)}, \quad (3.5)$$

where

$$p = \frac{Q_{n-1}}{Q_n}.$$

2.2 Rectangle aspect ratio

Consider some fixed center of rectangle $\{a_{i_1} \dots a_{i_2}\}$.

We will often extend it from the left (right) using some continuations $\Theta_{\gamma_1}, \Theta_{\gamma_2} (\Theta_{\delta_1}, \Theta_{\delta_2})$.

These values produce $\gamma_1, \gamma_2 (\delta_1, \delta_2)$ using (3.4).

In other words, finite continued fractions $[0; a_{-1}, a_{-2}, \dots, a_{i_1}] = \frac{P_{i_1}}{Q_{i_1}} \left([0; a_1, a_2, \dots, a_{i_2}] = \frac{P_{i_2}}{Q_{i_2}} \right)$

are convergents for $\gamma_1, \gamma_2 (\delta_1, \delta_2)$.

Then

$$\left| \frac{\gamma_1 - \gamma_2}{\delta_1 - \delta_2} \right| = \frac{1}{q} \left| \frac{\Theta_{\gamma_1} - \Theta_{\gamma_2}}{\Theta_{\delta_1} - \Theta_{\delta_2}} \right| \frac{1 + p'\Theta_{\delta_1}}{1 + p\Theta_{\gamma_1}} \frac{1 + p'\Theta_{\delta_2}}{1 + p\Theta_{\gamma_2}} \approx \frac{1}{q} \left| \frac{\Theta_{\gamma_1} - \Theta_{\gamma_2}}{\Theta_{\delta_1} - \Theta_{\delta_2}} \right|, \quad (4)$$

where

$$p = \frac{Q_{i_1+1}}{Q_{i_1}}, \quad p' = \frac{Q_{i_2-1}}{Q_{i_2}}, \quad q = \frac{Q_{i_1}^2}{Q_{i_2}^2}.$$

¹Here, as always in this book, \overline{abc}^k means k -times repetition of abc , and \overline{abc} means infinite repetition.

3 Rectangle boundaries

We will set boundaries for rectangles in a different way.

To distinguish rectangles, we will introduce the notion of left- or right-shortened rectangles.

It is given in subsections 3.1-3.3.

Subsections 3.4-3.5 provide the rules for boundaries $\Delta'_1, \Delta''_1, \Delta'_2, \Delta''_2$.

3.1 Resection

Definition. Call **resection** of a segment $A = [a; b]$ a process of removing subsegment $A_{12} = [a_1; b_1]$, leaving two segments $A_1 \sqcup A_2 = [a; a_1] \sqcup [b_1; b]$.

Definition. Call subsegment $A_{12} \subset A$ **normal**, if it is thicker than the two remaining subsegments:

$$|A_{12}| \leq \min \{|A_1|, |A_2|\} \quad (5)$$

We call a resection **normal** if the resected subsegment is normal.

Proposition. For any normal resection, having

$$A + A = (A_1 \sqcup A_2) + (A_1 \sqcup A_2). \quad (6)$$

3.2 Shortened rectangle

Consider a horizontal rectangle Δ .

3.2.1 Case $i_1 = i_2 = 0 \pmod{2}$

If both i_1 and i_2 are even, then left- and right-shortened rectangles are defined as follows.

Definition. Rectangle Δ is called left-shortened, if for subrectangle $\{3, 3\}$ the following condition takes place:

$$|\Delta_1(3)| \leq 1.4 \cdot |\Delta_2(3)|. \quad (7)$$

Here, as in (3), we allow terms a_s for $s < i_1$ and $s > i_2$ to be integers $\{1, 2, 3\}$, so (7) can be rewritten as

$$|[0; a_{-1}, \dots, a_{i_1}, 3, \overline{3}, \overline{1}] - [0; a_{-1}, \dots, a_{i_1}, \overline{3}, \overline{1}]| \leq |[0; a_1, \dots, a_{i_2}, 3, \overline{3}, \overline{1}] - [0; a_1, \dots, a_{i_2}, \overline{3}, \overline{1}]|.$$

For further convenience, we also introduce the opposite to (7) condition:

Definition. Rectangle Δ is called left-normal, if

$$|\Delta_1(3)| > 1.4 \cdot |\Delta_2(3)|. \quad (8)$$

Now we introduce the notion of right-shortened rectangle:

Definition. Rectangle Δ is called right-shortened, if for subrectangle $\{31, 13\}$ the following condition takes place:

$$|\Delta_1(13)| \leq 1.4 \cdot |\Delta_2(13)|. \quad (9)$$

As in 3.2.1, we allow terms a_s for $s < i_1$ and $s > i_2$ to be integers $\{1, 2, 3\}$, so (9) can be rewritten as

$$\left| [0; a_{-1}, \dots, a_{i_1}, 1, 3, \overline{3, 1}] - [0; a_{-1}, \dots, a_{i_1}, \overline{1, 3}] \right| \leq \left| [0; a_1, \dots, a_{i_2}, 1, 3, \overline{3, 1}] - [0; a_1, \dots, a_{i_2}, \overline{1, 3}] \right|.$$

We also introduce the opposite condition:

Definition. Rectangle Δ is called right-normal, if

$$|\Delta_1(13)| > 1.4 \cdot |\Delta_2(13)|. \quad (10)$$

3.2.2 Other cases

If $i_1 \equiv i_2 \pmod{2}$ and i_1 is odd, then the conditions for left- and right-shortened rectangles are swapped.

In case $i_1 \not\equiv i_2 \pmod{2}$, the notions of left- and right-shortened rectangles are inferred from subrectangles $\{1, \}$, $\{2, \}$, and $\{3, \}$.

We will refer to the relevant conditions from 3.5

3.3 Explanation of shortened

Notion left-shortened is used to set the lower bound Δ' for rectangle Δ .

Notion right-shortened is used to set the upper bound Δ'' .

For example, when interested in Δ' when i_1 is even and i_2 is odd, one has to consider larger a_{i_1-1} and smaller a_{i_2+1} . As the rectangle is horizontal, we will add terms to the left first. So we will check the condition (9) for $\{2, \}$ or $\{3, \}$.

In other words, check whether $\{2, \}$ or $\{3, \}$ is left-shortened or left-normal (recall that $i_1 - 1$ and i_2 are odd, so left-shortenence is defined by (9)).

Which subrectangle ($\{2, \}$ or $\{3, \}$) to take, however, depends on terms a_{i_1+1} and a_{i_1} (see 3.4).

3.4 How we ensure centeredness

Consider integers q_1, q_2, \dots, q_n . Let $\{\delta_n\}$ be the set of fractions $\delta_n = [0; q_1, q_2, \dots, q_n, \dots]$ with n fixed terms. We will suppose that n is even (for odd n the bounds are swapped).

At first, determine the smallest of fractions δ_n .

We will consider 2 cases: S (Shortened) and N (Normal):

$$S. \quad q_{n-1} = 3, q_n = 1. \quad (11a)$$

$$N. \quad \text{Otherwise.} \quad (11b)$$

The lower bound δ'_n for segment, containing δ_n , is defined by:

$$\begin{aligned} S. \quad \delta'_n &= [0; q_1, \dots, q_n, 213\overline{12}]. \\ N. \quad \delta'_n &= [0; q_1, \dots, q_n, 3\overline{12}]. \end{aligned}$$

To set the upper bound δ''_n , largest of δ_n , consider 2 other cases:

$$S. \quad q_n = 3. \quad (12a)$$

$$N. \quad q_n \neq 3. \quad (12b)$$

Then

$$\begin{aligned} S. \quad & \delta_n'' = [0; q_1, \dots, q_n, 1213\overline{12}]. \\ N. \quad & \delta_n'' = [0; q_1, \dots, q_n, 13\overline{12}]. \end{aligned}$$

These bounds will allow us to construct sequences \mathcal{M} , for which combination (31313) is forbidden and, therefore, the following condition takes place:

$$f_i(\mathcal{M}) \leq \lambda(\overline{31312}) \approx 4,5241, \quad i \neq 0,$$

which will ensure (1).

3.5 Formal

Let's now turn to concrete definitions and bounds. Remind that we are looking at the horizontal rectangle Δ .

Suppose that $i_1 = i_2 \pmod{2}$, i_1 is even.

3.5.1 Bounds for Δ'

- I. Suppose both $\{\gamma_0(\mathcal{M})\}$ and $\{\delta_0(\mathcal{M})\}$ satisfy (11b) and the rectangle Δ is left-normal, that is, satisfies (8). We will denote such situation by $N - N - N$ (segment Δ_1 is left-normal, Δ_2 is left-normal, rectangle Δ is normal).

In this case define Δ' by equation

$$\Delta' = f(\overline{21}3a_{i_1} \dots a_{i_2} \overline{312}). \quad (13)$$

- IIa. Sets $\{\gamma_0(\mathcal{M})\}$ and $\{\delta_0(\mathcal{M})\}$ meet condition (11b) and Δ is left-shortened, that is, meets condition (7). This is case $N - N - S$ (segments Δ_1 and Δ_2 are left-normal, rectangle Δ is left-shortened).

Then

$$\Delta' = f(\overline{21}3a_{i_1} \dots a_{i_2} 213\overline{12}). \quad (14)$$

- IIb. Set $\{\gamma_0(\mathcal{M})\}$ meets (11b), $\{\delta_0(\mathcal{M})\}$ meets (11a). No matter, what takes place, (??) of (??). It is case $N - S$ (segment Δ_1 is left-normal, segment Δ_2 left-shortened). Bound Δ_1 is defined by (14).

- III. Set $\{\gamma_0(\mathcal{M})\}$ meets (??), $\{\delta_0(\mathcal{M})\}$ doesn't. In this $(Y - H)$ case Δ' is defined by the following:

$$\Delta' = f(\overline{21}312a_{i_1} \dots a_{i_2} \overline{312}). \quad (11.11)$$

- IV. Both $\{\gamma_0(\mathcal{M})\}$ and $\{\delta_0(\mathcal{M})\}$ meet (??). In this $(Y - Y)$ case Δ' is defined by

$$\Delta' = f(\overline{21}312a_{i_1} \dots a_{i_2} 213\overline{12}). \quad (11.12)$$

Figure (1) illustrates the bounds. On the picture:

$$\Delta'_1 = \gamma(\Theta_i), \quad i = 3, 30, \quad \Delta'_2 = \delta(\Theta_i), \quad i = 3, 30, \quad \Delta' = \Delta'_1 + \Delta'_2.$$

Hatched areas correspond to values of a_{i_1-1} or a_{i_2+1} (equal 3), which can not appear in concrete case.

3.5.2 Bounds for Δ''

Now we will provide formulas for Δ'' :

$$\Delta'' = f(\overline{21}31a_{i_1}...a_{i_2}13\overline{12}) \quad (11.13)$$

$$\Delta'' = f(\overline{21}31a_{i_1}...a_{i_2}1213\overline{12}) \quad (11.14)$$

$$\Delta'' = f(\overline{21}3121a_{i_1}...a_{i_2}13\overline{12}) \quad (11.15)$$

$$\Delta'' = f(\overline{21}3121a_{i_1}...a_{i_2}1213\overline{12}) \quad (11.16)$$

Figure (2) regulates the choice of the formulas.

On the figure (1): $\Delta''_1 = \gamma(\Theta_i)$, $i = 90, 94$, $\Delta''_2 = \delta(\Theta_i)$, $i = 90, 94$, $\Delta'' = \Delta''_1 + \Delta''_2$. Hatched areas correspond to restricted value 3 of variables a_{i_1-2} or a_{i_2+2} .

3.5.3 Case $i_1 \not\equiv i_2 \pmod{2}$

Now take case $i_1 \not\equiv i_2 \pmod{2}$, i_1 is even. We will use rules from figure (3) to choose one of 4 formulas for Δ' .

$$I \quad \Delta' = f(\overline{21}3a_{i_1}...a_{i_2}13\overline{12}) \quad (11.17)$$

$$II \quad \Delta' = f(\overline{21}3a_{i_1}...a_{i_2}1213\overline{12}) \quad (11.18)$$

$$III \quad \Delta' = f(\overline{21}312a_{i_1}...a_{i_2}13\overline{12}) \quad (11.19)$$

$$IV \quad \Delta' = f(\overline{21}312a_{i_1}...a_{i_2}1213\overline{12}) \quad (11.20)$$

To determine Δ'' we will use rectangle

$$\{1, 0\}.$$

For this rectangle have $i_1 - 1 \equiv i_2 \pmod{2}$, so we can use all the previous formulas to determine Δ'' .




Figure 1: Bounds Δ' (left) and Δ'' (right).

N	Δ'_i	Δ''_i	Δ''
I	(5)	(5)	(5.I5) H-H-H (I3)
IIa	(5)	(5)	(5.I3) H-H-y (I4)
IIb	(5)	(4)	H-y (I4)
III	(4)	(5)	y-H (I5)
IV	(4)	(4)	y-y (I6)

Figure 2: Rules for choose of Δ'' in case $i_1 = i_2 \pmod{2}$.

N	Δ'_i	Δ'_i	Δ'
I	(I) не имеет места	(5)	H-H (I7)
II	(I) не имеет места	(4)	H-y (I8)
III	(I)	(5)	y-H (I9)
IV	(I)	(4)	y-y (20)

Figure 3: Rules for choose of Δ' in case $i_1 \neq i_2 \pmod{2}$, i_1 is even.

4 Good rectangle

4.1 Definition

Consider a horizontal rectangle $\Delta = \{a_{i_1} \dots a_{i_2}\}$.

Definition. Rectangle Δ is called **good**, if subrectangles $\{2, 0\}$ and $\{1, 0\}$ intersect.




Figure 4: Good rectangles if i_1 is odd (upper) and even (lower).

For example, in case i_1 is even, goodness is equivalent to

$$\{2, 0\}'' \geq \{1, 0\}' \quad (15)$$

Bounds of rectangles are determined by the rules from section 3.

If rectangle is not good (for example, (15) doesn't take place), then

$$(\{2, 0\}''; \{1, 0\}') \not\subset \Delta.$$

Clearly, if the rectangle is not good, then one can not split it into smaller rectangles. That's why we will only consider good rectangles during the proof.

4.2 Sufficient conditions of goodness

4.2.1 Results

A horizontal rectangle Δ is good, if

$$\frac{\Delta_1}{\Delta_2} < \begin{cases} 3.8, & i_1 \equiv i_2 \pmod{2}, \\ 3.43, & i_1 \not\equiv i_2 \pmod{2}. \end{cases} \quad (16)$$

4.2.2 Universal 2, 9 bound

Now let's introduce some sufficient conditions for rectangle to be good.

Remind that we suppose $|\Delta_1| \geq |\Delta_2|$, which means

$$q \frac{1+p\Theta_1}{1+p'\Theta_1} \frac{1+p\Theta_{95}}{1+p'\Theta_{95}} \leq 1.$$

Let's introduce the following designations:

$$\gamma = [0; a_{-1}, \dots, a_{i_1}, \frac{1}{\Theta_\gamma}],$$

$$\delta = [0; a_1, \dots, a_{i_2}, \frac{1}{\Theta_\delta}],$$

where Θ 's are some Θ 's from Figure (5), specified in each case separately.

If rectangle is good, then (12.1) should take place:

$$\gamma' - \gamma'' < |\delta' - \delta''|, \quad (12.1)$$

where $\Theta_{\gamma'} = \Theta_{66}$, $\Theta_{\gamma''} = \Theta_{63}$, $\Theta_{\delta'} = \Theta_{90}$, $\Theta_{\delta''} = \Theta_{30}$.

As I understand, these thetas are supposed to be $\gamma' = \gamma'_{\{1,0\}}$, $\gamma'' = \gamma''_{\{2,0\}}$, and (12.1) is just paraphrasing of (??) in the «worst» case. As usual, Freiman gives no comments.

Inequality (12.1) transforms into

$$0,313 < q \frac{1+p\Theta_{63}}{1+p'\Theta_{30}} \frac{1+p\Theta_{66}}{1+p'\Theta_{90}}. \quad (12.2)$$

Suppose

$$\frac{\Delta_1}{\Delta_2} < 2,9, \quad (12.3)$$

which is equivalent to

$$\frac{1}{q} \frac{1+p'\Theta_1}{1+p\Theta_1} \frac{1+p'\Theta_{95}}{1+p\Theta_{95}} < 2,9. \quad (12.4)$$

Then (12.2) takes place. Indeed, it is so, if

$$\frac{1}{2,9} \frac{1+p'\Theta_1}{1+p\Theta_1} \frac{1+p'\Theta_{95}}{1+p\Theta_{95}} > 0,313 \frac{1+p'\Theta_{30}}{1+p\Theta_{63}} \frac{1+p'\Theta_{90}}{1+p\Theta_{66}}.$$

or, equivalent,

$$1,1 > \frac{1+p\Theta_1}{1+p\Theta_{63}} \frac{1+p\Theta_{95}}{1+p\Theta_{66}} \frac{1+p'\Theta_{30}}{1+p'\Theta_1} \frac{1+p'\Theta_{90}}{1+p'\Theta_{95}},$$

Which is easily checked. *Ha-ha.*


Condition (12.3) is sufficient for rectangle to be good, no matter, are i_1 and i_2 equivalent (mod 2) or not, take conditions (??) and (??) place or not.

$\Theta_1 = [0; 3\bar{1}] = 0,263762$	$\Theta_{35} = [0; 2\bar{1}2\bar{1}3\bar{1}\bar{2}] = 0,36549$	$\Theta_{67} = [0; 1\bar{1}33\bar{1}\bar{2}] = 0,56635$
$\Theta_2 = [0; 3\bar{1}2\bar{1}\bar{3}] = 0,267649$	$\Theta_{36} = [0; 2\bar{1}] = 0,36602$	$\Theta_{68} = [0; 1\bar{1}33\bar{1}] = 0,566423$
$\Theta_3 = [0; 3\bar{1}\bar{2}] = 0,26794$	$\Theta_{37} = [0; 2\bar{1}2\bar{1}\bar{1}\bar{3}] = 0,36779$	$\Theta_{69} = [0; 1\bar{1}2\bar{1}\bar{3}] = 0,57600$
$\Theta_4 = [0; 3\bar{1}233\bar{1}] = 0,270448$	$\Theta_{38} = [0; 2\bar{1}23\bar{1}\bar{2}] = 0,37119$	$\Theta_{70} = [0; 1\bar{1}2\bar{1}3\bar{1}\bar{2}] = 0,57602$
$\Theta_5 = [0; 3\bar{1}23\bar{1}\bar{2}] = 0,270710$	$\Theta_{39} = [0; 2\bar{1}23\bar{1}] = 0,371249$	$\Theta_{71} = [0; 1\bar{1}\bar{2}] = 0,57735$
$\Theta_6 = [0; 3\bar{1}23\bar{1}] = 0,270738$	$\Theta_{40} = [0; 2\bar{1}\bar{1}\bar{3}] = 0,378537$	$\Theta_{72} = [0; 1\bar{1}23\bar{1}\bar{2}] = 0,59032$
$\Theta_7 = [0; 3\bar{1}\bar{1}\bar{1}\bar{3}] = 0,27459$	$\Theta_{41} = [0; 2\bar{1}\bar{1}\bar{1}3\bar{1}\bar{2}] = 0,37858$	$\Theta_{73} = [0; 1\bar{1}\bar{1}\bar{1}\bar{3}] = 0,609108$
$\Theta_8 = [0; 3\bar{1}\bar{1}\bar{1}3\bar{1}\bar{2}] = 0,27462$	$\Theta_{42} = [0; 2\bar{1}\bar{1}\bar{1}33\bar{1}] = 0,379018$	$\Theta_{74} = [0; 1\bar{1}\bar{1}\bar{1}3\bar{1}\bar{2}] = 0,60923$
$\Theta_9 = [0; 3\bar{1}\bar{1}\bar{1}33\bar{1}] = 0,274847$	$\Theta_{43} = [0; 2\bar{1}\bar{1}\bar{2}] = 0,37963$	$\Theta_{75} = [0; 1\bar{1}\bar{1}\bar{1}\bar{1}33\bar{1}] = 0,620981$
$\Theta_{10} = [0; 3\bar{1}\bar{1}\bar{2}] = 0,27517$	$\Theta_{44} = [0; 2\bar{1}\bar{1}\bar{1}\bar{1}33\bar{1}] = 0,383091$	$\Theta_{76} = [0; 1\bar{1}\bar{1}\bar{1}\bar{1}3\bar{1}\bar{2}] = 0,62141$
$\Theta_{11} = [0; 3\bar{1}\bar{1}\bar{2}] = 0,27954$	$\Theta_{45} = [0; 2\bar{1}\bar{1}\bar{1}\bar{1}3\bar{1}\bar{2}] = 0,38326$	$\Theta_{77} = [0; 1\bar{1}\bar{1}\bar{1}\bar{1}\bar{3}] = 0,621460$
$\Theta_{12} = [0; 3\bar{1}\bar{1}33\bar{1}] = 0,280392$	$\Theta_{46} = [0; 2\bar{1}\bar{1}\bar{1}\bar{3}] = 0,383272$	$\Theta_{78} = [0; 1\bar{1}\bar{1}23\bar{1}\bar{2}] = 0,62881$
$\Theta_{13} = [0; 3\bar{1}\bar{1}3\bar{1}\bar{2}] = 0,28097$	$\Theta_{47} = [0; 2\bar{1}\bar{1}23\bar{1}\bar{2}] = 0,38605$	$\Theta_{79} = [0; 1\bar{1}\bar{2}] = 0,63400$
$\Theta_{14} = [0; 3\bar{1}\bar{1}\bar{3}] = 0,28105$	$\Theta_{48} = [0; 2\bar{1}\bar{1}23\bar{1}] = 0,386033$	$\Theta_{80} = [0; 1\bar{1}\bar{1}33\bar{1}] = 0,63839$
$\Theta_{15} = [0; 323\bar{1}] = 0,290550$	$\Theta_{49} = [0; 2\bar{1}\bar{1}233\bar{1}] = 0,386237$	$\Theta_{81} = [0; 1\bar{1}\bar{1}3\bar{1}\bar{2}] = 0,63910$
$\Theta_{16} = [0; 323\bar{1}\bar{2}] = 0,29062$	$\Theta_{50} = [0; 2\bar{1}\bar{1}22\bar{1}] = 0,38651$	$\Theta_{82} = [0; 1\bar{1}\bar{1}\bar{3}] = 0,641742$
$\Theta_{17} = [0; 3233\bar{1}] = 0,291242$	$\Theta_{51} = [0; 2\bar{1}\bar{1}\bar{2}] = 0,38800$	$\Theta_{83} = [0; 123\bar{1}\bar{2}] = 0,69399$
$\Theta_{18} = [0; 322\bar{1}] = 0,29216$	$\Theta_{52} = [0; 2\bar{1}\bar{1}33\bar{1}] = 0,389648$	$\Theta_{84} = [0; 122\bar{1}3\bar{1}\bar{2}] = 0,70225$
$\Theta_{19} = [0; 32\bar{1}] = 0,29709$	$\Theta_{53} = [0; 2\bar{1}\bar{1}32\bar{1}] = 0,389916$	$\Theta_{85} = [0; 1223\bar{1}\bar{2}] = 0,70938$
$\Theta_{20} = [0; 32\bar{1}3\bar{1}\bar{2}] = 0,29774$	$\Theta_{54} = [0; 2\bar{1}\bar{1}3\bar{1}\bar{2}] = 0,39076$	$\Theta_{86} = [0; 1222\bar{1}] = 0,70783$
$\Theta_{21} = [0; 32\bar{1}\bar{3}] = 0,297773$	$\Theta_{55} = [0; 2\bar{1}\bar{1}\bar{3}] = 0,390891$	$\Theta_{87} = [0; 12\bar{1}\bar{1}3\bar{1}\bar{2}] = 0,71903$
$\Theta_{22} = [0; 333\bar{1}] = 0,302444$	$\Theta_{56} = [0; 223\bar{1}] = 0,409544$	$\Theta_{88} = [0; 12\bar{1}\bar{1}\bar{1}3\bar{1}\bar{2}] = 0,725381$
$\Theta_{23} = [0; 333\bar{1}\bar{2}] = 0,30025$	$\Theta_{57} = [0; 223\bar{1}\bar{2}] = 0,40968$	$\Theta_{89} = [0; \bar{1}\bar{2}] = 0,73206$
$\Theta_{24} = [0; 332\bar{1}] = 0,30330$	$\Theta_{58} = [0; 22\bar{1}] = 0,42265$	$\Theta_{90} = [0; 12\bar{1}3\bar{1}\bar{2}] = 0,73620$
$\Theta_{25} = [0; 33\bar{1}\bar{2}] = 0,30600$	$\Theta_{59} = [0; 22\bar{1}3\bar{1}\bar{2}] = 0,42398$	$\Theta_{91} = [0; 12\bar{1}\bar{3}] = 0,73624$
$\Theta_{26} = [0; 33\bar{1}2\bar{1}\bar{3}] = 0,30603$	$\Theta_{60} = [0; 22\bar{1}\bar{3}] = 0,424042$	$\Theta_{92} = [0; 133\bar{1}] = 0,765465$
$\Theta_{27} = [0; 33\bar{1}3\bar{1}\bar{2}] = 0,30638$	$\Theta_{61} = [0; 233\bar{1}] = 0,433577$	$\Theta_{93} = [0; 133\bar{1}\bar{2}] = 0,76569$
$\Theta_{28} = [0; 33\bar{1}\bar{3}] = 0,306394$	$\Theta_{62} = [0; 233\bar{1}\bar{2}] = 0,43365$	$\Theta_{94} = [0; 13\bar{1}\bar{2}] = 0,78868$
$\Theta_{29} = [0; 2\bar{1}\bar{3}] = 0,358258$	$\Theta_{63} = [0; 23\bar{1}\bar{2}] = 0,44093$	$\Theta_{95} = [0; \bar{1}\bar{3}] = 0,791287$
$\Theta_{30} = [0; 2\bar{1}3\bar{1}\bar{2}] = 0,35859$	$\Theta_{64} = [0; 23\bar{1}] = 0,441742$	
$\Theta_{31} = [0; 2\bar{1}333\bar{1}] = 0,361292$	$\Theta_{65} = [0; 1\bar{1}\bar{3}] = 0,558256$	
$\Theta_{32} = [0; 2\bar{1}33\bar{1}\bar{2}] = 0,36158$	$\Theta_{66} = [0; 1\bar{1}3\bar{1}\bar{2}] = 0,55905$	
$\Theta_{33} = [0; 2\bar{1}33\bar{1}] = 0,361602$		
$\Theta_{34} = [0; 2\bar{1}2\bar{1}\bar{3}] = 0,365455$		

Figure 5: Table of Θ 's.

4.2.3 Case $i_1 = i_2 \pmod{2}$

Now consider case $i_1 = i_2 \pmod{2}$ (see picture):



We will suppose case $a_{i_2} = 1$, $a_{i_2-1} = 3$ doesn't take place.

We can substitute the following γ 's and δ 's into (12.1):

$$\Theta_{\gamma'} = \Theta_{66}, \quad \Theta_{\gamma''} = \Theta_{63}, \quad \Theta_{\delta'} = \Theta_{90}, \quad \Theta_{\delta''} = \Theta_3,$$

and instead of (12.2) we will get

$$0,253 < q \frac{1+p\Theta_{63}}{1+p'\Theta_3} \frac{1+p\Theta_{66}}{1+p'\Theta_{90}}. \quad (12.5)$$

This choose of $\Theta_{\gamma'}$ and $\Theta_{\delta''}$ is fine, if the following inequality takes place:

$$\delta' - \delta'' > 1,4(\gamma' - \gamma''), \quad (12.6)$$

where

$$\Theta_{\gamma'} = \Theta_{68}, \quad \Theta_{\gamma''} = \Theta_{65}, \quad \Theta_{\delta'} = \Theta_{28}, \quad \Theta_{\delta''} = \Theta_1,$$

so we can rewrite (12.6) as

$$0,253 < q \frac{1+p\Theta_{65}}{1+p'\Theta_1} \frac{1+p\Theta_{68}}{1+p'\Theta_{28}}. \quad (12.7)$$

We can notice that (12.7) follows from (12.5). Indeed, that follows from inequality

$$0,253 \frac{1+p'\Theta_3}{1+p'\Theta_{63}} \frac{1+p\Theta_{90}}{1+p\Theta_{66}} > 0,269 \frac{1+p'\Theta_1}{1+p'\Theta_{65}} \frac{1+p\Theta_{28}}{1+p\Theta_{68}}$$

or inequality

$$\frac{1+p\Theta_{65}}{1+p\Theta_{63}} \frac{1+p\Theta_{68}}{1+p\Theta_{66}} \frac{1+p'\Theta_3}{1+p'\Theta_1} \frac{1+p'\Theta_{90}}{1+p'\Theta_{28}} > 1,064.$$

Here Freiman even omits comment «which is checked directly».

Overall, we proved that (12.5) is enough for condition to be good.

Suppose

$$\frac{\Delta_1}{\Delta_2} < 3,8 \quad (12.8)$$

or

$$\frac{1}{q} \frac{1+p'\Theta_1}{1+p\Theta_1} \frac{1+p'\Theta_{95}}{1+p\Theta_{95}} < 3,8.$$

Then (12.5) takes place. Indeed, it is so, if

$$\frac{1}{3,8} \frac{1+p'\Theta_1}{1+p'\Theta_1} \frac{1+p\Theta_{95}}{1+p\Theta_{95}} > 0,253 \frac{1+p'\Theta_3}{1+p'\Theta_{63}} \frac{1+p\Theta_{90}}{1+p\Theta_{66}}.$$


The last inequality is transformed into

$$1,04 > \frac{1+p'\Theta_3}{1+p'\Theta_1} \frac{1+p'\Theta_{90}}{1+p'\Theta_{95}} \frac{1+p\Theta_1}{1+p\Theta_{63}} \frac{1+p\Theta_{95}}{1+p\Theta_{66}},$$

which is easy to check.

4.2.4 Case $i_1 \neq i_2 \pmod{2}$

Now consider case $i_1 \neq i_2 \pmod{2}$ and, again, case $a_{i_2} = 1, a_{i_2-1} = 3$ doesn't take place. Suppose i_1 is even (see picture):



Taking

$$\Theta_{\gamma'} = \Theta_{66}, \Theta_{\gamma''} = \Theta_{59}, \Theta_{\delta'} = \Theta_3, \Theta_{\delta''} = \Theta_{90},$$

rewrite (12.1) as

$$0,2885 < q \frac{1+p\Theta_{59}}{1+p'\Theta_3} \frac{1+p\Theta_{66}}{1+p'\Theta_{90}}. \quad (12.9)$$

Suppose

$$\frac{\Delta_1}{\Delta_2} < 3,43 \quad (12.10)$$

or

$$\frac{1}{q} \frac{1+p'\Theta_1}{1+p\Theta_1} \frac{1+p'\Theta_{95}}{1+p\Theta_{95}} < 3,43.$$

Then (12.9) takes place. Indeed, it is so, if

$$\frac{1}{3,43} \frac{1+p'\Theta_1}{1+p\Theta_1} \frac{1+p'\Theta_{95}}{1+p\Theta_{95}} > 0,2885 \frac{1+p'\Theta_3}{1+p\Theta_{59}} \frac{1+p'\Theta_{90}}{1+p\Theta_{66}}.$$

or

$$1,0105 > \frac{1+p'\Theta_3}{1+p'\Theta_1} \frac{1+p'\Theta_{90}}{1+p'\Theta_{95}} \frac{1+p\Theta_1}{1+p\Theta_{59}} \frac{1+p\Theta_{95}}{1+p\Theta_{66}},$$

increasing the right part, obtain

$$1,0105 > 0,9896 \frac{1+p \cdot 1,0551 + p^2 \cdot 0,20875}{1+p \cdot 0,983 + p^2 \cdot 0,237},$$

which is easily checked.

5 Freiman's constant. Initial set of rectangles

All these rectangles satisfy *_here will be a complete list of conditions._*
 At least all of them are good, in terms of section 4.

$$\begin{aligned}
 1) & \quad \{ 3 \underset{\circ}{4} 3 \} \\
 2) & \quad \{ 3 \text{I} 3 \underset{\circ}{4} 3 \text{I} 2 \} \\
 3) & \quad \{ \text{I} 2 3 \underset{\circ}{4} 4 3 \text{I} \} \\
 4) & \quad \{ 3 \text{I} 3 \underset{\circ}{4} 3 \text{I} 3 \} \\
 5) & \quad \{ 2 \text{I} \text{I} 2 3 \underset{\circ}{4} 4 3 2 2 \} \\
 6) & \quad \{ 3 \text{I} \text{I} 2 3 \underset{\circ}{4} 4 3 2 3 \} \\
 7) & \quad \{ 3 \text{I} \text{I} 2 3 \underset{\circ}{4} 4 3 \underset{\circ}{2} 2 \} \\
 8+2n, \kappa) & \quad \{ \overset{\kappa}{3} \text{I} \overline{\overset{n}{3} \text{I} 3 \text{I} 2 \text{I} 3 \text{I} 3 \text{I} \text{I} 2 3 \underset{\circ}{4} 4 3 2 2 \overline{\overset{n}{3} \text{I} 3 \text{I} 2 \text{I} 3}}^{\kappa} \} \quad (\text{I3.I}) \\
 9+2n, \kappa) & \quad \{ \overset{\kappa+1}{3} 2 \text{I} 3 \overline{\overset{n}{3} \text{I} 3 \text{I} 2 \text{I} 3 \text{I} 3 \text{I} \text{I} 2 3 \underset{\circ}{4} 4 3 2 2 \overline{\overset{n}{3} \text{I} 3 \text{I} 2 \text{I} 3}}^{\kappa} 3 \text{I} 3 \} \quad (\text{I3.2}) \\
 9+2n, \kappa, p) & \quad \{ \overset{p}{3} \overline{\overset{\kappa}{3} \text{I} 3} 2 \text{I} 3 \text{I} 3 \overline{\overset{n}{3} \text{I} 3 \text{I} 2 \text{I} 3 \text{I} \text{I} 2 3 \underset{\circ}{4} 4 3 2 2 \overline{\overset{n}{3} \text{I} 3 \text{I} 2 \text{I} 3 \text{I} 3 \text{I} 2 \text{I} 3}}^{\kappa} \overset{p}{3} \text{I} 3 \} \quad (\text{I3.3})
 \end{aligned}$$

Figure 6: Freiman's constant. Initial set of rectangles.

Contents

1	Introductory notation	1
1.1	Markov and Lagrange values	1
1.2	Centered sequence	1
1.3	Rectangle	1
1.4	Horizontal rectangle	2
1.5	Subrectangle	2
1.6	Geometrical interpretation	2
2	Calculations	3
2.1	Length of Δ_1 or Δ_2	3
2.2	Rectangle aspect ratio	3
3	Rectangle boundaries	4
3.1	Resection	4
3.2	Shortened rectangle	4
3.2.1	Case $i_1 = i_2 = 0 \pmod{2}$	4
3.2.2	Other cases	5
3.3	Explanation of shortened	5
3.4	How we ensure centeredness	5
3.5	Formal	6
3.5.1	Bounds for Δ'	6
3.5.2	Bounds for Δ''	7
3.5.3	Case $i_1 \not\equiv i_2 \pmod{2}$	7
4	Good rectangle	9
4.1	Definition	9
4.2	Sufficient conditions of goodness	9
4.2.1	Results	9
4.2.2	Universal 2, 9 bound	10
4.2.3	Case $i_1 = i_2 \pmod{2}$	12
4.2.4	Case $i_1 \not\equiv i_2 \pmod{2}$	13
5	Freiman's constant. Initial set of rectangles	14