Supplementary material

Which drug–hearing-loss signals in FAERS survive adjustment and independent replication?

Oleg Tropynin

Table S1. READUS-PV checklist

Item	READUS-PV recommendation (abridged)	Where addressed
1a	Identify the study as a disproportionality analysis; name data type and database(s)	Title
1b	Name the adverse event(s) and/or drug(s)	Title (hearing loss); abstract
2a	Drugs, event, frequency, existing knowledge	Introduction, paragraph 1
2b	Rationale (routine PV, safety profile or pre-specified hypothesis)	Introduction, paragraphs 2–3
2c	Why ICSR databases and disproportionality suit the gap	Introduction, paragraphs 1–3
3	Objectives: event(s), drug(s), reference group, pre-specified hypotheses	Introduction, last paragraph; Methods 2.4
4a	Identify design and data type	Methods 2.1
4b	Outline primary and sensitivity analyses, case-by-case analysis	Methods 2.4–2.7; PROTOCOL.md (repository)
5a	Database names, custodians, coverage, thesaurus	Methods 2.1 (FDA, Health Canada; MedDRA PTs)
5b	Extraction dates; pre-processing and exclusions	Methods 2.1–2.2; Results 3.1
6a	Study population and restrictions	Methods 2.4 (restricted population)
6b	Key variables	Methods 2.3
6c	Grouping of drugs/events; ingredients vs trade names; drug role	Methods 2.2–2.3
6d	Additional data sources and how they interact with ICSRs	Methods 2.5 (US labels), 2.7 (PubMed)
7a	Descriptive analyses	Table S4
7b	Disproportionality measures and thresholds	Methods 2.4
7c	Sensitivity analyses and confounding control	Methods 2.4, 2.6; Results 3.3 (post hoc, labelled)
7d	Variables and criteria for case-by-case analysis	Methods 2.7; Table S3 (rule-based categories)
7e	Statistical methods for other data sources	Methods 2.5 (AUC, bootstrap), 2.6
8a	Number of ICSRs at each stage with reasons	Results 3.1
8b	Demographic and clinical characteristics of cases	Table S4
9	All results with confidence intervals; sensitivity analyses	Tables 1–3, S2, S5; Figures 1–2
10	Case-by-case analysis	Results 3.5; Table S3
11	Key results in context; expected vs emerging signals	Discussion, paragraphs 1–2
12a	External validity	Discussion; Limitations
12b	Relevance to clinical practice	Discussion, paragraph 4
12c	Further study designs	Discussion, paragraph 4; Conclusions
13	Limitations; disproportionality cannot prove causation or measure incidence	Limitations
14a	Funding	Declarations
14b	Conflicts of interest	Declarations
14c	Institutional approval	Declarations (not applicable)
14d	Data, code (software versions) and protocol availability	Declarations

Table S2. Established ototoxic drugs: crude and adjusted estimates (hearing loss)

FAERS crude: all reports; FAERS adjusted: Mantel–Haenszel OR in the restricted population (Methods); Canada: crude ROR, all reports.

Ingredient	FAERS cases	FAERS ROR (95% CI)	Adj. cases	FAERS adj. OR (95% CI)	Canada cases	Canada ROR (95% CI)
Gentamicin	315	9.66 (8.63–10.81)	181	11.37 (9.80–13.21)	131	33.59 (27.98–40.34)
Tobramycin	367	5.36 (4.83–5.94)	248	4.83 (4.25–5.47)	31	17.10 (11.89–24.58)
Amikacin	833	16.83 (15.69–18.07)	472	12.70 (11.57–13.94)	35	145.90 (97.39–218.57)
Streptomycin	18	5.76 (3.61–9.20)	12	7.92 (4.47–14.05)	9	6.28 (3.24–12.16)
Cisplatin	1206	4.61 (4.35–4.88)	730	4.26 (3.96–4.59)	138	31.21 (26.12–37.28)
Carboplatin	761	1.96 (1.83–2.11)	251	0.82 (0.72–0.92)	112	19.09 (15.73–23.16)
Oxaliplatin	121	0.44 (0.37–0.53)	106	0.46 (0.38–0.56)	1	0.14 (0.02–1.02)
Vancomycin	365	2.25 (2.03–2.49)	215	2.06 (1.80–2.36)	60	9.25 (7.14–11.98)
Furosemide	412	1.73 (1.57–1.91)	255	1.31 (1.16–1.48)	61	10.18 (7.87–13.16)
Hydroxychloroquine	556	2.10 (1.93–2.28)	327	1.73 (1.55–1.93)	88	2.21 (1.78–2.73)
Quinine	68	5.83 (4.59–7.42)	41	6.10 (4.47–8.31)	4	6.10 (2.27–16.42)
Sildenafil	894	3.99 (3.73–4.26)	725	3.93 (3.65–4.24)	32	4.18 (2.95–5.94)
Tadalafil	720	4.52 (4.20–4.87)	594	4.22 (3.89–4.58)	35	4.19 (2.99–5.85)
Vardenafil	6	1.31 (0.59–2.93)	4	0.93 (0.35–2.49)	4	17.01 (6.21–46.60)
Deferoxamine	51	8.97 (6.78–11.86)	50	11.78 (8.87–15.66)	34	36.29 (25.39–51.86)
Teprotumumab	61	124.18 (91.08–169.32)	61	145.26 (106.02–199.02)	1	–

Table S3. Case-level review of the 36 robust signals without a US label mention (post-hoc strict definition)

Category: A report cluster (≥50% of cases in reports listing ≥15 drugs); B ≥50% of cases co-report an established ototoxic drug; C stimulated or mass-exposure reporting; D indication or comorbidity linked to the event; K drug on the a-priori ototoxic list; E none of these. Rules applied in that order (K first). Profiles describe all FAERS cases (drug as suspect, event reported). Top country and year: share of cases from the most frequent country and receipt year.

Drug	Event	Cat.	Cases	Top country	Top year	Consumer	Median drugs	≥15 drugs	Ototoxic co-drug	Note
Linezolid	HL	B	350	US 18%	2025 11%	6%	9	21%	75%	nan
Clofazimine	HL	B	194	US 32%	2019 19%	8%	10	26%	78%	nan
Polymyxin B	HL	B	139	US 69%	2018 9%	78%	4	4%	78%	otic use: top indications ear infection / otitis externa
Mesna	HL	B	69	CA 35%	2016 26%	10%	7	4%	83%	nan
Rifabutin	HL	B	66	CA 30%	2024 30%	9%	10	18%	88%	nan
Rofecoxib	HL	C	245	US 63%	2006 42%	7%	6	16%	33%	nan
Tozinameran	HL	C	72	US 32%	2022 26%	74%	7.5	38%	8%	nan
Elasomeran	HL	C	20	US 85%	2023 35%	75%	8	35%	20%	nan
Minerals	HL	D	33	CA 27%	2018 18%	76%	9	15%	48%	non-specific supplement category; elderly multi-morbid patients
Etanercept	HL	E	2577	US 65%	2016 17%	45%	1	13%	18%	weak (adjusted OR 1.14) despite 2,001 cases

Pregabalin	HL	E	1795	US 81%	2021 22%	61%	3	15%	12%	1,795 cases, 61 % consumer, few co-reported ototoxics; alpha-2-delta ligand; partial label wording (ambiguous)
Amlodipine	HL	E	841	US 27%	2021 15%	69%	4	13%	15%	tinnitus labelled; hearing loss not; elderly hypertensive patients
Tocilizumab	HL	E	615	CA 45%	2020 23%	41%	8	37%	33%	weak; 45 % Canadian, 37 % drug-dense reports
Ondansetron	HL	E	157	US 87%	2020 22%	40%	3	18%	18%	supportive-care drug; median 28 reactions per report
Travoprost	HL	E	143	US 59%	2019 22%	76%	1	0%	3%	topical prostaglandin analogue; elderly glaucoma patients, 76 % consumer
Glycopyrronium	HL	E	120	US 72%	2025 32%	36%	3	3%	5%	inhaled LAMA in COPD; 32 % of cases in 2025
Interferon Alfa-2B	HL	E	27	US 41%	2013 22%	22%	2	4%	7%	hearing loss with interferon reported in the literature
Candesartan	Tin	A	540	GB 63%	2023 42%	32%	43	82%	42%	nan
Cholecalciferol	Tin	A	436	CA 33%	2025 23%	21%	75	82%	80%	nan
Tozinameran	Tin	A	183	CA 40%	2025 30%	35%	80	69%	62%	nan
Perindopril	Tin	A	162	CA 56%	2023 17%	23%	25.5	80%	74%	nan
Aminosalicylic Acid	Tin	A	97	CA 68%	2025 53%	5%	219	97%	95%	nan
Indapamide	Tin	A	45	PL 49%	2018 22%	20%	16	56%	60%	nan
Choline	Tin	A	4	US 100%	2013 100%	75%	23	100%	0%	4 cases, all in one 2013 HIV-treatment series
Finasteride	Tin	C	305	GB 37%	2024 18%	61%	1	1%	1%	nan
Rofecoxib	Tin	C	262	US 42%	2006 34%	19%	10	43%	47%	nan
Elasomeran	Tin	C	14	US 50%	2023 43%	64%	8	36%	29%	nan
Levothyroxine	Tin	D	992	GB 34%	2023 27%	53%	8	47%	23%	thyroid disease; 47 % of cases in drug-dense reports
Valsartan	Tin	D	476	US 41%	2022 10%	69%	5	17%	26%	hypertension / heart failure; also US patient-programme reporting
Olmesartan	Tin	D	85	US 34%	2020 20%	54%	5	21%	24%	hypertension, associated with tinnitus
Liothyronine	Tin	D	46	US 50%	2023 37%	76%	6	33%	2%	thyroid disease / over-replacement
Chlorthalidone	Tin	D	18	CA 44%	2026 39%	56%	10	39%	0%	hypertension; 18 cases
Galcanezumab	Tin	D	3	US 33%	2018 33%	33%	1	0%	0%	migraine, itself associated with tinnitus; 3 cases
Clonazepam	Tin	E	393	US 55%	2017 13%	69%	5	27%	6%	tinnitus on benzodiazepine withdrawal is described in the literature
Penicillin G Procaine	Tin	E	5	US 60%	2022 40%	40%	3	0%	0%	procaine (Hoigne) reactions include auditory symptoms; 5 cases
Paromomycin	Tin	K	4	US 50%	2017 25%	75%	2.5	0%	0%	aminoglycoside (oral); class effect plausible; 4 cases

Table S4. Characteristics of FAERS hearing-loss and tinnitus reports and all other reports (READUS-PV item 8b)

	Hearing loss	Tinnitus	All other reports
Reports, n	83,914	42,917	19,765,932
Age, median (years)	67	55	58

Age			
<18 (%)	2.4	1.2	3.5
18-44 (%)	6.9	16.3	12.5
45-64 (%)	15.6	30.0	20.2
≥65 (%)	30.6	17.4	20.7
unknown (%)	44.5	35.1	43.1
Sex			
female (%)	52.9	58.3	52.7
male (%)	39.3	32.2	34.3
unknown (%)	7.8	9.5	13.0
Reporter			
consumer (%)	60.0	53.7	44.3
physician (%)	16.2	16.9	21.6
pharmacist (%)	3.2	4.7	6.1
other health professional (%)	12.6	14.2	18.6
lawyer (%)	0.8	1.8	2.5
unknown (%)	7.2	8.7	6.9
Country			
United States (%)	69.9	64.0	71.3
other (%)	28.6	32.8	27.0
unknown (%)	1.5	3.2	1.7
FDA receipt			
2004-2012 (%)	13.6	21.2	15.6
2013-2018 (%)	30.7	34.0	31.8
2019-2026 (%)	55.7	44.8	52.6

Table S5. All robust signals (primary definition: adjusted FAERS signal replicated in Canada Vigilance)

FAERS: adjusted MH OR; Canada: crude ROR, all reports; Label: any US label mentions the event.

Ingredient	Event	FAERS cases	FAERS adj. OR (95% CI)	Canada cases	Canada ROR (95% CI)	Label
Amikacin	HL	833	12.70 (11.57–13.94)	35	145.90 (97.39–218.57)	yes
Gentamicin	HL	315	11.37 (9.80–13.21)	131	33.59 (27.98–40.34)	yes
Deferoxamine	HL	51	11.78 (8.87–15.66)	34	36.29 (25.39–51.86)	yes
Agalsidase Alfa	HL	14	17.79 (7.75–40.82)	10	10.65 (5.66–20.02)	no
Ethacrynic Acid	HL	11	12.64 (4.61–34.64)	28	274.38 (164.45–457.82)	yes
Quinine	HL	68	6.10 (4.47–8.31)	4	6.10 (2.27–16.42)	yes
Streptomycin	HL	18	7.92 (4.47–14.05)	9	6.28 (3.24–12.16)	yes
Tobramycin	HL	367	4.83 (4.25–5.47)	31	17.10 (11.89–24.58)	yes
Elosulfase Alfa	HL	80	5.22 (4.17–6.53)	3	9.15 (2.90–28.86)	no
Azithromycin	HL	711	4.52 (4.10–4.98)	45	7.03 (5.22–9.46)	yes
Cisplatin	HL	1206	4.26 (3.96–4.59)	138	31.21 (26.12–37.28)	yes

Tadalafil	HL	720	4.22 (3.89–4.58)	35	4.19 (2.99–5.85)	yes
Laronidase	HL	49	5.08 (3.80–6.78)	3	37.74 (11.36–125.37)	no
Neomycin	HL	139	4.94 (3.78–6.45)	13	7.65 (4.41–13.28)	yes
Clofazimine	HL	194	5.03 (3.70–6.86)	5	13.85 (5.65–33.97)	no
Galsulfase	HL	51	4.86 (3.67–6.43)	5	68.64 (25.98–181.36)	yes
Sildenafil	HL	894	3.93 (3.65–4.24)	32	4.18 (2.95–5.94)	yes
Alendronate Sodium	HL	673	3.94 (3.60–4.31)	6	2.80 (1.25–6.27)	no
Deferasirox	HL	344	3.83 (3.41–4.29)	20	5.12 (3.29–7.97)	yes
Agalsidase Beta	HL	115	3.92 (3.22–4.78)	4	6.94 (2.57–18.71)	yes
Travoprost	HL	143	3.67 (3.10–4.34)	3	4.27 (1.37–13.36)	no
Pregabalin	HL	1795	3.22 (3.06–3.39)	34	2.68 (1.91–3.76)	no
Tofacitinib	HL	1885	3.09 (2.94–3.25)	78	1.32 (1.05–1.65)	no
Clarithromycin	HL	377	3.16 (2.75–3.63)	32	4.00 (2.82–5.68)	yes
Formoterol	HL	1018	2.84 (2.65–3.04)	24	1.62 (1.08–2.42)	no
Glycopyrrolate	HL	207	3.00 (2.61–3.45)	3	3.51 (1.12–10.96)	no
Ciprofloxacin	HL	638	2.75 (2.50–3.03)	50	3.51 (2.65–4.65)	yes
Covid-19 Vaccine	HL	28	3.97 (2.38–6.62)	7	2.33 (1.11–4.91)	no
Tozinameran	HL	72	2.97 (2.32–3.80)	116	5.17 (4.29–6.23)	no
Glycopyrronium	HL	120	2.69 (2.24–3.24)	9	3.07 (1.59–5.92)	no
Amlodipine	HL	841	2.30 (2.14–2.48)	22	2.23 (1.46–3.39)	no
Moxifloxacin	HL	355	2.46 (2.12–2.86)	35	6.21 (4.44–8.69)	yes
Latanoprost	HL	278	2.35 (2.07–2.66)	5	2.56 (1.06–6.17)	no
Ergocalciferol	HL	36	2.93 (2.04–4.20)	4	3.25 (1.21–8.70)	no
Rofecoxib	HL	245	2.32 (1.96–2.74)	15	3.19 (1.91–5.30)	no
Octreotide	HL	268	2.23 (1.95–2.54)	86	4.11 (3.32–5.10)	yes
Anifrolumab	HL	9	3.65 (1.89–7.06)	4	4.05 (1.51–10.87)	no
Vancomycin	HL	365	2.06 (1.80–2.36)	60	9.25 (7.14–11.98)	yes
Canakinumab	HL	83	2.23 (1.78–2.79)	18	19.27 (11.95–31.06)	no
Celecoxib	HL	501	1.98 (1.78–2.21)	30	1.71 (1.20–2.46)	yes
Erythromycin	HL	76	2.07 (1.60–2.68)	30	4.20 (2.93–6.04)	yes
Metoprolol	HL	385	1.79 (1.59–2.02)	14	2.28 (1.35–3.87)	no
Aflibercept	HL	167	1.84 (1.58–2.16)	28	2.02 (1.39–2.94)	no
Hydroxychloroquine	HL	556	1.73 (1.55–1.93)	88	2.21 (1.78–2.73)	yes
Polymyxin B	HL	139	2.98 (1.54–5.74)	9	6.36 (3.28–12.31)	no
Imatinib	HL	430	1.66 (1.51–1.83)	13	1.84 (1.07–3.18)	yes
Tiotropium	HL	659	1.59 (1.46–1.73)	18	1.70 (1.07–2.70)	no
Vitamin D	HL	21	2.72 (1.41–5.24)	7	2.20 (1.05–4.63)	no
Influenza Virus Vaccine	HL	48	2.05 (1.39–3.01)	9	2.72 (1.41–5.26)	no
Imipenem	HL	158	1.87 (1.37–2.57)	3	4.44 (1.42–13.88)	yes
Ondansetron	HL	157	1.64 (1.36–1.99)	13	4.53 (2.62–7.84)	no
Bisoprolol	HL	172	1.58 (1.30–1.91)	8	2.53 (1.26–5.08)	yes
Elasomeran	HL	20	2.11 (1.29–3.46)	19	2.89 (1.84–4.55)	no
Tocilizumab	HL	615	1.41 (1.28–1.55)	87	1.43 (1.16–1.77)	no
Piroxicam	HL	19	2.08 (1.25–3.45)	4	3.16 (1.18–8.47)	yes
Linezolid	HL	350	1.53 (1.24–1.89)	10	9.30 (4.95–17.47)	no
Tramadol	HL	482	1.36 (1.23–1.51)	17	2.79 (1.73–4.50)	yes
Interferon Alfa-2B	HL	27	1.81 (1.22–2.68)	6	2.72 (1.22–6.07)	no
Amphotericin B	HL	73	1.56 (1.19–2.06)	5	3.90 (1.61–9.43)	yes
Peginterferon Alfa-2B	HL	68	1.50 (1.18–1.92)	8	3.10 (1.54–6.22)	yes
Furosemide	HL	412	1.31 (1.16–1.48)	61	10.18 (7.87–13.16)	yes
Ravulizumab	HL	17	1.81 (1.12–2.91)	5	3.76 (1.56–9.10)	no
Alemtuzumab	HL	71	1.43 (1.12–1.82)	14	2.63 (1.55–4.45)	no
Etanercept	HL	2577	1.15 (1.10–1.20)	144	1.85 (1.57–2.19)	no

Minerals	HL	33	1.77 (1.10–2.85)	3	6.00 (1.91–18.81)	no
Rifabutin	HL	66	2.14 (1.07–4.30)	3	17.76 (5.54–56.92)	no
Mesna	HL	69	1.78 (1.01–3.14)	17	32.80 (19.87–54.15)	no
Gabapentin	HL	543	1.09 (0.97–1.21)	33	4.16 (2.94–5.87)	yes
Caffeine	HL	93	1.30 (0.96–1.75)	20	1.68 (1.08–2.60)	yes
Felodipine	Tin	417	69.99 (60.67–80.75)	6	10.23 (4.54–23.04)	yes
Mefloquine	Tin	99	27.21 (21.81–33.96)	9	21.97 (11.20–43.08)	yes
Amikacin	Tin	355	17.24 (15.12–19.65)	8	29.48 (14.34–60.62)	yes
Paromomycin	Tin	4	32.77 (11.83–90.75)	3	220.78 (52.74–924.20)	no
Ciprofloxacin	Tin	1242	12.10 (11.36–12.89)	61	5.27 (4.08–6.80)	yes
Duloxetine	Tin	1971	11.81 (11.25–12.41)	41	5.67 (4.16–7.73)	yes
Bisoprolol	Tin	571	12.50 (11.20–13.96)	8	3.09 (1.54–6.20)	yes
Candesartan	Tin	540	10.63 (9.49–11.89)	25	2.77 (1.87–4.11)	no
Finasteride	Tin	305	9.72 (8.65–10.92)	5	3.23 (1.34–7.80)	no
Choline	Tin	4	20.36 (7.46–55.54)	5	5.92 (2.44–14.33)	no
Bupropion	Tin	1224	7.50 (7.06–7.97)	85	5.99 (4.82–7.44)	yes
Tozinameran	Tin	183	7.18 (5.67–9.10)	186	10.51 (9.05–12.21)	no
Tobramycin	Tin	206	6.09 (5.18–7.15)	12	7.76 (4.38–13.77)	yes
Paroxetine	Tin	608	5.31 (4.88–5.79)	38	5.59 (4.05–7.72)	yes
Quinine	Tin	41	6.65 (4.76–9.28)	10	19.20 (10.16–36.30)	yes
Levothyroxine	Tin	992	5.09 (4.72–5.49)	12	2.92 (1.65–5.16)	no
Azithromycin	Tin	382	5.00 (4.42–5.66)	29	5.46 (3.78–7.89)	yes
Covid-19 Vaccine	Tin	32	7.78 (4.40–13.77)	10	4.08 (2.18–7.61)	no
Venlafaxine Hydrochloride	Tin	796	4.65 (4.30–5.04)	6	2.40 (1.07–5.35)	yes
Desvenlafaxine	Tin	244	4.69 (4.11–5.35)	11	6.48 (3.57–11.77)	yes
Gentamicin	Tin	63	5.40 (4.05–7.20)	33	9.32 (6.59–13.19)	yes
Hydrochlorothiazide	Tin	777	4.29 (3.93–4.69)	23	1.72 (1.14–2.59)	yes
Liothyronine	Tin	46	5.24 (3.89–7.06)	3	15.55 (4.89–49.38)	no
Doxycycline	Tin	229	4.46 (3.83–5.20)	9	2.52 (1.31–4.85)	no
Tadalafil	Tin	320	4.13 (3.67–4.65)	13	1.87 (1.08–3.23)	yes
Escitalopram Oxalate	Tin	616	3.98 (3.55–4.46)	4	15.02 (5.52–40.85)	yes
Clarithromycin	Tin	258	4.15 (3.55–4.84)	17	2.57 (1.59–4.14)	yes
Cholecalciferol	Tin	436	4.36 (3.52–5.39)	10	3.68 (1.97–6.86)	no
Dimenhydrinate	Tin	89	5.66 (3.40–9.42)	5	2.46 (1.02–5.93)	no
Moxifloxacin	Tin	320	3.92 (3.39–4.54)	26	5.59 (3.79–8.25)	yes
Clonazepam	Tin	393	3.56 (3.20–3.96)	18	4.23 (2.66–6.74)	no
Esomeprazole	Tin	966	3.43 (3.16–3.72)	13	2.48 (1.44–4.29)	yes
Telmisartan	Tin	240	3.93 (3.14–4.91)	14	3.17 (1.87–5.37)	yes
Losartan	Tin	193	3.46 (2.96–4.04)	7	2.68 (1.28–5.65)	yes
Penicillin G Procaine	Tin	5	7.07 (2.92–17.12)	5	10.34 (4.25–25.17)	no
Minocycline	Tin	57	3.79 (2.88–4.99)	8	3.38 (1.68–6.79)	yes
Amlodipine	Tin	580	3.00 (2.74–3.29)	21	2.59 (1.69–3.99)	yes
Naproxen	Tin	658	2.99 (2.72–3.28)	34	3.00 (2.14–4.21)	yes
Perindopril	Tin	162	3.62 (2.67–4.91)	5	6.53 (2.69–15.81)	no
Buspirone Hydrochloride	Tin	46	3.57 (2.65–4.81)	4	10.22 (3.78–27.63)	yes
Aspirin	Tin	812	2.83 (2.62–3.06)	126	7.18 (6.00–8.59)	yes
Irbesartan	Tin	90	3.26 (2.62–4.06)	7	2.41 (1.15–5.08)	yes
Mirtazapine	Tin	310	2.99 (2.60–3.45)	9	2.74 (1.42–5.29)	yes
Metoprolol	Tin	297	2.79 (2.43–3.19)	12	2.38 (1.35–4.21)	yes
Influenza Virus Vaccine	Tin	29	3.67 (2.41–5.59)	10	3.70 (1.98–6.90)	no
Amitriptyline	Tin	232	2.79 (2.35–3.31)	22	3.06 (2.01–4.67)	yes
Naltrexone	Tin	218	2.70 (2.35–3.10)	18	4.13 (2.59–6.58)	yes
Atenolol	Tin	113	2.77 (2.24–3.43)	9	3.62 (1.88–6.98)	yes

Vortioxetine	Tin	224	2.73 (2.21–3.37)	5	3.68 (1.52–8.89)	yes
Cisplatin	Tin	373	2.48 (2.19–2.80)	26	6.45 (4.37–9.52)	yes
Terbinafine	Tin	68	2.83 (2.19–3.67)	7	3.27 (1.55–6.90)	yes
Cortisone	Tin	87	3.17 (2.15–4.66)	11	2.55 (1.41–4.62)	no
Rofecoxib	Tin	262	2.51 (2.11–2.99)	14	3.62 (2.14–6.14)	no
Isotretinoin	Tin	223	2.40 (2.10–2.75)	11	3.64 (2.01–6.59)	yes
Nortriptyline	Tin	40	2.87 (2.09–3.95)	8	4.20 (2.09–8.43)	yes
Dexchlorpheniramine	Tin	8	4.14 (2.06–8.31)	5	2.73 (1.13–6.58)	yes
Gabapentin	Tin	631	2.22 (2.01–2.45)	38	5.87 (4.25–8.10)	yes
Sildenafil	Tin	258	2.30 (2.01–2.63)	13	2.05 (1.19–3.53)	yes
Norfloxacin	Tin	138	4.21 (2.00–8.87)	4	4.72 (1.76–12.66)	yes
Fremanezumab-Vfrm	Tin	52	2.61 (1.97–3.46)	4	5.91 (2.20–15.88)	no
Citalopram	Tin	218	2.23 (1.93–2.57)	14	2.25 (1.33–3.81)	yes
Chlorthalidone	Tin	18	2.92 (1.84–4.64)	4	4.27 (1.59–11.44)	no
Omeprazole	Tin	450	2.05 (1.81–2.31)	15	1.84 (1.11–3.07)	yes
Doxepin	Tin	49	2.50 (1.73–3.63)	5	4.92 (2.03–11.90)	yes
Codeine	Tin	344	2.02 (1.70–2.40)	42	2.93 (2.16–3.98)	yes
Elasomeran	Tin	14	3.13 (1.68–5.82)	21	3.91 (2.54–6.02)	no
Tramadol	Tin	439	1.83 (1.62–2.07)	15	2.99 (1.80–4.98)	yes
Diltiazem	Tin	227	2.01 (1.56–2.58)	17	4.41 (2.73–7.13)	yes
Celecoxib	Tin	400	1.76 (1.51–2.04)	30	2.09 (1.46–3.00)	yes
Pregabalin	Tin	685	1.64 (1.50–1.79)	29	2.78 (1.93–4.02)	yes
Acetazolamide	Tin	15	2.63 (1.49–4.63)	5	7.90 (3.26–19.17)	yes
Pantoprazole	Tin	200	1.73 (1.47–2.04)	20	1.72 (1.11–2.68)	yes
Lidocaine	Tin	105	1.75 (1.44–2.12)	15	3.60 (2.16–5.99)	yes
Riociguat	Tin	73	1.84 (1.43–2.38)	8	2.08 (1.04–4.17)	no
Nitrofurantoin	Tin	102	1.98 (1.37–2.85)	11	2.10 (1.16–3.81)	no
Hydroxychloroquine	Tin	181	1.56 (1.33–1.82)	45	1.36 (1.01–1.82)	yes
Ibuprofen	Tin	388	1.48 (1.32–1.65)	46	3.26 (2.43–4.36)	yes
Olmesartan	Tin	85	1.67 (1.30–2.13)	9	4.27 (2.21–8.25)	no
Vancomycin	Tin	107	1.57 (1.28–1.93)	13	2.36 (1.37–4.07)	yes
Perindopril Erbumine	Tin	25	2.52 (1.26–5.05)	17	4.39 (2.72–7.09)	yes
Pravastatin	Tin	179	1.70 (1.23–2.37)	6	4.54 (2.03–10.15)	yes
Galcanezumab	Tin	3	3.81 (1.22–11.86)	6	6.95 (3.10–15.59)	no
Indomethacin	Tin	223	1.97 (1.20–3.22)	14	3.24 (1.91–5.49)	yes
Indapamide	Tin	45	1.90 (1.19–3.02)	7	3.83 (1.82–8.07)	no
Desloratadine	Tin	13	1.99 (1.15–3.43)	4	4.17 (1.56–11.18)	no
Quetiapine	Tin	340	1.20 (1.07–1.34)	19	2.08 (1.32–3.27)	yes
Cyclobenzaprine	Tin	39	1.40 (1.00–1.98)	5	3.79 (1.57–9.16)	yes
Trazodone Hydrochloride	Tin	70	1.24 (0.96–1.60)	12	5.14 (2.90–9.09)	yes
Aminosalicylic Acid	Tin	97	2.26 (0.85–6.04)	6	4.91 (2.19–10.99)	no