Draft version October 25, 2026
Typeset using LaTeX **twocolumn** style in AASTeX631

Rotation Periods of 41,842 Asteroids from Gaia DR3 Sparse Photometry, with Calibrated Reliability and a Null Test

Oleg Tropynin[1]

[1] *Independent researcher, Kazakhstan*

ABSTRACT

I derive rotation periods for every numbered asteroid with at least 25 Gaia DR3 transits (42,195 objects; 41,842 solved). I use a periodogram with season offsets, a linear phase term and a second-order Fourier series, and Gaia-centric geometry from numerical orbit integration. Asteroids with a known period are split once into a tuning half and a test half. Every choice, including the period-doubling rule and a logistic model for the probability that a period is correct, is fixed on the tuning half or on a null test in which the residuals are permuted within seasons. On the 4,834 test asteroids, confident (tier A) periods are correct within 3% in 86.3% of cases (1079/1250), and the probabilities are well calibrated. Without the null test, a threshold chosen for 80% precision on known asteroids would have let 6.9% of signal-free asteroids through. On known asteroids that resemble the unknown population, tier A is 85% correct (137/161). Agreement with three other Gaia DR3 analyses is 98%. Agreement with TESS (Vavilov & Carry) is 89%. I list 837 tier-A periods for asteroids with no published period in LCDB, DAMIT, about 180 VizieR tables or the Minor Planet Bulletin. The pipeline processes DR3 in about ten minutes on ten cores and is intended to run on Gaia DR4 on release day.

Keywords: Asteroid rotation (2211) — Light curves (918) — Main belt asteroids (2036) — Photometry (1234)

1. INTRODUCTION

Gaia DR3 gave sparse photometry for about 157,000 numbered asteroids (Tanga et al. 2023). Three groups have derived rotation periods or shapes from it. Ďurech & Hanuš (2023) used convex inversion, Cellino et al. (2024) used a genetic search and MacLennan et al. (2026) used a period search with phase-curve modelling. All three use the same data, and a large fraction of the asteroids with enough transits remains without a published period. Two things are needed before such periods can be used: an honest estimate of how often a period is wrong, and the same estimate for the asteroids that are *not* already in other catalogues. This paper provides both and makes the pipeline fast enough to rerun on Gaia DR4 (expected in December 2026, with about twice the time baseline). A pilot on 600 asteroids was reported separately (Tropynin 2026).

2. DATA AND GEOMETRY

All 20 bulk files of `sso_observation` (23.3 M CCD rows) were averaged per `transit_id`, giving 3,069,170 transits of 156,789 numbered asteroids. The median asteroid has 17 transits; 42,195 have at least 25 and are analysed here. Each transit takes the largest relative flux error of its CCDs, with a 0.03 mag floor.

Heliocentric positions come from osculating elements (JPL, epoch 2026 June) integrated back to 2014–2017 with a fourth-order Runge–Kutta scheme. The step is one day, and the integration includes the Sun and the four giant planets, using JPL Horizons vectors. Gaia's position is given in the data. Light time is iterated. Against JPL Horizons on 19 asteroids, the reduced magnitude agrees to 0.0007 mag, the observer distance to 0.0005 au and the phase angle to 0.009°. A two-body solution was rejected because its errors reach 0.13 mag.

3. PERIOD SEARCH

The reduced magnitude $G - 5\log_{10}(r\Delta)$ is fitted by weighted least squares. The model has one constant per observing season (seasons are separated by gaps longer than 60 d), a term linear in phase angle and a second-order Fourier series in rotational phase. Outliers beyond 5 median absolute deviations from the no-rotation model are removed in two passes.

The search scans periods of 2–1000 h on a grid with a step of $0.5/T$, where T is the time span. The 30 deepest

minima are then refined. A grid five times finer changed test precision by less than one point.

Most lightcurves are double-peaked, so at the true period the second harmonic dominates. If the first-harmonic amplitude A_1 exceeds $1.15A_2$, the doubled period is adopted. The factor 1.15 was chosen on the tuning half. A BIC comparison of fourth-order models at P and $2P$ was tried first and rejected, because it favoured the half period.

4. CALIBRATION, NULL TEST AND TIERS

The truth set is LCDB V4.0 (Warner et al. 2009), quality $U \geq 2$: 9,613 of the solved asteroids. A period counts as correct if it is within 3% of the LCDB value. The asteroids are split by a hash of their number into TUNE (4,779) and TEST (4,834). On TUNE I fitted a logistic model of the probability of a correct period. Its inputs are the χ^2 gain over the no-rotation model, the separation from the next non-alias minimum, the number of transits, the amplitude, the reduced χ^2, the period and A_1/A_2. On TEST the predicted probabilities match the observed fractions (Figure 2).

A threshold that gives 80% precision on TUNE (0.478) is not safe for unknown asteroids. To test it, I permuted the residuals of the no-rotation model within each season for 4,000 random unknown asteroids. This keeps the sampling, the errors and the season levels but destroys rotation. At 0.478, 6.9% of these signal-free asteroids still pass. At 0.70 the rate falls to 0.8% (Figure 3). Tier A is therefore defined as probability ≥ 0.70 and period < 100 h. Tier B covers 0.478–0.70. Best periods of 100–1000 h are flagged as slow candidates and never treated as confident, because only about 40% of them are correct.

5. RESULTS

Of 41,842 solutions, 5,594 are tier A and 3,727 are tier B; 769 are flagged as slow candidates. On TEST the results are:

- tier A: 1079 of 1250 correct (86.3%), with 87%, 85% and 80% in the 2–10, 10–30 and 30–100 h bands;
- the 171 tier-A errors: 156 other periods, 11 at $2P$ and 4 at $P/2$;
- tier B: 67% correct.

The comparison with LCDB is shown in Figure 1. Correct periods agree with LCDB to a median of 3×10^{-4}. The curvature error of the χ^2 minimum is about 25 times smaller than the scatter against Ďurech & Hanuš (2023), because a single period ignores the synodic drift. I therefore add $2.4 \times 10^{-4}P$ in quadrature, which gives a median $|\Delta P|/\sigma$ of 0.65.

Table 1. Agreement of tier-A periods with published catalogues (within 3%).

Catalogue	Data	n	Agree
Ďurech & Hanuš (2023)	Gaia DR3	3117	97.7%
Cellino et al. (2024)	Gaia DR3	2765	98.0%
MacLennan et al. (2026)	Gaia DR3	2580	97.9%
Vavilov & Carry (2025)	TESS	660	89.2%
Erasmus et al. (2020)	ATLAS	228	69.3%
McNeill et al. (2023)	TESS	774	52.6%

NOTE—Rows that use Gaia DR3 test consistency, not accuracy. For TEST asteroids that are also in LCDB, this work and the catalogue agree with LCDB in 320 vs 316 of 362 cases (Vavilov & Carry), 86 vs 89 of 114 (Erasmus et al.), and 302 vs 211 of 332 (McNeill et al.).

Transfer to unknown asteroids. The TEST asteroids are brighter and better studied than the asteroids without a period. The closest known analogue is the set of TEST asteroids that appear in none of the six catalogues of Table 1. Tier A is 85% correct on them (137/161; 95% interval 80–91%). I adopt this figure, not the overall 86%, for the new periods.

New periods. 912 tier-A asteroids have no period in LCDB or in the six catalogues. 75 of them do have a period in DAMIT (Ďurech et al. 2010), in about 180 VizieR tables with rotation periods (5.0 M rows) or in the Minor Planet Bulletin, volumes 50–53. Where such a value exists, this work agrees with it in 38 of 56 cases. The remaining 837 (602 at 2–10 h, 140 at 10–30 h, 95 at 30–100 h) are listed as new, with an expected precision of about 85%. Behrend's Geneva page and ZTF catalogues outside VizieR were not searched.

Pilot candidates. Of the 42 candidates of Tropynin (2026):

- 35 agree with this work;
- 4 flagged "low" are doubled here;
- for 3 flagged "high" (39932, 109412, 13686), the extended 1000 h search prefers slow solutions of low probability. These three should be treated as ambiguous.

6. LIMITATIONS AND PLANS FOR GAIA DR4




Figure 1. Period from this work against the LCDB period for TEST asteroids. Tier A is shown in blue and tier B in orange. The lines mark P, $2P$ and $P/2$.




Figure 2. Reliability diagram on TEST, all solutions. Bars are 95% binomial intervals; numbers give the count per bin.

The A_1/A_2 rule is a heuristic. A model-based comparison of P and $2P$ (ellipsoid or convex fit) would be more robust. The synodic effect is not modelled; a phase-angle-bisector correction is planned. Tumbling and slow rotators fall outside a single-period model. For DR4, all thresholds, the null test and the matched-control test will be fixed in advance, and injection–recovery tests on the real DR4 sampling will be added, so that results can be released within a day without re-tuning.

7. DATA AVAILABILITY

The catalogue (41,842 rows), the list of 837 new periods, the validation tables and the code will be archived on Zenodo [DOI to be assigned].

This work used data from the ESA mission Gaia, processed by the Gaia DPAC; JPL Horizons and the JPL Small-Body Database; LCDB; and VizieR (CDS, Strasbourg). The analysis was designed, coded and checked with the help of an AI assistant (Claude, Anthropic) under the author's direction. Two AI-based referee passes were used to stress-test the validation before submission.

REFERENCES

Cellino, A., et al. 2024, A&A, 687, A277

Ďurech, J., & Hanuš, J. 2023, A&A, 675, A24

Ďurech, J., Sidorin, V., & Kaasalainen, M. 2010, A&A, 513, A46

Erasmus, N., et al. 2020, ApJS, 247, 13

MacLennan, E. M., et al. 2026, A&A, 707, A131

McNeill, A., et al. 2023, AJ, 166, 152

Gaia Collaboration, Tanga, P., et al. 2023, A&A, 674, A12

Tropynin, O. 2026, Candidate rotation periods for 42 main-belt asteroids from Gaia DR3 sparse photometry, Zenodo, doi: 10.5281/zenodo.23624115

Vavilov, D., & Carry, B. 2025, A&A, 693, A66




Figure 3. Left: fraction of signal-free (permuted) asteroids that pass as a function of the probability threshold. Right: precision on TEST. Dashed: the TUNE 80%-precision threshold. Solid: the tier-A threshold.

Warner, B. D., Harris, A. W., & Pravec, P. 2009, Icarus, 202, 134