

Shapiro-Wilk Test for Normality (done on subsets because  $N > 5000$ ):

Dataset A subset: statistic = 0.9997, p-value = 0.8377

Dataset B subset: statistic = 0.9535, p-value = 0.0000

Anderson-Darling Test for Normality (on full datasets):

Dataset A subset:  $A_2 = 0.2776$

Significance level: 15.0%, Critical Value: 0.5760

Significance level: 10.0%, Critical Value: 0.6560

Significance level: 5.0%, Critical Value: 0.7870

Significance level: 2.5%, Critical Value: 0.9180

Significance level: 1.0%, Critical Value: 1.0920

Dataset B subset:  $A_2 = 8917.0256$

Significance level: 15.0%, Critical Value: 0.5760

Significance level: 10.0%, Critical Value: 0.6560

Significance level: 5.0%, Critical Value: 0.7870

Significance level: 2.5%, Critical Value: 0.9180

Significance level: 1.0%, Critical Value: 1.0920

Data is not normally distributed. Using Pearson Chi-square test on binned data...

Chi-square Test Results:

Chi-square statistic: 361918.8269772363

Degrees of freedom: 18


P-value: 0.0

KS Test Results (For Distributional Differences):


KS-statistic: 0.217063

P-value: 0.0


Histogram of Data



Boxplot Comparison



CDF of Datasets



Density (KDE) Comparison

