Practice workbook · Completing the square

Rewriting ax^2 + bx + c in the form a(x + p)^2 + q, or as q - a(x + p)^2 when the x squared term is negative, and stating the constants.

12 question parts: 1 worked-example retry and 11 further practice parts.
Name: _____ Class: _____ Date: _____

Practise one question type at a time. Attempt the part inside the red boundary; the rest of the question and any figures are kept for context. Write your working in the answer space.

The first question is a retry of the website worked example, without its solution. Check your answers online with 'Check my answer', or ask your teacher for the answer booklet.

OLevels.com SCHOOL

Type ID: FPM-QUA-03

1. Question 3 · part a · 3 marks

2023 JAN · Paper 02

Worked-example retry · Attempt part a only.

3

$$f(x) = 8x^2 + 10x - 3$$

Given that $f(x)$ can be written in the form $A(x + B)^2 + C$ where A, B and C are constants,

(a) find the value of A, the value of B and the value of C.

(3)

(b) Hence, or otherwise, find,

(i) the value of x for which $f(x)$ has a minimum,

(ii) the minimum value of $f(x)$.

(2)

The curve C has equation $y = f(x)$.

(c) Find the x coordinate of each of the points where C crosses the x-axis.

(2)

The straight line l has equation $y = 2x + 13$

(d) Use algebra to find the coordinates of the two points of intersection of C and l.

(4)

Using the same axes and the results of parts (b), (c) and (d),

(e) sketch the curve C and the straight line l.

(2)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

OLevels.COM SCHOOL

P 7 1 8 1 9 A 0 6 3 6

1. Question 3 · part a · 3 marks · Working and answer

Working and answer

OLevels.com SCHOOL

2. Question 10 · part a · 3 marks

2019 JUN · Paper 01 (regional R)

Further practice · Attempt part a only.

10 $f(x) = 6x - x^2 \qquad x \in \mathbb{R}$

Given that $f(x)$ can be written in the form $D(x + E)^2 + F$ where D, E and F are integers,

(a) find the value of D, the value of E and the value of F.

(3)

(b) Find

(i) the maximum value of $f(x)$,

(ii) the value of x for which the maximum occurs.

(2)

The curve C has equation $y = f(x)$

The curve S has equation $y = x^2 - 4x + 8$

The curve S intersects the curve C at two points.

(c) Find the coordinates of each of these two points.

(4)

The finite region R is bounded by the curve C and the curve S.

(d) Use algebraic integration to find the area of R.

(4)

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

OLevels.COM — SCHOOL —

2. Question 10 · part a · 3 marks · Working and answer

Working and answer

OLevels.com SCHOOL

3. Question 7 · part a · 2 marks

2020 NOV · Paper 01

Further practice · Attempt part a only.

7

$$f(x) = x^2 - 9x + 14$$

Given that $f(x)$ can be written in the form $(x + a)^2 + b$, where a and b are constants,

(a) find the value of a and the value of b.

(2)

(b) Hence, or otherwise, find

(i) the minimum value of $f(x)$

(ii) the value of x for which this minimum occurs.

(2)

The curve C has equation $y = f(x)$

The line l has equation $y = x + 5$

(c) Use algebra to find the coordinates of the points of intersection of C and l.

(4)

(d) Use algebraic integration to find the exact area of the finite region bounded by C and l.

(5)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

OLevels.com SCHOOL

P 6 2 2 8 0 A 0 1 4 3 2

3. Question 7 · part a · 2 marks · Working and answer

Working and answer

OLevels.com SCHOOL

4. Question 2 · part a · 3 marks

2022 JAN · Paper 01

Further practice · Attempt part a only.

2 $f(x) = 2x^2 - 12x + 5$

Given that $f(x)$ can be written in the form $f(x) = a(x + b)^2 + c$ where a, b and c are integers,

(a) find the value of a, the value of b and the value of c

(3)

(b) Hence find the set of values of x for which $f(x) - 37 > 0$

(4)

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

OLevels.com SCHOOL

4. Question 2 · part a · 3 marks · Working and answer

Working and answer

OLevels.com SCHOOL

5. Question 2 · part a · 3 marks

2024 JUN · Paper 01

Further practice · Attempt part a only.

2 $f(x) = 2x^2 + 4x + 9$

Given that $f(x)$ can be written in the form $A(x + B)^2 + C$, where A, B and C are integers,

(a) find the value of A, the value of B and the value of C

(3)

(b) Hence, or otherwise, find

(i) the value of x for which $\frac{1}{f(x)}$ is a maximum

(ii) the maximum value of $\frac{1}{f(x)}$

(2)

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

OLevels.COM — SCHOOL —

P 7 6 5 0 6 A 0 4 3 2

5. Question 2 · part a · 3 marks · Working and answer

Working and answer

OLevels.com SCHOOL

6. Question 2 · part a · 3 marks

2020 JAN · Paper 01 (regional R)

Further practice · Attempt part a only.

2 $f(x) = 7 + 4x - x^2$

(a) Write $f(x)$ in the form $a - b(x + c)^2$ where a, b and c are integers to be found. **(3)**

(b) Hence, or otherwise, find

(i) the maximum value of $f(x)$

(ii) the value of x for which this maximum occurs. **(2)**

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

OLevels.com SCHOOL

P 6 0 4 7 5 A 0 4 3 6

6. Question 2 · part a · 3 marks · Working and answer

Working and answer

OLevels.com SCHOOL

7. Question 2 · part a · 3 marks

2025 NOV · Paper 01

Further practice · Attempt part a only.

2 $f(x) = 2x^2 - 10x + 7$

Given that $f(x)$ can be written in the form $f(x) = a(x+b)^2 + c$ where a, b and c are rational numbers,

(a) find the value of a, the value of b and the value of c

(3)

(b) Hence, or otherwise, write down

(i) the minimum value of $f(x)$

(ii) the value of x at which this minimum value occurs.

(2)

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

OLevels.COM SCHOOL

P 7 8 9 6 4 A 0 4 3 6

7. Question 2 · part a · 3 marks · Working and answer

Working and answer

OLevels.com SCHOOL

8. Question 2 · part a · 4 marks

2021 JUN · Paper 02

Further practice · Attempt part a only.

2 $f(x) = 2 + \frac{4}{5}x - \frac{1}{25}x^2$

Given that $f(x)$ can be expressed in the form $A - B(x + C)^2$ where A, B and C are constants,

(a) find the value of A, the value of B and the value of C.

(4)

(b) Hence write down

(i) the maximum value of $f(x)$,

(ii) the value of x for which this maximum occurs.

(2)

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

OLevels.com SCHOOL

8. Question 2 · part a · 4 marks · Working and answer

Working and answer

OLevels.com SCHOOL

9. Question 11 · part a · 4 marks

2023 JUN · Paper 01

Further practice · Attempt part a only.

11 $f(x) = 10 + 6x - x^2$

Given that $f(x)$ can be written in the form $A(x + B)^2 + C$ where A, B and C are constants,

(a) find the value of A, the value of B and the value of C

(4)

(b) Hence, or otherwise, find

(i) the value of x for which $f(x)$ has its greatest value

(ii) the greatest value of $f(x)$

(2)

The curve C has equation $y = f(x)$

The curve S with equation $y = x^2 - x + 13$ intersects curve C at two points.

(c) Find the x coordinate of each of these two points.

(3)

(d) Use algebraic integration to find the exact area of the finite region bounded by the curve C and the curve S.

(5)

DO NOT WRITE IN THIS AREA DO NOT WRITE IN THIS AREA DO NOT WRITE IN THIS AREA

OLevels.com SCHOOL

9. Question 11 · part a · 4 marks · Working and answer

Working and answer

OLevels.com SCHOOL

10. Question 10 · part a · 4 marks

2024 NOV · Paper 01

Further practice · Attempt part a only.

10 $f(x) = 3 - 4x - 9x^2$

Given that $f(x)$ can be expressed in the form $A - B(x + C)^2$ where A, B and C are positive constants

(a) find the value of A, the value of B and the value of C

(4)

(b) Hence write down the maximum value of $f(x)$

(1)

The equation $f(x) = 0$ has roots α and β

Without solving the equation $f(x) = 0$

(c) form a quadratic equation, with integer coefficients, that has roots $\frac{3\alpha}{\beta}$ and $\frac{3\beta}{\alpha}$

(6)

(d) Show that $(x + y)^3 = x^3 + y^3 + 3xy(x + y)$

(1)

$g(x) = 3x^2 + qx + r$ where q and r are constants

The equation $g(x) = 0$ has roots $\alpha^2 - \beta$ and $\beta^2 - \alpha$ where α and β are the roots of the equation $f(x) = 0$

(e) Using your answer to part (d), find in simplified exact form, the value of q and the value of r

(6)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

OLevels.com SCHOOL

10. Question 10 · part a · 4 marks · Working and answer

Working and answer

OLevels.com SCHOOL

11. Question 2 · part a · 2 marks

2020 NOV · Paper 01 (regional R)

Further practice · Attempt part a only.

2 (a) Express $x^2 + 4x - 8$ in the form $(x + a)^2 + b$ where a and b are constants whose values are to be found.

(2)

(b) Use algebra to solve the simultaneous equations

$$y = x^2 + 4x - 8$$
$$y = 2x + 7$$

(5)

The curve C has equation $y = x^2 + 4x - 8$
The straight line L has equation $y = 2x + 7$

Using the same axes and the results of part (a) and part (b),

(c) sketch C and L, showing clearly the coordinates of the turning point of C and the coordinates of the points of intersection of C and L.

(4)

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

OLevels.com SCHOOL

11. Question 2 · part a · 2 marks · Working and answer

Working and answer

OLevels.com SCHOOL

12. Question 2 · part a · 3 marks

2023 NOV · Paper 01

Further practice · Attempt part a only.

2 $g(x) = 2x^2 + \frac{1}{2}x - 3$

(a) Express g(x) in the form $p(x+q)^2 + r$ where p, q and r are rational numbers to be found.

(3)

(b) Find

(i) the minimum value of g(x)

(ii) the value of x at which this minimum occurs.

(2)

$$h(x) = 2x^6 + \frac{1}{2}x^3 - 3$$

(c) Hence, or otherwise, write down

(i) the minimum value of h(x)

(ii) the value of x at which this minimum occurs.

(2)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

OLevels.COM — SCHOOL —

12. Question 2 · part a · 3 marks · Working and answer

Working and answer

OLevels.com SCHOOL