Type MS · Visual agnosia, prosopagnosia and pre-frontal damage

Answer 1 of 6 · Question 18 part a · 2 marks

2024 JUN · Paper 01 · extract 1 of 2 · find part a in this mark scheme extract

Question number	Answer	Mark
18(a)	**One** mark for accurate understanding of a relevant concept of neurological damage. **One** mark for exemplification of how a relevant concept of neurological damage can be used to explain the scenario. For example: • Visual agnosia is where a person is able to physically see an object with no problem but is unable to recognise it (1). The patient cannot identify the objects, such as a set of keys, but is able to recognise a familiar face, so this indicates they may be suffering from visual agnosia (1). Accept any other appropriate response.	(2)

Question number	Answer	Mark
18(b)	**One** mark for identification of a relevant improvement. **One** mark for justification, which must be linked to the improvement identified. For example: • Matthew could have used objects that may be considered more familiar to people than the ones he chose in his assessment of the patients (1) such as using a toothbrush instead of a games console which would be more familiar to everyone so that errors in recognition are not due to being less familiar with the object, giving his assessment more credibility (1). Accept any other appropriate response.	(2)

Type MS · Visual agnosia, prosopagnosia and pre-frontal damage

Answer 1 of 6 · Question 18 part a · 2 marks

2024 JUN · Paper 01 · extract 2 of 2 · find part a in this mark scheme extract

Type MS · Visual agnosia, prosopagnosia and pre-frontal damage

Answer 2 of 6 · Question 14 part whole · 2 marks

2019 JUN · Paper 01 · find part whole in this mark scheme extract

Question number	Answer	Mark
14	Up to **two** marks for defining what is meant by 'visual agnosia'. For example: • Visual agnosia is a condition where an individual can see but is unable to recognise what they can see (1). It may be that their eyes can see an everyday object like a table but are unable to name what it is when asked (1). Accept any other appropriate response.	**(2)**

Type MS · Visual agnosia, prosopagnosia and pre-frontal damage

Answer 3 of 6 · Question 14 part whole · 2 marks

2020 NOV · Paper 01 · find part whole in this mark scheme extract

Question number	Answer	Mark
14	**One** mark for accurate understanding of prosopagnosia. **One** mark for exemplification of how prosopagnosia can be used to explain the scenario. For example: • Prosopagnosia is a neurological condition which means the sufferer is unable to recognise faces of familiar people (1) which means Neha would struggle to continue working as an Art teacher as she would struggle to recognise each of her students (1). Accept any other appropriate response.	**(2)**

Type MS · Visual agnosia, prosopagnosia and pre-frontal damage

Answer 4 of 6 · Question 19 part whole · 2 marks

2021 NOV · Paper 01 · find part whole in this mark scheme extract

Question number	Answer	Mark
19	**One** mark for accurate understanding of visual agnosia. **One** mark for exemplification of how visual agnosia can be used to explain the scenario. For example: • Visual agnosia is where an individual can see but is unable to recognise and name what they can see (1). Carl can say that the patient is likely to see objects in the factory such as different ingredients for new chocolate bars, but he will be unable to recognise them, so will struggle to do his job (1). Accept any other appropriate response.	**(2)**

Type MS · Visual agnosia, prosopagnosia and pre-frontal damage

Answer 5 of 6 · Question 20 part whole · 2 marks

2022 JUN · Paper 01 · extract 1 of 2 · find part whole in this mark scheme extract

Question number	Answer	Mark
20	**One** mark for a conclusion made. **One** mark for justification of the conclusion through analysis/interpretation. For example: • The participants all appeared to have prosopagnosia (1) because none of them could recognise the familiar face but they were able to all recognise familiar objects and this is a symptom of prosopagnosia (1). Accept any other appropriate response.	**(2)**

Type MS · Visual agnosia, prosopagnosia and pre-frontal damage

Answer 5 of 6 · Question 20 part whole · 2 marks

2022 JUN · Paper 01 · extract 2 of 2 · find part whole in this mark scheme extract

Section E
Social influence – How do others affect you?

Type MS · Visual agnosia, prosopagnosia and pre-frontal damage

Answer 6 of 6 · Question 16 part whole · 2 marks

2024 JUN · Paper 01 · find part whole in this mark scheme extract

Question number	Answer	Mark
16	Up to **two** marks for description. For example: • Damage to the pre-frontal cortex can lead to emotional blunting which means the individual may find it difficult to feel emotions such as joy or sadness (1). Impulsive behaviour can also occur where an individual may not consider the consequences of their actions and therefore react with more violent behaviour (1). Accept any other appropriate response.	**(2)**