

# Introduction to biostatistics and machine learning

Olga Dethlefsen, Eva Freyhult, Bengt Sennblad, Payam Emami

2020-10-24



# Contents

<b>Preface</b>	<b>5</b>
<b>I Preliminary Mathematics</b>	<b>7</b>
<b>1 Mathematical notations</b>	<b>9</b>
1.1 Numbers . . . . .	9
1.2 Variables, constants and letters . . . . .	10
1.3 A precise language . . . . .	10
1.4 Using symbols . . . . .	10
1.5 Inequalities . . . . .	11
1.6 Indices and powers . . . . .	12
1.7 Exercises: notations . . . . .	12
Answers to selected exercises (notations) . . . . .	13
<b>2 Sets</b>	<b>15</b>
2.1 Definitions . . . . .	15
2.2 Basic set operations . . . . .	16
2.3 Venn diagrams . . . . .	16
2.4 Exercises: sets . . . . .	17
Answers to selected exercises (sets) . . . . .	18
<b>3 Functions</b>	<b>19</b>
3.1 Definitions . . . . .	19
3.2 Evaluating function . . . . .	19
3.3 Plotting function . . . . .	20
3.4 Standard classes of functions . . . . .	20
3.5 Piecewise functions . . . . .	21
3.6 Exercises: functions . . . . .	23
Answers to selected exercises (functions) . . . . .	23

<b>II</b>	<b>Introduction to R, R Studio and R markdown</b>	<b>25</b>
<b>4</b>	<b>R</b>	<b>27</b>
<b>5</b>	<b>R Studio</b>	<b>29</b>
<b>6</b>	<b>R Markdown</b>	<b>31</b>
<b>III</b>	<b>Probability</b>	<b>33</b>
<b>7</b>	<b>Probability: reasoning under uncertainty</b>	<b>35</b>
7.1	Introduction . . . . .	35
7.2	Basic set definitions . . . . .	35
7.3	Basic set operations . . . . .	36
7.4	Exercises . . . . .	36
7.5	Answers to exercises . . . . .	36
<b>8</b>	<b>Probability: random variables</b>	<b>37</b>
8.1	Random variables . . . . .	37
8.2	Discrete random variables . . . . .	38
<b>9</b>	<b>Summarising and visualising data</b>	<b>39</b>
<b>10</b>	<b>Linear regression</b>	<b>41</b>
10.1	Simple regression . . . . .	41
10.2	Multiple regression . . . . .	41

# Preface

This “bookdown” book contains teaching and learning materials prepared and used during “Introduction to biostatistics and machine learning” course organised by NBIS, National Bioinformatics Infrastructure Sweden. The course is open for PhD students, postdoctoral researcher and other employees in need of biostatistical skills within Swedish universities. The materials are geared towards life scientists wanting to be able to understand and use basic statistical and machine learning methods. It may also suits those already applying biostatistical methods but who have never gotten a chance to reflect on the basic statistical concepts, such as the commonly misinterpreted p-value.

More about the course <https://nbisweden.github.io/workshop-mlbiostatistics/>



## Part I

# Preliminary Mathematics





# Chapter 1

## Mathematical notations

### Aims

- to recapitulate the basic notations and conventions used in mathematics and statistics

### Learning outcomes

- to recognize natural numbers, integrals and real numbers
- to understand the differences between variables and constants
- to use symbols, especially Sigma and product notations, to represent mathematical operations

## 1.1 Numbers

- **Natural numbers,  $\mathbf{N}$ :** numbers such as 0, 1, 3, ...
- **Integers,  $\mathbf{Z}$ :** include negative numbers ..., -2, -1, 0, 1, 2
- **Rational numbers:** numbers that can be expressed as a ratio two integers, i.e. in a form  $\frac{a}{b}$ , where  $a$  and  $b$  are integers, and  $b \neq 0$
- **Real numbers,  $\mathbf{R}$ :** include both rational and irrational numbers
- **Reciprocal** of any number is found by dividing 1 by the number, e.g. reciprocal of 5 is  $\frac{1}{5}$
- **Absolute value** of a number can be viewed as its distance from zero, e.g. the absolute value of 6 is 6, written as  $|6| = 6$  and absolute value of -5 is 5, written as  $|-5| = 5$
- **Factorial** of a non-negative integer number  $n$  is denoted by  $n!$  and it is a product of all positive integers less than or equal to  $n$ , e.g.  $4! = 4 \cdot 3 \cdot 2 \cdot 1 = 24$

## 1.2 Variables, constants and letters

Mathematics gives us a precise language to communicate different concepts and ideas. To be able to use it it is essential to learn symbols and understand how they are used to represent physical quantities as well as understand the rules and conventions that have been developed to manipulate them.

- **variables:** things that can vary, e.g. temperature and time
- **constants:** fixed and unchanging quantities used in certain calculations, e.g. 3.14159
- in principle one could freely choose letters and symbols to represent variables and constants, but it is helpful and choose letters and symbols that have meaning in a particular context. Hence, we
- $x, y, z$ , the end of the alphabet is reserved for variables
- $a, b, c$ , the beginning of the alphabet is used to represent constants
- $\pi, \omega$  and Greek letters below are used frequently used to represent common constant, e.g.  $\pi = 3.14159$

Table 1.1: Uppercase and lowercase letters of the Greek alphabet

Letter	Upper case	Lower case	Letter	Upper case	Lower case
Alpha	A	$\alpha$	Nu	N	$\nu$
Beta	B	$\beta$	Xi	$\Xi$	$\xi$
Gamma	$\Gamma$	$\gamma$	Omicron	O	$o$
Delta	$\Delta$	$\delta$	Pi	$\Pi$	$\pi$
Epsilon	E	$\epsilon$	Rho	P	$\rho$
Zeta	Z	$\zeta$	Sigma	$\Sigma$	$\sigma$
Eta	H	$\eta$	Tau	T	$\tau$
Theta	$\Theta$	$\theta$	Upsilon	Y	$\upsilon$
Iota	i	$\iota$	Phi	$\Phi$	$\phi$
Kappa	K	$\kappa$	Chi	$\Gamma$	$\gamma$
Lambda	$\Gamma$	$\gamma$	Psi	$\Psi$	$\psi$
Mu	M	$\mu$	Omega	$\Omega$	$\omega$

## 1.3 A precise language

- Mathematics is a precise language meaning that a special attention has to be paid to the exact position of any symbol in relation to other.
- Given two symbols  $x$  and  $y$ ,  $xy$  and  $x^y$  and  $x_y$  can mean different things
- $xy$  stands for multiplication,  $x^y$  for superscript and  $x_y$  for subscript

## 1.4 Using symbols

If the letters  $x$  and  $y$  represent two numbers, then:

- their **sum** is written as  $x + y$
- subtracting  $y$  from  $x$  is  $x - y$ , known also as **difference**
- to multiply  $x$  and  $y$  we written as  $x \cdot y$  or also with the multiplication signed omitted as  $xy$ . The quantity is known as **product of x and y**
- multiplication is **associative**, e.g. when we multiply three numbers together,  $x \cdot y \cdot z$ , the order of multiplication does not matter, so  $x \cdot y \cdot z$  is the same as  $z \cdot x \cdot y$  or  $y \cdot z \cdot x$
- division is denoted by  $\frac{x}{y}$  and means that  $x$  is divided by  $y$ . In this expression  $x$ , on the top, is called **numerator** and  $y$ , on the bottom, is called **denominator**
- division by 1 leaves any number unchanged, e.g.  $\frac{x}{1} = x$  and division by 0 is not allowed

Equal sign

- the equal sign  $=$  is used in **equations**, e.g.  $x - 5 = 0$  or  $5x = 1$
- the equal sign  $=$  can be also used in **formulae**. Physical quantities are related through a formula in many fields, e.g. the formula  $A = \pi r^2$  relates circle area  $A$  to its radius  $r$  and the formula  $s = \frac{d}{t}$  defines speed as distance  $d$  divided by time  $t$
- the equal sign  $=$  is also used in identities, expressions true for all values of the variable, e.g.  $(x - 1)(x + 1) = (x^2 - 1)$
- opposite to the equal sign is “is not equal to” sign  $\neq$ , e.g. we can write  $1 + 2 \neq 4$

**Sigma and Product notation**

- the  $\Sigma$  notation, read as **Sigma notation**, provides a convenient way of writing long sums, e.g. the sum of  $x_1 + x_2 + x_3 + \dots + x_{20}$  is written as  $\sum_{i=1}^{i=20} x_i$
- the  $\Pi$  notation, read as **Product notation**, provides a convenient way of writing long products, e.g.  $x_1 \cdot x_2 \cdot x_3 \cdot \dots \cdot x_{20}$  is written as  $\prod_{i=1}^{i=20} x_i$

## 1.5 Inequalities

Given any two real numbers  $a$  and  $b$  there are three mutually exclusive possibilities:

- $a > b$ , meaning that  $a$  is greater than  $b$
- $a < b$ , meaning that  $a$  is less than  $b$
- $a = b$ , meaning that  $a$  is equal to  $b$

Strict and weak

- inequality in  $a > b$  and  $a < b$  is **strict**
- as oppose to **weak** inequality denoted as  $a \geq b$  or  $a \leq b$

Some useful relations are:

- if  $a > b$  and  $b > c$  then  $a > c$
- if  $a > b$  then  $a + c > b$  for any  $c$
- if  $a > b$  then  $ac > bc$  for any positive  $c$
- if  $a > b$  then  $ac < bc$  for any negative  $c$

## 1.6 Indices and powers

- **Indices**, also known as **powers** are convenient when we multiply a number by itself several times
- e.g.  $5 \cdot 5 \cdot 5$  is written as  $5^3$  and  $4 \cdot 4 \cdot 4 \cdot 4 \cdot 4$  is written as  $4^5$
- in the expression  $x^y$ ,  $x$  is called the *base* and  $y$  is called *index* or *power*

The laws of indices state:

- $a^m \cdot a^n = a^{m+n}$
- $\frac{a^m}{a^n} = a^{m-n}$
- $(a^m)^n = a^{m \cdot n}$

Rules derived from the laws of indices:

- $a^0 = 1$
- $a^1 = a$

Negative and fractional indices:

- $a^{-m} = \frac{1}{a^m}$  e.g.  $5^{-2} = \frac{1}{5^2} = \frac{1}{25}$  for negative indices
- e.g.  $4^{\frac{1}{2}} = \sqrt{4}$  or  $8^{\frac{1}{3}} = \sqrt[3]{8}$  for fractional indices

## 1.7 Exercises: notations

**Exercise 1.1.** Classify numbers as natural, integers or real. If real, specify if they are rational or irrational.

- $\frac{1}{3}$
- 2
- $\sqrt{4}$
- 2.3
- $\pi$
- $\sqrt{5}$
- 7
- 0
- 0.25

**Exercise 1.2.** Classify below descriptors as variables or constants. Do you know the letters or symbols commonly used to represent these?

- speed of light in vacuum
- mass of an apple
- volume of an apple

- d) concentration of vitamin C in an apple
- e) distance from Stockholm central station to Uppsala central station
- f) time on the train to travel between the above stations
- g) electron charge

**Exercise 1.3.** Write out explicitly what is meant by the following:

- a)  $\sum_{i=1}^{i=6} k_i$
- b)  $\prod_{i=1}^{i=6} k_i$
- c)  $\sum_{i=1}^{i=6} i^k$
- d)  $\prod_{i=1}^{i=3} i^k$
- e)  $\sum_{i=1}^n i$
- f)  $\sum_{i=1}^{i=4} (i+1)^k$
- g)  $\prod_{i=1}^{i=4} (k+1)^i$
- h)  $\prod_{i=0}^n i$

**Exercise 1.4.** Use Sigma or Product notation to represent the long sums and products below:

- a)  $1 + 2 + 3 + 4 + 5 + 6$
- b)  $2^2 + 3^2 + 4^2 + 5^2$
- c)  $4 \cdot 5 \cdot 6 \cdot 7 \cdot 8$
- d)  $1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{5} + \dots + \frac{1}{n}$
- e)  $2 - 2^2 + 2^3 - 2^4 + \dots + 2^n$
- f)  $3 + 6 + 9 + 12 + \dots + 60$
- g)  $3x + 6x^2 + 9x^3 + 12x^4 + \dots + 60x^{20}$
- h)  $3x \cdot 6x^2 \cdot 9x^3 \cdot 12x^4 \cdot \dots \cdot 60x^{20}$

## Answers to selected exercises (notations)

Exr. 1.1

- a) real, rational
- b) natural and integers, integers include natural numbers
- c)  $\sqrt{4} = 2$  so it is a natural number and/subset of integers
- d) real number, rational as it could be written as  $\frac{23}{10}$
- e) real number, irrational as it cannot be explained by a simple fraction
- f) real number, irrational as it cannot be explained by a simple fraction
- g) integer, non a natural number as these do not include negative numbers
- h) natural number, although there is some argument about it as some define natural numbers as positive integers starting from 1, 2 etc. while others include 0.
- i) real, rational number, could be written as  $\frac{25}{100}$

## Exr. 1.2

- a) constant, speed of light in vacuum is a constant, denoted  $c$  with  $c = 299792458 \frac{m}{s}$
- b) variable, mass of an apple is a variable, different for different apple sizes, for instance 138 grams, denoted as  $m = 100g$
- c) variable, like mass volume can be different from apple to apple, denoted as  $V$ , e.g.  $V = 200cm^3$
- d) variable, like volume and mass can vary, denoted as  $\rho_i$  and defined as  $\rho_i = \frac{m}{V}$ . So given 6.3 milligrams of vitamin C in our example apple, we have  $\rho_i = \frac{0.0063}{2} \frac{g}{cm^3} = 0.000315 \frac{g}{cm^3}$  concentration of vitamin D
- e) constant, the distance between Stockholm and Uppsala is fixed; it could be a variable though if we were to consider an experiment on a very long time scale; distance is often denoted in physics as  $d$
- f) variable, time on the train to travel between the stations varies, often denoted as  $t$  with speed being calculated as  $s = \frac{d}{t}$
- g) constant, electron charge is  $e = 1.60217663 \cdot 10^{-19}C$

## Exr. 1.3

- a)  $\sum_{i=1}^{i=6} k_i = k_1 + k_2 + k_3 + k_4 + k_5 + k_6$
- b)  $\prod_{i=1}^{i=6} k_i = k_1 \cdot k_2 \cdot k_3 \cdot k_4 \cdot k_5 \cdot k_6$
- c)  $\sum_{i=1}^{i=3} i^k = 1^k + 2^k + 3^k$
- d)  $\prod_{i=1}^{i=3} i^k = 1^k \cdot 2^k \cdot 3^k$
- e)  $\sum_{i=1}^n i = 1 + 2 + 3 + \dots + n$  we are using dots (...) to represent all the number until  $n$ . Here, thanks to Gauss we can also write  $\sum_{i=1}^n i = \frac{n(n+1)}{2}$ , i.e. Gauss formula for sum of first  $n$  natural numbers

## Exr. 1.4

- a)  $1 + 2 + 3 + 4 + 5 + 6 = \sum_{k=1}^6 k$
- b)  $2^2 + 3^2 + 4^2 + 5^2 = \sum_{x=2}^5 x^2$
- c)  $4 \cdot 5 \cdot 6 \cdot 7 \cdot 8 = \prod_{x=4}^8 x$
- d)  $1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{5} + \dots + \frac{1}{n} = \sum_{n=1}^n \frac{1}{n}$

# Chapter 2

## Sets

### Aims

- to introduce sets and basic operations on sets

### Learning outcomes

- to be able to explain what a set is
- to be able to construct new sets from given sets using the basic set operations
- to be able to use Venn diagrams to show all possible logical relations between two and three sets

## 2.1 Definitions

- **set**: a well-defined collection of distinct objects, e.g.  $S = \{2, 4, 6\}$
- **elements**: the objects that make up the set are also known as **elements** of the set
- if  $x$  is an element of  $S$ , we say that  $x$  belongs to  $S$  and write  $x \in S$  and if  $x$  is not an element of  $S$  we say that  $x$  does not belong to  $S$  and write  $x \notin S$
- a set may contain **finitely** many or **infinitely** many elements
- **subset**,  $\subseteq$ : if every element of set  $A$  is also in  $B$ , then  $A$  is said to be a subset of  $B$ , written as  $A \subseteq B$  and pronounced  $A$  is contained in  $B$ , e.g.  $A \subseteq B$ , when  $A = \{2, 4, 6\}$  and  $B = \{2, 4, 6, 8, 10\}$ . Every set is a subset of itself.
- **superset**: for our sets  $A$  and  $B$  we can also say that  $B$  is a **superset** of  $A$  and write  $B \supset A$
- **cardinality**: the number of elements within a set  $S$ , denoted as  $|S|$
- **empty set**,  $\emptyset$ : is a unique set with no members, denoted by  $E = \emptyset$  or  $E = \{\}$ . The empty set is a subset of every set.


## 2.2 Basic set operations

- **union of two sets**,  $\cup$  : two sets can be “added” together, the union of A and B, written as  $A \cup B$ , e.g.  $\{1, 2\} \cup \{2, 3\} = \{1, 2, 3\}$  or  $\{1, 2, 3\} \cup \{1, 4, 5, 6\} = \{1, 2, 3, 4, 5, 6\}$
- **intersection of two sets**,  $\cap$ : a new set can be constructed by taking members of two sets that are “in common”, written as  $A \cap B$ , e.g.  $\{1, 2, 3, 4, 5, 6\} \cap \{2, 3, 7\} = \{2, 3\}$  or  $\{1, 2, 3\} \cap \{7\} = \emptyset$
- **complement of a set**,  $A'$ ,  $A^c$ : are the elements not in A
- **difference of two sets**,  $\setminus$ : two sets can be “subtracted”, denoted by  $A \setminus B$ , by taking all elements that are members of A but are not members of B, e.g.  $\{1, 2, 3, 4\} \setminus \{1, 3\} = \{2, 4\}$ . This is also in other words a relative complement of A with respect to B.
- **partition of a set**: a partition of a set S is a set of nonempty subset of S, such that every element x in S is in exactly one of these subsets. That is, the subset are pairwise *disjoint*, meaning no two sets of the partition contain elements in common, and the union of all the subset of the partition is S, e.g. Set  $\{1, 2, 3\}$  has five partitions: i)  $\{1\}, \{2\}, \{3\}$ , ii)  $\{1, 2\}, \{3\}$ , iii)  $\{1, 3\}, \{2\}$ , iv)  $\{1\}, \{2, 3\}$  and v)  $\{1, 2, 3\}$

## 2.3 Venn diagrams


Venn diagram is a diagram that shows all possible logical relations between a finite collection of different sets. A Venn diagrams shows elements as points in the plane, and sets as regions inside closed curves. A Venn diagram consists of multiple overlapping closed curves, usually circles, each representing a set.

E.g. given  $A = \{1, 2, 5\}$  and  $B = \{1, 6\}$  Venn diagram of A and B:




And given  $A = \{1, 2, 5\}$ ,  $B = \{1, 6\}$  and  $C = \{4, 7\}$  Venn diagram of A, B and C:





And given  $A = \{1, 2, 3, 4, 5, 6\}$  and  $B = \{2, 4, 6\}$  Venn diagram of  $A$  and  $B$ :



## 2.4 Exercises: sets

**Exercise 2.1.** Given set  $S = \{1, 2, 3, 4, 5, 6\}$ :

- what is the subset  $T$  of  $S$  consisting of its even elements?
- what is the complement  $T^c$  of  $T$ ?
- what is the subset  $U$  of  $S$  containing of the prime numbers in  $S$ ?
- What is the intersection  $T \cap U$ ?
- What is the union of  $T \cup U$ ?
- What is the set difference  $U \setminus T$ ?

**Exercise 2.2.** Given set  $A = \{cat, elephant, dog, turtle, goldfish, hamster, parrot, tiger, guineapig, lion\}$

- what is the subset  $D$  of  $A$  consisting of domesticated animals?
- what is the subset  $C$  of  $A$  consisting of Felidae (cat) family?

- c) what is the intersection of  $D$  and  $C$ ?
- d) what is the complement of  $D$ ,  $D^c$ ?
- e) what is the union of  $D$  and  $C$ ?
- f) what is the set difference of  $A \setminus C$ ?
- g) can you draw Venn diagram showing relationship between  $D$  and  $C$ ?

## Answers to selected exercises (sets)

Exr. 2.1

- a)  $T = \{2, 4, 6\}$
- b)  $T^c = \{1, 3, 5\}$ , i.e.  $T^c$  contains all the elements of  $S$  not in  $T$
- c)  $U = \{2, 3, 5\}$ , the primes in  $S$
- d)  $T \cap U = \{2\}$ , common elements of  $T$  and  $U$ , i.e. the even and prime numbers
- e)  $T \cup U = \{2, 3, 4, 5, 6\}$
- f)  $U \setminus T = \{3, 5\}$ , consisting of the elements of  $U$  that are not in  $T$

# Chapter 3

## Functions

### Aims

- to revisit the concept of a function

### Learning outcomes

- to be able to explain what function, function domain and function range are
- to be able to identify input, output, argument, independent variable, dependent variable
- to be able to evaluate function for a given value and plot the function

### 3.1 Definitions

- A **function**,  $f(x)$ , can be viewed as a rule that relates input  $x$  to an output  $f(x)$
- In order for a rule to be a function it must produce a single output for any given input
- Input  $x$  is also known as **argument** of the function
- **Domain of a function**: the set of all values that the function “maps”
- **Range**: the set of all values that the function maps into

### Many names are used interchangeably

Functions have been around for a while and there are many alternative names and writing conventions are being used. Common terms worth knowing:

### 3.2 Evaluating function

To evaluate a function is to replace (substitute) its variable with a given number or expression. E.g. given a rule (function) that maps temperature measurements

from Celsius to Fahrenheit scale:

$$f(x) = 1.8x + 32$$

where  $x$  is temperature measurements in Celsius and  $f(x)$  is the associated value in Fahrenheit, we can find for a given temperature in Celsius corresponding temperature in Fahrenheit. Let's say we measure 10 Celsius degrees one autumn day in Uppsala and we want to share this information with a friend in USA. We can find the equivalent temperature in Fahrenheit by evaluating our function at 10,  $f(10)$ , giving us

$$f(10) = 1.8 \cdot 10 + 32 = 50$$

### 3.3 Plotting function

Function graphs are a convenient way of showing functions, by looking at the graph it is easier to notice function's properties, e.g. for which input values functions yields positive outcomes or whether the function is increasing or decreasing. To graph a function, one can start by evaluating function at different values for the argument  $x$  obtaining  $f(x)$ , plotting the points by plotting the pairs  $(x, f(x))$  and connecting the dots. E.g. evaluating our above temperature rule at -20, -10, 0, 10, 20, 30 Celsius degrees results in:

x (Celsius degrees)	evaluates	f(x) (Fahrenheit degrees)
-20	$f(-20) = 1.8 \cdot (-20) + 32$	-4
-10	$f(-10) = 1.8 \cdot (-10) + 32$	14
0	$f(0) = 1.8 \cdot (0) + 32$	32
10	$f(10) = 1.8 \cdot (10) + 32$	50
20	$f(20) = 1.8 \cdot (20) + 32$	68
30	$f(30) = 1.8 \cdot (30) + 32$	86


### 3.4 Standard classes of functions

**Algebraic function:** functions that can be expressed as the solution of a polynomial equation with integer coefficients, e.g.

- constant function  $f(x) = a$
- identity function  $f(x) = x$
- linear function  $f(x) = ax + b$
- quadratic function  $f(x) = a + bx + cx^2$
- cubic function  $f(x) = a + bx + cx^2 + dx^3$

**Transcendental functions:** functions that are not algebraic, e.g.

- exponential function  $f(x) = e^x$

Figure 3.1: Graph of  $f(x)$  for the temperature rule

- logarithmic function  $f(x) = \log(x)$
- trigonometric function  $f(x) = -3\sin(2x)$

### 3.5 Piecewise functions

A function can be in pieces, i.e. we can create functions that behave differently based on the input  $x$  value. They are useful to describe situations in which a rule changes as the input value crosses certain “boundaries”. E.g. a function value could be fixed in a given range and equal to the input value (identity function) for input values outside this range

$$f(x) = \begin{cases} 2 & \text{if } x \leq 1 \\ x & \text{if } x > 1 \end{cases} \quad (3.1)$$

The function can be split in many pieces, e.g. the personal training fee in SEK may depend whether the personal trainer is hired for an hour, two hours or three or more hours:

$$f(h) = \begin{cases} 500 & \text{if } h \leq 1 \\ 750 & \text{if } 1 < h \leq 2 \\ 500 + 250 \cdot h & \text{if } h > 2 \end{cases} \quad (3.2)$$




Figure 3.2: Examples of the standard classess of functions




Figure 3.3: Examples of piece-wise functions

### 3.6 Exercises: functions

**Exercise 3.1.** Given the function for the personal trainer costs:

$$f(h) = \begin{cases} 500 & \text{if } h \leq 1 \\ 750 & \text{if } 1 < h \leq 2 \\ 500 + 250 \cdot h & \text{if } h > 2 \end{cases} \quad (3.3)$$

How much would you pay

- a) for a 4-hours session? Evaluate function  $f(h)$  for value 4.
- b) for a 2-hour session? Evaluate function  $f(h)$  for value 2.

**Exercise 3.2.** A museum charges 50 SEK per person for a guided tour with a group of 1 to 9 people or a fixed \$500 SEK fee for a group of 10 or more people. Write a function relating the number of people  $n$  to the cost  $C$ .

**Exercise 3.3.** Given function

$$f(x) = \begin{cases} x^2 & \text{if } x \leq 1 \\ 3 & \text{if } 1 < x \leq 2 \\ x & \text{if } x > 2 \end{cases} \quad (3.4)$$

- a) sketch a graph of a function for  $x$  range  $(-4, 4)$
- b) evaluate function at  $f(1)$
- c) evaluate function at  $f(4)$

### Answers to selected exercises (functions)

Exr. 3.1

- a)  $f(4) = 500 + 250 \cdot 4 = 1500$
- b)  $f(2) = 750$  as  $h \leq 2$  means less or equal to 2, that is including 2





## Part II

# Introduction to R, R Studio and R markdown



## Chapter 4

# R

bla bla bla



## Chapter 5

# R Studio

bla bla bla



## Chapter 6

# R Markdown

bla bla bla





Part III

Probability



## Chapter 7

# Probability: reasoning under uncertainty

### Learning outcomes

- understand the concept of probability
- manipulate probabilities by their rules
- assign probabilities in very simple cases

### 7.1 Introduction

Some things are more likely to occur than others. Compare:

- the chance of the sun rising tomorrow with the chance that no-one is infected with COVID-19 tomorrow
- the chance of a cold dark winter in Stockholm with the chance of no rainy days over the summer months in Stockholm

We intuitively believe that the chance of sun rising or dark winter occurring are enormously higher than COVID-19 disappearing over night or having no rain over the entire summer. **Probability** gives us a scale for measuring the likeliness of events to occur. **Probability rules** enable us to reason about uncertain events. The probability rules are expressed in terms of sets, a well-defined collection of distinct objects.

### 7.2 Basic set definitions

- **set**: a well-defined collection of distinct objects, e.g.  $A = \{2, 4, 6\}$
- **subset**,  $\subseteq$ : if every element of set  $A$  is also in  $B$ , then  $A$  is said to be a subset of  $B$ , written as  $A \subseteq B$  and pronounced  $A$  is contained in  $B$ ,

e.g.  $A \subseteq B$ , when  $B = \{2, 4, 6, 8, 10\}$ . Every set is a subset of itself.

- **empty set**,  $\emptyset$ : is a unique set with no members, denoted by  $E = \emptyset$  or  $E = \{\}$ . The empty set is a subset of every set.

### 7.3 Basic set operations

- **union of two sets**,  $\cup$ : two sets can be “added” together, the union of A and B, written as  $A \cup B$ , e.g.  $\{1, 2\} \cup \{2, 3\} = \{1, 2, 3\}$  or  $\{1, 2, 3\} \cup \{1, 4, 5, 6\} = \{1, 2, 3, 4, 5, 6\}$
- **intersection of two sets**,  $\cap$ : a new set can be constructed by taking members of two sets that are “in common”, written as  $A \cap B$ , e.g.  $\{1, 2, 3, 4, 5, 6\} \cap \{2, 3, 7\} = \{2, 3\}$  or  $\{1, 2, 3\} \cap \{7\} = \emptyset$
- **complement of a set**,  $A^c$ ,  $A^C$ : are the elements not in A
- **difference of two sets**,  $\setminus$ : two sets can be “subtracted”, denoted by  $A \setminus B$ , by taking all elements that are members of A but are not members of B, e.g.  $\{1, 2, 3, 4\} \setminus \{1, 3\} = \{2, 4\}$ . This is also in other words a relative complement of A with respect to B.
- **partition of a set**: a partition of a set S is a set of nonempty subset of S, such that every element x in S is in exactly one of these subsets. That is, the subset are pairwise *disjoint*, meaning no two sets of the partition contain elements in common, and the union of all the subset of the partition is S, e.g. Set  $\{1, 2, 3\}$  has five partitions: i)  $\{1\}, \{2\}, \{3\}$ , ii)  $\{1, 2\}, \{3\}$ , iii)  $\{1, 3\}, \{2\}$ , iv)  $\{1\}, \{2, 3\}$  and v)  $\{1, 2, 3\}$

### 7.4 Exercises

**Exercise 7.1.** Here is my exercise.

### 7.5 Answers to exercises

## Chapter 8

# Probability: random variables

### Learning outcomes

- understand the concept of random discrete and continuous variables
- to be able to use probability density/mass functions and cumulative distribution functions and to understand the relationship between them
- describe properties of binomial, geometric, Poisson, uniform, exponential and normal distributions and identify which distributions to use in practical problems

### 8.1 Random variables

The outcome of a random experiment can be described by a random variable.

Example random variables:

- The weight of a random newborn baby
- The smoking status of a random mother
- The hemoglobin concentration in blood
- The number of mutations in a gene
- BMI of a random man
- Weight status of a random man (underweight, normal weight, overweight, obese)
- The result of throwing a die

Whenever chance is involved in the outcome of an experiment the outcome is a random variable.

A random variable is usually denoted by a capital letter,  $X, Y, Z, \dots$ . Values collected in an experiment are observations of the random variable, usually

denoted by lowercase letters  $x, y, z, \dots$

A random variable can not be predicted exactly, but the probability of all possible outcomes can be described.

The population is the collection of all possible observations of the random variable. Note, the population is not always countable.

A sample is a subset of the population.

## 8.2 Discrete random variables

A discrete random variable can be described by its *probability mass function*.

->

->

## Chapter 9

# Summarising and visualising data





## Chapter 10

# Linear regression

10.1 Simple regression

10.2 Multiple regression