

## Derivatives Review:

$$1. (\cos x)' = -\sin x$$

$$2. (e^{2x})' = e^{2x} \cdot 2$$

$$3. (\arcsin x)' = \frac{1}{\sqrt{1-x^2}}$$

$$4. (\sec 3x)' = \tan(3x) \sec(3x) \cdot 3$$

$$5. \left(-\frac{1}{x}\right)' = \frac{1}{x^2}$$

$$6. (\arctan x)' = \frac{1}{1+x^2}$$


$$7. (\ln(5x))' = \frac{1}{5x} \cdot 5$$

$$8. (\cot(2x))' = -\operatorname{csc}^2(2x) \cdot 2$$

$$9. (\sqrt{x})' = \frac{1}{2\sqrt{x}}$$

SECTION 3.9: RELATED RATES – DAY 2  
SECTION 3.10 TANGENT LINE APPROXIMATION INTRO

1. A Ferris wheel with a radius of 13 meters is rotating at a rate of one revolution every three minutes. How fast is a rider rising when her seat is 18 meters above the ground? (Assume the wheel is tangent to the ground at the bottom.) *Hint: Label useful things in the diagram sketch.*



- (a) In terms of the labels given in the picture and calculus-type language:

- What do we KNOW? (Hint: how many radians in one revolution?)

$$r = 13 \text{ meters} \quad \frac{d\theta}{dt} = \frac{2\pi}{3} \frac{\text{rad}}{\text{min}}$$

- What do we WANT?

$$\frac{dh}{dt} \text{ when } h = 18 \text{ meters}$$

- (b) Determine an equation that relates the variables in your WANT and KNOW.

$$h = 13 + y = 13 + 13 \cdot \sin \theta$$

$$h = 13 + 13 \sin \theta$$

$$\sin \theta = \frac{y}{13}$$

$$y = 13 \cdot \sin \theta$$

- (c) Solve the related rates problem.

(Hint: use what you know about right-triangle trigonometry! You don't actually need to know the angle from horizontal she's at when she's 18 feet above the ground.)

$$\frac{dh}{dt} = \frac{d}{dt} (13 + 13 \sin \theta) = 0 + 13 \cdot \cos \theta \cdot \frac{d\theta}{dt}$$

$$\frac{dh}{dt} = 13 \cos \theta \cdot \frac{d\theta}{dt} = 13 \cdot \cos \theta \cdot \frac{2\pi}{3}$$

$$\cos \theta = \frac{x}{13}$$


$$x^2 + y^2 = 13^2$$

$$y = 5$$

$$x^2 + 25 = 169$$

$$x^2 = 144$$

$$x = 12$$



$$\cos \theta = \frac{x}{13} = \frac{12}{13}$$

$$\frac{dh}{dt} = 13 \cdot \frac{12}{13} \cdot \frac{2\pi}{3} = 8\pi \text{ m/min}$$

2. Consider the function  $f(x) = x^3$ .

(a) At the point  $x = 2$ , what is  $f(x)$ ?

$$f(2) = \underline{8}$$

(b) Let  $L(x)$  be the function that is the tangent line to  $f(x)$  at  $x = 2$ . This tangent line is sometimes called the *linearization* of  $f(x)$  at  $x = 2$ . Finish the equation (you will need to show some work).

$$L(x) = f'(a)(x-a) + f(a) \quad L(x) = 12(x-2) + 8$$

$$a = 2$$

$$f(a) = 8$$

$$f'(x) = 3x^2 \quad f'(2) = 12 \quad L(x) = \underline{12(x-2) + 8}$$

(c) Observe that the value  $x = 2.1 = 2 + \frac{1}{10}$  is very close to  $x = 2$ . Evaluate  $L(x)$  at  $x = 2.1$ . Do not use a calculator until your very last step (that is you can get a decimal approximation of a fraction, but you should compute the fraction by hand).

$$(2.1)^3 \approx L(2.1) = 12(2.1-2) + 8 = 12 \cdot 0.1 + 8 =$$

$$= \frac{12}{10} + 8 = \frac{6}{5} + 8 = \frac{46}{5} \approx 9.2$$

$$L(2.1) \text{ as a fraction } \underline{\frac{46}{5}} \quad L(2.1) \text{ as a decimal approximation. } \underline{9.2}$$

(d) Use a calculator or a computer to evaluate  $f(2.1)$ .  $f(2.1) = \underline{9.261}$

(e) What is the error if you use  $L(2.1)$  to approximate  $f(2.1)$ ? (That is, what is the difference between the two quantities?) What is the percent error, calculated as (approx value - actual value)/(actual value)?

$$\text{error} = |9.2 - 9.261| = 0.061$$

(f) Draw a rough sketch of  $f(x)$  and  $L(x)$ , and use the picture and your computations to explain, in a sentence or two, why using  $L(2.1)$  to approximate the cube of 2.1 is a reasonable thing to do.

