

1. Fill in the table below.

Derivatives of Trigonometric Functions:

$$\bullet \frac{d}{dx}(\sin x) = \underline{\cos(x)}$$

$$\bullet \frac{d}{dx}(\cos x) = \underline{-\sin(x)}$$

$$\bullet \frac{d}{dx}(\tan x) = \underline{\sec^2(x)}$$

$$\bullet \frac{d}{dx}(\csc x) = \underline{-\cot(x) \cdot \csc(x)}$$

$$\bullet \frac{d}{dx}(\sec x) = \underline{\tan(x) \cdot \sec(x)}$$

$$\bullet \frac{d}{dx}(\cot x) = \underline{-\csc^2(x)}$$

2. Find the derivative of $y = \frac{\sec x}{1 - x \tan x}$. Quotient Rule

$$y' = \frac{f' \cdot g - g' \cdot f}{g^2} = \frac{(\sec(x))' \cdot (1 - x \tan(x)) - (1 - x \tan(x))' \cdot \sec(x)}{(1 - x \tan(x))^2}$$

$$= \frac{\tan(x) \cdot \sec(x) \cdot (1 - x \tan(x)) + (\tan(x) + x \cdot \sec^2(x)) \cdot \sec(x)}{(1 - x \tan(x))^2}$$

$(\sec(x))' = \tan(x) \cdot \sec(x)$

$(1 - x \tan(x))' = (1)' - (x \tan(x))' = -(x \tan(x))' = -(\tan(x) + x \sec^2(x))$

3. If $f(\theta) = e^\theta \sin(\theta)$, find $f''(\theta)$. Simplify your answers.

$$f'(\theta) = e^\theta \cdot \sin(\theta) + \cos(\theta) \cdot e^\theta$$

$$= e^\theta (\sin(\theta) + \cos(\theta))$$

$$f''(\theta) = (f'(\theta))'$$

theta θ

$$f''(\theta) = (f'(\theta))' = e^\theta \cdot (\sin(\theta) + \cos(\theta)) + (\cos(\theta) - \sin(\theta)) \cdot e^\theta$$

4. Find $\frac{d}{dt}(t \sin t \cos t)$.

$$\left(\underbrace{(t \cdot \cos t)}_f \cdot \underbrace{(\sin t)}_g \right) \text{ or } \left(\underbrace{\left(\frac{t}{f} \right)}_f \cdot \underbrace{(\sin(t) \cdot \cos(t))}_g \right)$$

Product Rule:

$$\left(\underbrace{t \cdot \sin(t)}_f \cdot \underbrace{\cos(t)}_g \right)' = f' \cdot g + g' \cdot f = (t \cdot \sin t)' \cdot \cos(t) + (\cos t)' \cdot t \cdot \sin(t)$$

$$= (\sin(t) + t \cos(t)) \cdot \cos(t) - t \cdot \sin^2(t)$$

$(\cos(t))' = -\sin(t)$

$(t \cdot \sin(t))' = (t)' \cdot \sin(t) + (\sin(t))' \cdot t = 1 \cdot \sin(t) + \cos(t) \cdot t$

5. An elastic band is hung on a hook and a mass is hung on the lower end of the band. When the mass is pulled down 2 cm past its rest position and then released, it vibrates vertically. The equation of motion is

$$s = 2 \cos t + 3 \sin t, \text{ for } t \geq 0,$$

where s is measured in centimeters and t is measured in seconds. (We are taking the positive direction to be downward.)

- (a) Why might you expect to use sines and cosines to model this particular problem?

displacement velocity acceleration

- (b) Find $s(0)$, $s'(0)$, and $s''(0)$ including units.

$$s(0) = 2 \cos(0) + 3 \sin(0) = 2 \cdot 1 + 3 \cdot 0 = 2 \text{ (cm)}$$

$$s'(t) = -2 \sin(t) + 3 \cos(t)$$

$$s'(0) = -2 \sin(0) + 3 \cos(0) = -2 \cdot 0 + 3 \cdot 1 = 3 \text{ (cm/sec)}$$

$$s''(t) = -2 \cos(t) - 3 \sin(t) \quad s''(0) = -2 \text{ (cm/sec}^2\text{)}$$

- (c) What do the numbers from part (a) indicate about the mass in the context of the problem?

$s(0)$ is an initial displacement of the mass

$s'(0)$ is an initial velocity of the mass


$s''(0)$ is an initial acceleration of the mass

6. A 12 foot ladder rests against a wall. Let θ be the angle between the ladder and the wall and let x be the distance from the base of the ladder and the wall.

- (a) Compute x as a function of θ . (Drawing a picture will help.)

$$x = x(\theta)$$

$$\sin(\theta) = \frac{x}{12} \Rightarrow x = 12 \sin(\theta)$$



- (b) How fast does x change with respect to θ when $\theta = \pi/6$? (Get an exact answer and a decimal approximation.)

$$\left. \frac{dx}{d\theta} \right|_{\theta = \frac{\pi}{6}} = x'(\frac{\pi}{6}) = 12 \cos(\theta) \Big|_{\theta = \frac{\pi}{6}} = \frac{12\sqrt{3}}{2} = 6\sqrt{3} \text{ (foot/sec)}$$

- (c) Interpret your answer from part (b) in the context of the problem. (Units will help you here.)

The displacement of the ladder (distance from the wall) is increasing at the rate of $6\sqrt{3}$ foot/sec when $\theta = \frac{\pi}{6}$.

- (d) Determine how far the ladder is from the wall when $\theta = \pi/6$.

$$x(\theta) \text{ when } \theta = \frac{\pi}{6} ?$$

$$x\left(\frac{\pi}{6}\right) = 12 \cdot \sin\left(\frac{\pi}{6}\right) = 6 \text{ foot}$$