

THEORETICAL PART:**Definition (y-Axis Symmetry):**

The graph of a function f has **y-axis symmetry**, or is **symmetric with respect to the y-axis**, if $f(-x) = f(x)$ for all x in the domain of f . Such functions are called **even functions**.

Definition (Origin Symmetry):

The graph of a function f has **origin symmetry**, or is **symmetric with respect to the origin**, if $f(-x) = -f(x)$ for all x in the domain of f . Such functions are called **odd functions**.

Definition (Symmetry of Equations):

We say an equation in x and y is **symmetric with respect to**

1. the **y-axis** if replacing x with $-x$ results in an equivalent equation;
2. the **x-axis** if replacing y with $-y$ results in an equivalent equation;
3. the **origin** if replacing x with $-x$ and y with $-y$ results in an equivalent equation.

Definition (Increasing, decreasing, and Constant):

We say that a function f is

1. **increasing on an interval** if for any x_1 and x_2 in the interval with $x_1 < x_2$, it is the case that $f(x_1) < f(x_2)$;
2. **decreasing on an interval** if for any x_1 and x_2 in the interval with $x_1 < x_2$, it is the case that $f(x_1) > f(x_2)$;
3. **constant on an interval** if for any x_1 and x_2 in the interval, it is the case that $f(x_1) = f(x_2)$.

Definition (Local Extrema):

A function f has a **local maximum at c** if there is an open interval (a, b) containing c for which $f(x) \leq f(c)$ for all x in (a, b) . In this case we say $f(c)$ is the **local maximum value** of f .

Similarly, f has a **local minimum at c** if there is an open interval (a, b) containing c for which $f(x) \geq f(c)$ for all x in (a, b) , and in this case we say $f(c)$ is the **local minimum value** of f . The local maxima and minima of a function are collectively referred to as **local extrema**.

Definition (Average Rate of Change):

Given function f defined on an interval $[a, b]$, $a \neq b$, the **average rate of change** of f over $[a, b]$ is

$$\frac{f(b) - f(a)}{b - a}.$$

If $y = f(x)$, then any of the following expressions may be used to represent the average rate of change of f over $[a, b]$:

$$\frac{\Delta y}{\Delta x} = \frac{\Delta f}{\Delta x} = \frac{f(b) - f(a)}{b - a}.$$

The average rate of change of f over $[a, b]$ represents the slope of the **secant line** drawn between the points $(a, f(a))$ and $(b, f(b))$ on the graph of f .


PRACTICAL PART:

1. Sketch the graphs of the following relations, making use of symmetry:


(a) $f(x) = \frac{1}{x^2}$

(b) $g(x) = x^3 - x$


(b) $x = y^2$



2. Determine the open intervals of monotonicity of the function $f(x) = (x - 2)^2 - 1$.




3. The water level of a certain river varied over the course of a year as follows. In January, the level was 13 feet. From that level, the water increased linearly to a level of 18 feet in May. The water remained constant at that level until July, at which point it began to decrease linearly to a final level of 11 feet in December. Graph the water level as a function of time and determine the open intervals of monotonicity.



4. For the given graph of the function f below determine:

- locations and types of the local extrema of f ;
- the values of the local extrema of f .



5. Given the function $f(x) = 3x^2 - 5x + 2$, determine the average rate of change over each of the following intervals:

- $[1, 3]$
- $[-2, 2]$
- $[c, c + h], h \neq 0$