# Analysis of Gamma Ray Spectra from Reference Isotopes with Multi-Channel Analyzers


Oliver Kirkpatrick School of Engineering, RMIT University Email: s3725341@student.rmit.edu.au

Abstract-stuff


#### I. RESULTS AND DISCUSSION

#### A. Calibration Process


# iiiiiii HEAD




## (a) Recording 1, hh:mm:ss, T+ minutes



(b) Recording 2, hh:mm:ss, T+ minutes



(c) Recording 3, hh:mm:ss, T+ minutes



(d) <sup>137</sup>Cs Gamma Ray Spectra



# B. Background Radiation




Fig. 2: Background radiation profile.

## C. Analysis of Unknown Sample




TABLE I: Total counts and peak energies with with increasing time since first measurement.

Sample Number	T+ (s)	<b>Total Counts</b>	Peak 1 Energy	Peak 2 Energy	Peak 3 Energy	Peak 4 Energy
1	0s	68493	401.24 keV	815.92 keV	1096.59 keV	1296.46 keV
2	420s	63755	400.52 keV	799.32 keV	1106.59 keV	1295.25 keV
3	819s	59850	401.25 keV	796.80 keV	1099.32 keV	1293.16 keV
4	1199s	56863	400.35 keV	759.04 keV	1097.07 keV	1295.73 keV
5	1559s	53111	401.94 keV	805.54 keV	1104.55 keV	1296.78 keV
6	1936s	50483	400.41 keV	812.17 keV	1085.92 keV	1292.84 keV

		4

#### APPENDIX




Fig. 3: Raw count data for each recording of gamma rays from the unknown isotope.