

# PROBABILITY THEORY

*Introduction to Probability, Statistics and Random Processes*

Hossein Pishro-Nik

## Contents

<b>1</b>	<b>Chapter 1</b>	<b>1</b>
1.1	Set operations . . . . .	1
1.1.1	Cardinality and countable sets . . . . .	2
1.1.2	Functions . . . . .	4
1.2	Random experiments . . . . .	4
1.2.1	Probability . . . . .	5
1.3	Conditional probability . . . . .	7
1.3.1	Independence . . . . .	11
1.3.2	Law of Total Probability . . . . .	12
1.3.3	Conditional Independence . . . . .	14
<b>2</b>	<b>Chapter 2</b>	<b>15</b>
2.1	Counting . . . . .	15
2.1.1	Ordered with replacement . . . . .	15
2.1.2	Ordered without replacement (Permutations) . . . . .	16
2.1.3	Unordered without replacement (Combinations) . . . . .	17
2.1.4	Unordered without replacement . . . . .	21
<b>3</b>	<b>Chapter 3</b>	<b>23</b>
3.1	Random variables . . . . .	23
3.1.1	Discrete random variables . . . . .	23
3.1.2	Probability mass function . . . . .	23
3.1.3	Independent random variables . . . . .	24
3.2	Special distributions . . . . .	24
3.2.1	Benroulli distribution . . . . .	24
3.2.2	Geometric distribution . . . . .	24
3.2.3	Binomial distribution . . . . .	25

3.2.4	Negative binomial (Pascal) distribution . . . . .	26
3.2.5	Hypergeometric distribution . . . . .	27
3.2.6	Poisson distribution . . . . .	28
3.3	Cumulative distribution function . . . . .	29
<b>A</b>	<b>Chapter problems</b>	<b>30</b>
A.1	Chapter 1 . . . . .	30
A.2	Chapter 2 . . . . .	51
A.3	Chapter 3 . . . . .	58

# 1 Chapter 1

## 1.1 Set operations

A union of 2 sets is given by the combination of their elements:

$$A \cup B = \{1, 2\} \cup \{2, 3\} = \{1, 2, 3\}$$

The intersection of 2 sets is instead given by their shared elements:

$$A \cap B = \{1, 2\} \cap \{2, 3\} = \{2\}$$

**Theorem 1** (De Morgan's law). *For any sets  $A_1, A_2, \dots, A_n$  we have*

$$\begin{aligned}\overline{A_1 \cup A_2 \cup \dots \cup A_n} &= \overline{A_1} \cap \overline{A_2} \cap \dots \cap \overline{A_n} \\ \overline{A_1 \cap A_2 \cap \dots \cap A_n} &= \overline{A_1} \cup \overline{A_2} \cup \dots \cup \overline{A_n}\end{aligned}$$

**Theorem 2** (Distributive law). *For any sets  $A, B$  and  $C$  we have*

$$\begin{aligned}A \cap (B \cup C) &= (A \cap B) \cup (A \cap C) \\ A \cup (B \cap C) &= (A \cup B) \cap (A \cup C)\end{aligned}$$

The complement of a set is given by all elements that are in the universal set, but not the set itself:

$$S = \{1, 2, 3, 4, 5\} \quad \overline{A} = S \setminus A = \{1, 2, 3, 4, 5\} \setminus \{1, 2\} = \{3, 4, 5\}$$

The difference between two sets is given by elements in the first but not the second:

$$A \setminus B = \{1, 2\} - \{2, 3\} = \{1\} \quad A \setminus B = A \cap \overline{B}$$

Two sets are disjoint if their intersection is an empty set

$$A \cap B = \emptyset$$

Sets can be partitioned into smaller parts. The sets  $A_1, A_2, \dots, A_n$  are a partition of  $S$  if they're disjoint and:

$$\bigcup_{i=1}^n A_i = S$$

The cartesian product of two sets are given by the ordered pairs of both sets:

$$A \times B = \{1, 2\} \times \{2, 3\} = \{(1, 2), (1, 3), (2, 2), (2, 3)\}$$

Which can be expressed more generally as:

$$A \times B = \{(x, y) | x \in A \text{ and } y \in B\}$$

The number of elements contained in a (finite) sets is given by its cardinality:

$$|A| = |\{1, 2\}| = 2$$

For determining the cardinality of (finite) sets, the inclusion-exclusion principle is often used:

$$|A \cup B| = |A| + |B| - |A \cap B|$$

This can once again be expanded to more sets:

$$\begin{aligned} \left| \bigcup_{i=1}^n A_i \right| &= \sum_{i=1}^n |A_i| - \sum_{i < j} |A_i \cap A_j| \\ &+ \sum_{i < j < k} |A_i \cap A_j \cap A_k| - \dots + (-1)^{n+1} |A_1 \cap \dots \cap A_n| \end{aligned}$$

### 1.1.1 Cardinality and countable sets

Finite sets are obviously countable, however when we move onto infinite sets they are divided into countable **and** uncountable sets. A countable set is characterised by the ability to write it in one-to-one correspondance with the natural numbers, e.g.:

$$A = \{a_1, a_2, \dots, a_n\}$$

Meaning you can list the elements, this is true for sets like the natural numbers,  $\mathbb{N}$ , and the integers,  $\mathbb{Z}$ , but also the rational numbers,  $\mathbb{Q}$ . Uncountable sets (such as the real- and complex numbers) on the other hand cannot be written as lists, but instead have to be denoted as intervals.

**Definition 1** (Countability of a set). A set,  $A$ , is called countable if one of the following is true:

- It is a finite set,  $|A| < \infty$ .
- The set can be written as a list with one-to-one correspondance with the natural numbers.

This means that any subset of  $\mathbb{N}$ ,  $\mathbb{Z}$  and  $\mathbb{Q}$  are countable, whilst any set containing an interval on the real line is uncountable.

**Theorem 3** (Countability of sub- and supersets). *Any subset of a countable set is countable and any superset of an uncountable set is uncountable.*

*Proof.* Let  $A$  be a countable set and  $B \subset A$ . If  $A$  is finite, then it follows that  $|B| \leq |A| < \infty$ , thus  $B$  must be countable as its cardinality cannot exceed that of  $A$ , which must be smaller than  $\infty$ .

If  $A$  is instead countably infinite, then it follows that as  $B$  is a subset of  $A$  it must be possible to construct it by removing  $\overline{B}$  from  $A$ , whereby it must also be countable, as it can be constructed as a list.

The opposite can be argued by assuming  $B$  is **not** countable, whereby a contradiction would occur in both proofs.  $\square$

**Theorem 4** (Countability of union). *If  $A_1, A_2, \dots, A_n$  are countable sets, then the union of those must also be countable.*

*Proof.* As the sets are countable it must be possible to write them in the form

$$\begin{aligned} A_1 &= \{a_{11}, a_{12}, \dots, a_{1n}\} \\ A_2 &= \{a_{21}, a_{22}, \dots, a_{2n}\} \\ A_3 &= \{a_{31}, a_{32}, \dots, a_{3n}\} \end{aligned}$$

As such the union of those sets must also be possible to construct as a list

$$\bigcup_{i=1}^m A_i = \{a_{11}, a_{12}, a_{21}, a_{22}, a_{31}, a_{32}, \dots, a_{mn}\}$$

And as a result must be countable.  $\square$

**Theorem 5** (Countability of cartesian product). *If  $A$  and  $B$  are countable, then  $A \times B$  is also countable.*

*Proof.* As  $A$  and  $B$  are countable it must be possible to write them in the form

$$\begin{aligned} A &= \{a_1, a_2, \dots, a_n\} \\ B &= \{b_1, b_2, \dots, b_n\} \end{aligned}$$

In accordance with the definition of the cartesian product, the two sets can be constructed as a list with the form

$$A \times B = \{(a_i, b_j) | i, j \in \mathbb{N}\}$$

Whereby it must be countable as it can be constructed as a list.  $\square$

As a result of this proof it also becomes clear that any set that can be written in the form

$$C = \bigcup_i \bigcup_j \{a_{ij}\} \text{ where } i, j \text{ belong to a countable set}$$

Must also be countable, the set of rational numbers is an example of this as it can be written as

$$\mathbb{Q} = \bigcup_{i \in \mathbb{Z}} \bigcup_{j \in \mathbb{N}} \left\{ \frac{i}{j} \right\}$$

### 1.1.2 Functions

Functions take an input from its domain, apply a rule to said input, whereby an output from the co-domain is produced.

$$f : A \rightarrow B \quad f(x \in A) \in B$$

**Definition 2** (Function). A function maps elements from the domain set to elements in the co-domain with the property that each input is mapped to exactly one output.

In the same context the range operand is defined, as it is not necessary for a function to be able to output all elements of the codomain:

$$f : \mathbb{R} \xrightarrow{x^2} \mathbb{R}$$

Here both the domain- and co-domain are the real numbers, however it is clear that no value  $x \in \mathbb{R}$  would ever produce a negative number, therefore:

$$\text{Range}(f) = \mathbb{R}^+$$

## 1.2 Random experiments

A random experiment will always have an **outcome** corresponding to an element from the **sample space**,  $S$ .

**Definition 3** (Random experiment). A random experiment is a process by which we observe something uncertain.

When a random experiment is repeated, each repetition is called a **trial**. The goal of analyzing a random experiment is to assign probabilities to **events**, which correspond to subsets of the sample space.

### 1.2.1 Probability

A probability is assigned to an event,  $P(A) \in [0, 1]$ . The derivation of probability theorem is based on 3 axioms:

- Axiom 1: For any event  $A$ ,  $1 \geq P(A) \geq 0$ .
- Axiom 2: Probability of the sample space,  $S$ , is  $P(S) = 1$ .
- Axiom 3: If  $A_1, A_2, \dots, A_n$  are disjoint events, then  $P(\bigcup_{i=1}^n A_i) = \sum_{i=1}^n P(A_i)$

Notationally unions and intersections can be read as:

$$\begin{aligned} P(A \cap B) &= P(A \text{ and } B) = P(A, B) \\ P(A \cup B) &= P(A \text{ or } B) \end{aligned}$$

**Theorem 6** (Probability of complement). *For any event  $A$ ,  $P(\bar{A}) = 1 - P(A)$ .*

*Proof.* As the complement of a set contains all elements of the sample space that are not in the set

$$\bar{A} = S \setminus A$$

It is clear that their unions must be  $S$  and they must be disjoint whereby

$$P(A \cup \bar{A}) = P(S) = 1$$

As they are disjoint we can write the probability of their union as the sum of their probabilities

$$P(A) + P(\bar{A}) = 1 \Leftrightarrow P(A) = 1 - P(\bar{A})$$

□

**Theorem 7** (Probability of empty set). *The probability of the empty is zero,  $P(\emptyset) = 0$ .*

*Proof.* As the empty set must be the complement of the sample space we have that

$$P(\emptyset) = P(\bar{S}) = 1 - P(S) = 1 - 1 = 0$$

□

**Theorem 8** (Probability must be equal to or less than 1). *For any event  $A$ ,  $P(A) \leq 1$ .*

*Proof.* By the first axiom we have that

$$P(\overline{A}) \geq 0$$

It becomes clear that

$$P(A) \leq 1$$

As  $P(A) + P(\overline{A}) = 1$ . □

**Theorem 9** (Probability of a difference). *The probability of a difference is given by  $P(A \setminus B) = P(A) - P(A \cap B)$ .*

*Proof.* As  $A \cap B$  and  $A \setminus B$  must be disjoint, whilst their union must be  $A$

$$(A \cap B) \cup (A \setminus B) = A$$

We have by the third axiom that

$$P(A) = P((A \cap B) \cup (A \setminus B)) = P(A \cap B) + P(A \setminus B)$$

By rearranging it becomes clear that

$$P(A \setminus B) = P(A) - P(A \cap B)$$

□

**Theorem 10** (Probability of a union). *The probability of a union is given by  $P(A \cup B) = P(A) + P(B) - P(A \cap B)$ .*

*Proof.* As  $A$  and  $B \setminus A$  must be disjoint sets whilst their union must be  $A \cup B$ , it is clear that

$$P(A \cup B) = P(A \cup (B \setminus A))$$

As we know these are disjoint we write

$$P(A \cup B) = P(A) + P(B \setminus A)$$

Rewriting using the previous theorem we then have

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

□



**Theorem 11** (Probability of a subset must be less than or equal to its superset). *If  $A \subset B$  then  $P(A) \leq P(B)$ .*

*Proof.* As  $A \subset B$  it is clear that their union must be  $B$

$$P(B) = P(A \cap B) + P(B \setminus A)$$

As their intersection is  $A$  we have that

$$P(B) = P(A) + P(B \setminus A)$$

As

$$P(B \setminus A) \geq 0$$

By the first axiom, we have that

$$P(B) \geq P(A)$$

□

### 1.3 Conditional probability

Conditional probabilities are written as

$$P(A|B)$$

And are read as "*the probability of  $A$ , given that  $B$  has occurred*". The conditional probability will therefore be given by

$$\begin{aligned} P(A|B) &= \frac{|A \cap B|}{|B|} \\ &= \frac{\frac{|A \cap B|}{|S|}}{\frac{|B|}{|S|}} \text{ dividing by } |S| \\ &= \frac{P(A \cap B)}{P(B)} \end{aligned}$$

This is as when we know  $B$  has occurred, the sample space of  $A$  is shrunk to  $B$ , whereby the cardinality of their intersection must be equal to the amount of favourable outcomes.

The earlier established probability axioms can also be formulated for conditional probabilities

- Axiom 1: For any event  $A$ ,  $P(A|B) \geq 0$ .
- Axiom 2: Conditional probability of  $B$  given  $B$  is 1, i.e.,  $P(B|B) = 1$ .
- Axiom 3: If  $A_1, A_2, \dots, A_n$  are disjoint events, then  $P(\bigcup_{i=1}^n A_i|B) = \sum_{i=1}^n P(A_i|B)$ .

And the same applies to the established fomulas

**Theorem 12.** For any conditional event,  $A|C$ ,  $P(\bar{A}|C) = 1 - P(A|C)$ .

*Proof.* We know that

$$P(A|B) = \frac{P(A|B)}{P(B)}$$

Assuming the theorem is correct we then have that

$$1 - P(\bar{A}|C) = 1 - \frac{P(\bar{A}|C)}{P(B)}$$

We wish to show that

$$\frac{P(A|B)}{P(B)} = 1 - \frac{P(\bar{A}|C)}{P(B)}$$

Multiplying by  $P(B)$  gets us

$$P(A|B) = P(B) - P(\bar{A}|C)$$

Adding  $P(\bar{A}|C)$  on the LHS we get

$$P(A|B) + P(\bar{A}|C) = P(B)$$

As the two are mutually exclusive by the definition of the complement

$$P((A|B) \cup (\bar{A}|C)) = P(B)$$

Which is true. □

**Theorem 13.** The probability of the empty set is zero,  $P(\emptyset|C) = 0$ .

*Proof.* As the empty is the complement of the sample set we have from the previous theorem that

$$P(S|C) = 1$$

By applying previous equation

$$P(\bar{S}|C) = 1 - 1 = 0$$

□

**Theorem 14.** *The probability of a conditional probability occurring must always be less than or equal to 1,  $P(A|C) \leq 1$ .*

*Proof.* From the definition of conditional probability

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

It is clear that the denominator and numerator can never be more than even as  $A \cap B = B$  even if  $AB$ , resulting in the maximum value being 1.

$$P(A|B) \leq 1$$

□

**Theorem 15.** *The probability of a difference is given by  $P(A \setminus B|C) = P(A|C) - P(A \cap B|C)$ .*

*Proof.* As  $A \cap B|C$  and  $A \setminus B|C$  must be disjoint whilst their union must be  $A|C$ , we have that

$$(A \cap B|C)(A \setminus B|C) = A|C$$

By the third axiom we have that

$$P(A|C) = P(A \cap B|C) + P(A \setminus B|C)$$

Rearranging the terms we get

$$P(A \setminus B|C) = P(A|C) - P(A \cap B|C)$$

□

**Theorem 16.** *The probability of a union is given by  $P(A \cup B|C) = P(A|C) + P(B|C) - P(A \cap B|C)$ .*

*Proof.* As  $A|C$  and  $B \setminus A|C$  must be disjoint and their union must be equal to  $A \cup B|C$  we have that

$$P(A \cup B) = P(A|C(B \setminus A|C))$$

As these sets are disjoint we rewrite using the third axiom

$$P(A \cup B) = P(A|C) + P(B \setminus A|C)$$

By the previous theorem the difference is rewritten as

$$P(A \cup B) = P(A|C) + P(B|C) - P(A \cap B|C)$$

□

**Theorem 17.** *If  $A \subset B$  then  $P(A|C) \leq P(B|C)$ .*

*Proof.* As  $A \subset B$  it is clear that their union must be  $B$

$$P(B|C) = P(A \cap B|C) + P(B \setminus A|C)$$

Since their intersection is  $A$  due to it being the subset

$$P(B|C) = P(A|C) + P(B \setminus A|C)$$

By the first axiom

$$P(B \setminus A|C) \geq 0$$

As such

$$P(B|C) \geq P(A|C)$$

□

This introduces some special cases

$$\begin{aligned} P(A|B) &= \frac{P(\emptyset)}{P(B)} = 0, \text{ for } A \cap B = \emptyset \\ P(A|B) &= \frac{P(B)}{P(B)} = 1, \text{ for } B \subset A \\ P(A|B) &= \frac{P(A)}{P(B)}, \text{ for } A \subset B \end{aligned}$$

Furthermore we also write the chain rule for conditional probability using the definition as a starting point

**Theorem 18.** *The extended chain rule is given by  $P(A_1 \cap A_2 \cap A_3 \cap \dots \cap A_n) = P(A_1)P(A_2|A_1)P(A_3|A_2, A_1) \cdots P(A_n|A_{n-1}, A_{n-2}, \dots, A_1)$*

*Proof.* From the definition of conditional probability we have that

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

By isolation for  $P(A \cap B)$  we get

$$P(A \cap B) = P(A|B)P(B) = P(B|A)P(A)$$

Extending to 3 or more events we get that

$$P(A \cap B \cap C) = P(A \cap (B \cap C)) = P(A)P(B \cap C|A)$$

Applying the first equation

$$P(B \cap C) = P(B)P(C|B)$$

By conditioning both sides on  $A$  we get

$$P(B \cap C|A) = P(B|A)P(C|A, B)$$

Inserting in the original equation we then get

$$P(A \cap B \cap C) = P(A)P(B|A)P(C|A, B)$$

Which can be generalised to

$$P(A_1 \cap A_2 \cap \dots \cap A_n) = P(A_1)P(A_2|A_1)P(A_3|A_2, A_1) \dots \\ P(A_n|A_{n-1}, A_{n-2}, \dots, A_1)$$

□

### 1.3.1 Independence

Conditional probabilities are only relevant if two events are not independent.

**Definition 4** (Independence). Two events  $A, B$  are independent if  $P(A \cap B) = P(A)P(B)$ .

Independence for two or more events requires that all the individual events are independent, as well as all of them together, this means that for 3 events, all of the following must hold

$$\begin{aligned} P(A \cap B) &= P(A)P(B) \\ P(A \cap C) &= P(A)P(C) \\ P(B \cap C) &= P(B)P(C) \\ P(A \cap B \cap C) &= P(A)P(B)P(C) \end{aligned}$$

**Theorem 19.** *If  $A$  and  $B$  are independent, then  $A$  and  $\overline{B}$ ,  $\overline{A}$  and  $B$ ,  $\overline{A}$  and  $\overline{B}$  are also independent.*

*Proof.* The first statement is proven as the others can be concluded from it. As the statement is equivalent to

$$P(A \cap \overline{B}) = P(A) - P(A \cap B)$$

As we know  $A$  and  $B$  are independent we have

$$P(A \cap \overline{B}) = P(A) - P(A)P(B)$$

We factor out  $P(A)$  and get that

$$P(A \cap \overline{B}) = P(A)(1 - P(B))$$

And as

$$1 - P(B) = P(\overline{B})$$

It becomes clear that

$$P(A \cap \overline{B}) = P(A)P(\overline{B})$$

□

To determine the probability of several unions of independent events, we make use of De Morgans law

$$P\left(\bigcup_{i=1}^n A_i\right) = 1 - P\left(\bigcap_{i=1}^n \overline{A_i}\right)$$

Which is equivalent to

$$P\left(\bigcup_{i=1}^n A_i\right) = 1 - \prod_{i=1}^n (1 - P(A_i))$$

### 1.3.2 Law of Total Probability

The law of total probability states that the probability of an event,  $A$  must be the sum of the probability of it occurring in every partition,  $B_1, B_2, \dots, B_n$  of the sample space

$$P(A) = \sum_{i=1}^n P(A \cap B_i) = \sum_{i=1}^n P(A|B_i)P(B_i)$$

*Proof.* As  $B_1, B_2, \dots, B_n$  are partitions of the sample space we write

$$S = \bigcup_{i=1}^n B_i$$

Now the event  $A$  occurring must be given by its intersection of the sample space

$$A = A \cap \left( \bigcup_{i=1}^n B_i \right)$$

By the distributive property it becomes clear that

$$A = \bigcup_{i=1}^n (A \cap B_i)$$

Now as the partitions by definition are disjoint we can determine the probability as the sum of probabilities

$$P(A) = P\left(\bigcup_{i=1}^n (A \cap B_i)\right) = \sum_{i=1}^n P(A \cap B_i)$$

Rewriting using the definition of conditional probability (as  $A \in B_i$  can only occur if  $B_i$  has occurred) we get

$$P(A) = \sum_{i=1}^n P(A|B_i)P(B_i)$$

□

**Theorem 20.** Bayes rule states that:  $P(B|A) = \frac{P(A|B)P(B)}{P(A)}$

*Proof.* From the definition of conditional probability

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

Multiplying by  $P(B)$  on both sides

$$P(A|B)P(B) = P(A \cap B)$$

Dividing by  $P(A)$  on both sides

$$\frac{P(A|B)P(B)}{P(A)} = \frac{P(A \cap B)}{P(A)} = P(B|A)$$

□

### 1.3.3 Conditional Independence

**Definition 5** (Conditional independence). Two events  $A$  and  $B$  are conditionally independent given an event  $C$  if  $P(A \cap B|C) = P(A|C)P(B|C)$ .

*Proof.* From the definition of conditional probability we have

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

Conditioning both sides on  $C$  it becomes apparent that

$$P(A|B, C) = \frac{P(A \cap B|C)}{P(B|C)}$$

Assuming that  $A$  and  $B$  are conditionally independent we have

$$P(A|B, C) = \frac{P(A|C)P(B|C)}{P(B|C)} = P(A|C)$$

□



## 2 Chapter 2

### 2.1 Counting

For a finite sample space with equal probabilities we recall that

$$P(A) = \frac{|A|}{|S|}$$

As such determining the probability is a counting problem, determining the cardinality of  $A$  and  $S$ .

**Definition 6** (Multiplication principle). Suppose that we perform  $r$  experiments such that the  $k$ 'th experiment has  $n_k$  possible outcomes for  $k = \{1, 2, \dots, r\}$ . Then there are a total of  $n_1 \times n_2 \times \dots \times n_r$  possible outcomes for the sequence of  $r$  experiments.

For counting problems, some general terminology is relevant:

- Sampling: Choosing a random element from a set.
- With replacement, the sampled element is returned to the set and can therefore be drawn multiple times with repeated sampling.
- Without replacement, the sampled element is not returned to the set and can therefore not be drawn multiple times.
- Ordered means that the order at which elements are written matters,  $\{a_1, a_2, a_3\} \neq \{a_3, a_1, a_2\}$ .
- Unordered means that the order at which elements are written does not matter,  $\{a_1, a_2, a_3\} = \{a_3, a_1, a_2\}$ .

#### 2.1.1 Ordered with replacement

Suppose we have a set consisting of  $n$  elements and we wish to draw  $k$  samples from the set, for example, say  $A = \{1, 2, 3\}$  where we wish to sample  $k = 2$ , we then get 9 different possibilities

$$(1, 1), (1, 2), (1, 3), (2, 1), (2, 2), (2, 3), (3, 1), (3, 2), (3, 3)$$

From this it's clear that we create a list consisting of  $k$ -valued elements where each position has  $n$  options for values

$$\begin{array}{cccc} a_1 & a_2 & \dots & a_k \\ \uparrow & \uparrow & & \uparrow \\ n & n & & n \end{array}$$

Meaning that we can determine the total amount of possibilities as

$$n \times n \times \dots \times n = n^k$$

### 2.1.2 Ordered without replacement (Permutations)

As opposed to the previous circumstance, a element is now removed every time we draw, resulting in there being one less options every time we move to the next position, as such, using the same example with  $A$  and  $k = 2$  we get

$$(1, 2), (1, 3), (2, 1), (2, 3), (3, 1), (3, 2)$$

Here we create a list of  $k$ -valued elements where each position has one less option than the previous

$$\begin{array}{cccc} (a_1 & a_2 & \dots & a_k) \\ \uparrow & \uparrow & & \uparrow \\ n & n-1 & & n-k+1 \end{array}$$

Which is called a  $k$ -permutation of the elements in the set, the following notation is used to show the number of  $k$ -permutations of an  $n$ -element set

$$P_k^n = n \times (n-1) \times \dots \times (n-k+1)$$

A special case can also occur here when  $n < k$ , as there then wont be enough options for every position and there therefore are no possible lists. Another special case is an  $n$ -permutation where  $k = n$  which results in the sequence

$$\begin{aligned} P_n^n &= n \times (n-1) \times (n-2) \times \dots \times (n-n+1) \\ &= n \times (n-1) \times (n-2) \times \dots \times 1 \\ &= n! \end{aligned}$$

As such the factorial operator simply denotes the total number of permutations of an  $n$  element set, aka the total number of ways you can order  $n$  different objects. By definition  $0! = 1$ , using this we rewrite the formula for  $P_k^n$

**Theorem 21.** *The amount of  $k$ -permutations of an  $n$ -element set is given by  $P_k^n = \frac{n!}{(n-k)!}$ .*

*Proof.* From the original expression for  $P_k^n$  we have that

$$P_k^n = n \times (n-1) \times (n-2) \times \dots \times (n-k+1)$$

By multiplying by  $\frac{(n-k)!}{(n-k)!}$  we get

$$\begin{aligned} &= n \times (n-1) \times (n-2) \times \dots \times (n-k+1) \cdot \frac{(n-k)!}{(n-k)!} \\ &= \frac{n!}{(n-k)!} \end{aligned}$$

As multiplying by  $(n-k)!$  in the numerator results in the sequence “finishing” and being equal to  $n!$ .  $\square$

### 2.1.3 Unordered without replacement (Combinations)

We now wish to determine the amount of possible lists when we sample  $k$  elements from an  $n$ -element set. This means that we want to determine the amount of possible  $k$ -element subsets of the  $n$ -element set. Using the same example as before with  $A$  and  $k = 2$  we get 3 different combinations

$$(1, 2), (1, 3), (2, 3)$$

We show the number of  $k$ -element subsets of  $A$  as

$$\binom{n}{k}$$

Which is read as “ $n$  choose  $k$ ”, to determine the value of this we compare with  $P_k^n$  as the only difference between the two is ordering. This is because for any  $k$ -element subset of an  $n$ -element set, we can order the elements in  $k!$  different ways, as such

$$P_k^n = \binom{n}{k} \times k!$$

Rewriting using our previously established formula for  $P_k^n$  and dividing by  $k!$  we get

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}, \text{ for } 0 \leq k \leq n$$

This term is used extensively in the binomial theorem, which states that

$$(a+b)^n = \sum_{k=0}^n \binom{n}{k} a^k b^{n-k}$$

**Theorem 22.** *For any non-negative integers  $n, k$  it follows that  $\binom{n}{k} = \binom{n}{n-k}$ .*

*Proof.* Assume we wish to determine the amount of possible sequences consisting of  $k$  A's and  $j$  B's, as such we have  $n = j + k$  positions to fill with either A or B, from these positions we need to choose  $j$  for A's and whatever is left is filled with B's, as such the amount of ways is

$$\binom{n}{j}$$

If we instead observe this from the point of B's, it is clear that the amount of ways would then be given by

$$\binom{n}{k}$$

As these must be equivalent we have that

$$\binom{n}{j} = \binom{n}{k}$$

From the initial determination of  $n$  we get that

$$n = j + k \implies j = n - k \vee k = n - j$$

As such

$$\binom{n}{j} = \binom{n}{n-k} = \binom{n}{k} = \binom{n}{n-j}$$

□

**Theorem 23.** For any non-negative integers  $k, n$  it follows that  $\sum_{k=0}^n \binom{n}{k} = 2^n$ .

*Proof.* From the Binomial theorem we know that

$$(a + b)^n = \sum_{k=0}^n \binom{n}{k} a^k b^{n-k}$$

We let  $a = b = 1$  and as such get

$$2^n = \sum_{k=0}^n \binom{n}{k}$$

□

**Theorem 24.** For non-negative integers  $0 \leq k \leq n$  it follows that  $\binom{n+1}{k+1} = \binom{n}{k+1} + \binom{n}{k}$ .

*Proof.* We define an arbitrary set,  $A$  with  $n+1$  elements

$$A = \{a_1, a_2, \dots, a_n, a_{n+1}\}$$

From this set we wish to choose a  $k+1$  element subset, call it  $B$ , by combinations we know this is equal to

$$\binom{n+1}{k+1}$$

$B$  can also be constructed as the union of two subsets of  $B$  that are defined by either containing- or not containing  $a_{n+1}$

$$B = B_1 \cup B_2, \text{ where } a_{n+1} \notin B_1, a_{n+1} \in B_2, B_1 \cap B_2 = \emptyset$$

To define  $B_1$  we need to choose  $k+1$  elements from the set  $A \setminus a_{n+1}$  which is equal to

$$\binom{n}{k+1}$$

To complete the set we then need to choose  $k$  elements from  $A$  which can be done in

$$\binom{n+1}{k}$$

ways. As such the sum of the two must be equal to the initial expression resulting in

$$\binom{n+1}{k+1} = \binom{n}{k+1} + \binom{n}{k}$$

□

**Theorem 25.** Vandermonde's identity states that  $\binom{m+n}{k} = \sum_{i=0}^k \binom{m}{i} \binom{n}{k-i}$

*Proof.* We construct a set  $A$  with  $m+n$  elements, as such

$$A = \{a_1, a_2, \dots, a_m, b_1, b_2, \dots, b_n\}$$

Determining the number of  $k$ -element subsets of  $A$  is equal to

$$\binom{m+n}{k}$$

This can also be done by choosing  $i$  elements from  $\{a_1, a_2, \dots, a_m\}$  first, and then  $k-i$  elements from  $\{b_1, b_2, \dots, b_n\}$ , which then can be done in

$$\binom{m}{i} \binom{n}{k-i}$$

ways. But as  $i$  can be any number from  $0 \rightarrow k$  it is necessary to sum all the possible options whereby we write

$$\binom{m+n}{k} = \sum_{i=0}^k \binom{m}{i} \binom{n}{k-i}$$

□

An important class of random experiments are Bernoulli trials, a random experiment where there are two possible outcomes, success and failure (which can be extended to any experiment with 2 outcomes as we can arbitrarily define success and failure).

In Bernoulli trials the probability of success is usually denoted by  $p$  and the probability of failure as its complement  $q = 1 - p$ . If we perform  $n$  independent Bernoulli trials and count the number of successes, it is called a binomial experiment, for example a coin toss where we define success as heads and count the number of heads.

**Theorem 26.** *The binomial formula is given by  $P(k) = \binom{n}{k} p^k (1-p)^{n-k}$*

*Proof.* Imagine we toss a coin with  $P(H) = p$  and  $P(T) = 1 - p$   $n$  times, we define  $C$  as the event of observing  $k$  heads (and  $n-k$  tails), the probability of observing  $k$  heads will be given by

$$P(k) = |C| p^k (1-p)^{n-k}$$

To determine  $|C|$ , we realise that we can see the event as ordered sampling without replacement, as such we have that

$$|C| = \binom{n}{k}$$

Which we insert into the previous expression and get

$$P(k) = \binom{n}{k} p^k (1-p)^{n-k}$$

□

**Theorem 27.** *If we have  $n$  people and wish to separate them into  $r$  groups, the number of ways this is possible is given by the multinomial coefficients,*

$$\binom{n}{n_1, n_2, \dots, n_r} = \frac{n!}{n_1! n_2! \dots n_r!}.$$

*Proof.* Say we have  $n$  people we wish to divide into  $r$  groups, using combinations, this would be given as

$$\begin{aligned} \binom{n}{k_1} \binom{n-k_1}{k_2} \dots \binom{n-k_1-\dots-k_{r-1}}{k_r} &= \frac{n!}{k_1!(n-k_1)!} \cdot \frac{(n-k_1)!}{k_2!(n-k_1-k_2)!} \\ &\dots \frac{(n-k_1-\dots-k_{r-1})!}{k_r!(n-k_1-\dots-k_{r-1}-k_r)!} = \frac{n!}{k_1 k_2! \dots k_r!} \end{aligned}$$

The simplification of this expression results from the following term always cancelling the “excess” in the previous, until we reach the final term where the numerator cancels the excess term from the previous fraction, whilst the denominator equals one as  $n = \sum_{i=1}^r k_i \implies (n - k_1 - \dots - k_r)! = (0)! = 1$  □

### 2.1.4 Unordered without replacement

As opposed to the previous section we’re now working with replacement, as such we again use the example of  $A$  with  $k = 2$  and get 6 possibilities given by

$$(1, 1), (1, 2), (1, 3), (2, 2), (2, 3), (3, 3)$$

One way to represent this sample is to list them as  $n$ -element vectors where each position corresponds to a number, eg.

$$\begin{aligned} (a, b) &\rightarrow (x_1, x_2, x_3) = (n_1, n_2, n_3) \\ (1, 1) &\rightarrow (x_1, x_2, x_3) = (2, 0, 0) \\ (1, 2) &\rightarrow (x_1, x_2, x_3) = (1, 1, 0) \\ (1, 3) &\rightarrow (x_1, x_2, x_3) = (1, 0, 1) \end{aligned}$$

Constructing these vectors a pattern emerges, that is  $\sum_{i=1}^3 x_i = 2$ , as such we can determine the amount of possibilities as the amount of integer solutions to  $x_1 + x_2 + x_3 = 2$ .

**Theorem 28.** *The number of distinct solutions to the equation  $x_1 + x_2 + \dots + x_n = k$  where  $x_n \in \{0, 1, 2, 3, \dots\}$  is equal to  $\binom{n+k-1}{k} = \binom{n+k-1}{n-1}$ .*

*Proof.* We first define a mapping where an integer  $x_n \geq 0$  is replaced with  $x_n$  vertical lines. Suppose we now have a solution to the equation where  $k = 6$ , for example

$$3 + 0 + 2 + 1 \Leftrightarrow ||| + + || + |$$

Here we realise that the equation can be represented by  $k$  vertical lines and  $n - 1$  plus signs, as such we can use combinations to determine the amount of distinct sequences we can create using  $k$  vertical lines and  $n - 1$  plus signs, as such

$$n_{\text{solutions}} = \binom{k+n-1}{k} = \binom{k+n-1}{n-1}$$

□



### 3 Chapter 3

#### 3.1 Random variables

**Definition 7** (Random variable). A random variable is a real-valued value determined by an underlying random experiment. Mathematically this can be expressed as a function that assigns a value to a possible outcome,  $S \xrightarrow{X} \mathbb{R}$ .

##### 3.1.1 Discrete random variables

What defines whether a random variable is discrete, continuous or mixed is its range. If the range of the random variable is strictly countable, the random variable is said to be discrete, this for example occurs if we wish to determine the amount of heads in a coin throw.

##### 3.1.2 Probability mass function

Let  $X$  be a discrete random variable, as such

$$\text{Range}(X) = \{x_1, x_2, x_3, \dots\}$$

As the possible outcomes are discrete, they must each have their own probability of occurring, we define the event

$$A = \{s \in S | X(s) = x_i\}$$

Where the probabilities of each individual event  $\{X = x_i\}$  is given by the probability mass function of  $X$

$$P_X(x_i) = P(X = x_i), \text{ for } i = 1, 2, 3, \dots$$

To ensure the function is well-defined, the definition of the mass probability function is frequently extended to all real numbers by a piecewise function

$$P_X(x) = \begin{cases} P(X = x) & x \in \text{Range}(X) \\ 0 & x \notin \text{Range}(X) \end{cases}$$

As this is a plottable discrete function, this is also called the probability distribution of the random variable  $X$ .

### 3.1.3 Independent random variables

Random variables can, like probabilities and conditional probabilities, be independent. Independence is defined in a similar way to in those examples, 2 random variables,  $X$  and  $Y$ , are independent if

$$P(X = x \cap Y = y) = P(X = x)P(Y = y), \text{ for all } x, y$$

This means that if they are independent we can write

$$P(X \in A \cap Y \in B) = P(X \in A)P(Y \in B), \text{ for all sets } A \text{ and } B$$

We can also condition random variables on eachother, for two independent random variables it then follows that

$$P(X = x|Y = y) = P(X = x), \text{ for all } x, y$$

As knowing the outcome of  $Y$  does not impact the probability of  $X$ .

## 3.2 Special distributions

### 3.2.1 Benroulli distribution

We recall that a Bernoulli experiment is an experiment with only two possible outcomes, success or failure, defining  $x = 1$  as success and  $x = 0$  as failure, we get the Bernoulli PDF

$$P_X(x) = \begin{cases} p & x = 1 \\ 1 - p & x = 0 \\ 0 & \text{otherwise} \end{cases}$$

### 3.2.2 Geometric distribution

For a geometric experiment, that being a Bernoulli experiment where we repeat the experiment until one of the two outcomes is observed has the probability  $P(A) = p(1-p)^{k-1}$ , where  $p$  is the probability of observing what we want, as such the PDF is given by

$$P_X(x) = \begin{cases} p(1-p)^{k-1} & k \in \mathbb{N} \\ 0 & \text{otherwise} \end{cases}$$

### 3.2.3 Binomial distribution

We recall that the binomial theorem states that

$$P(k) = \binom{n}{k} p^k (1-p)^{n-k}$$

As such the PDF is given by

$$P_X(k) = \begin{cases} \binom{n}{k} p^k (1-p)^{n-k} & k \in \{0, 1, 2, \dots, n\} \\ 0 & \text{otherwise} \end{cases}$$

Simultaneously we can observe each cointoss that makes up the sequence as individual Bernoulli distributions, as such we can express a binomial as the sum of a sequence of Bernoulli experiments, i.e. we have the PDF  $X$  defined as the binomial giving the number of heads, which can be expressed as

$$X = \sum_{i=1}^n X_i$$

Where  $i$  is the  $i$ th cointoss in the sequence as given by the Bernoulli PDF.

**Theorem 29.** *Given two independent random variables described by binomial distributions,  $X, Y$ , with  $k_1 = n$  and  $k_2 = m$  the random variable defined as  $Z = X + Y$  is given by*

$$P_Z(k) = \begin{cases} \binom{n+m}{k} p^k (1-p)^{n+m-k} & k \in \{0, 1, 2, \dots, m+n\} \\ 0 & \text{otherwise} \end{cases}$$

*Proof.* As the random variable can be described as the sum of the individual Bernoulli random variables we rewrite that

$$\begin{aligned} Z &= X + Y \\ &= X_1 + X_2 + \dots + X_n + Y_1 + Y_2 + \dots + Y_n \end{aligned}$$

A way to express this as probabilities is

$$\begin{aligned}
P_Z(k) &= P(Z = k) \\
&= P(X + Y = k) \\
&= \sum_{i=0}^n P(X + Y = k | X = i) P(X = i) \\
&= \sum_{i=0}^n P(X = k - i) P(X = i) \\
&= \sum_{i=0}^n \binom{m}{k-i} p^{k-i} (1-p)^{m-k+i} \binom{n}{i} p^i (1-p)^{n-i} \\
&= \sum_{i=0}^n \binom{m}{k-i} \binom{n}{i} p^k (1-p)^{m-k+n} \\
&= p^k (1-p)^{m-k+n} \sum_{i=0}^n \binom{m}{k-i} \binom{n}{i} \\
&= \binom{m+n}{k} p^k (1-p)^{m-k+n} \\
&= \begin{cases} \binom{m+n}{k} p^k (1-p)^{m-k+n} & k \in \{0, 1, 2, \dots, m+n\} \\ 0 & \text{otherwise} \end{cases}
\end{aligned}$$

□

### 3.2.4 Negative binomial (Pascal) distribution

The Pascal distribution is a generalization of the geometric distribution where we wish to observe  $m$  successes instead of just 1, by this it follows that a Pascal distribution with  $m = 1$  is simply a geometric distribution.

Suppose we toss a coin until  $m$  heads are observed, where  $X$  is the total number of tosses, in this context we define the event  $A = \{X = k\}$  as the event that  $m$  heads are observed in  $k$  tosses. We rewrite this event as the intersection  $A = B \cap C$  where

- $B$  is the event that  $m - 1$  heads are observed in the first  $k - 1$  tosses.
- $C$  is the event that heads is observed in the  $k$ th toss.

As these two events are independent (seperate coin tosses), we have that

$$P(A) = P(B \cap C) = P(B)P(C)$$

We can determine these probabilities.  $B$  is given by a binomial distribution whilst  $C$  is a constant value, as such we write that

$$\begin{aligned} P(B) &= \binom{k-1}{m-1} p^{m-1} (1-p)^{(k-1)-(m-1)} \\ &= \binom{k-1}{m-1} p^{m-1} (1-p)^{k-m} \\ P(C) &= p \end{aligned}$$

Since  $A$  is given by the product of these, we multiply them together whereby we get

$$\begin{aligned} P(A) &= P(B)P(C) \\ &= \binom{k-1}{m-1} p^{m-1} (1-p)^{k-m} p \\ &= \binom{k-1}{m-1} p^m (1-p)^{k-m} \end{aligned}$$

Whereby we can write the PDF for the Pascal distribution as

$$P_X(k) = \begin{cases} \binom{k-1}{m-1} p^m (1-p)^{k-m} & k \in \{m, m+1, m+2, \dots\} \\ 0 & \text{otherwise} \end{cases}$$

### 3.2.5 Hypergeometric distribution

Imagine a bag containing  $b$  blue marbles and  $r$  red ones. Choosing  $k \leq b+r$  marbles at random without replacements will at best give us  $X \leq \min(k, b)$  blue marbles, whilst the number of red marbles in our sample must be  $X \geq \max(0, k-r)$ . As such we can define the range of  $X$  as

$$\text{Range}(X) = \{\max(0, k-r), \max(0, k-r) + 1, \dots, \min(k, b)\}$$

Simultaneously we can create an expression for determining the probability of getting  $x$  blue marbles (and  $k-x$  red ones) by

$$\frac{\binom{b}{x} \binom{r}{k-x}}{\binom{b+r}{k}}, \text{ for } x \in \text{Range}(X)$$

Using this expression we create the PDF as

$$P_X(x) = \begin{cases} \frac{\binom{b}{x} \binom{r}{k-x}}{\binom{b+r}{k}} & x \in \text{Range}(X) \\ 0 & \text{otherwise} \end{cases}$$

### 3.2.6 Poisson distribution

The Poisson distribution is typically used to model occurrences in an interval using a variable,  $\lambda$ , as the known average.

**Theorem 30.** *Let  $X$  be a binomial PDF with  $p = \frac{\lambda}{n}, \lambda > 0$ . Then for any  $k \in \{0, 1, 2, \dots\}$  we have that  $\lim_{n \rightarrow \infty} P_X(k) = \frac{e^{-\lambda} \lambda^k}{k!}$ .*

*Proof.* The binomial will be given by

$$P_X(k) = \binom{n}{k} \left(\frac{\lambda}{n}\right)^k \left(1 - \frac{\lambda}{n}\right)^{n-k}$$

Taking the limit of this binomial we get

$$\begin{aligned} \lim_{n \rightarrow \infty} P_X(k) &= \lim_{n \rightarrow \infty} \binom{n}{k} \left(\frac{\lambda}{n}\right)^k \left(1 - \frac{\lambda}{n}\right)^{n-k} \\ &= \lambda^k \lim_{n \rightarrow \infty} \frac{n!}{k!(n-k)!} \left(\frac{1}{n^k}\right) \left(1 - \frac{\lambda}{n}\right)^{n-k} \\ &= \frac{\lambda^k}{k!} \lim_{n \rightarrow \infty} \frac{n!}{(n-k)!} \left(\frac{1}{n^k}\right) \left(1 - \frac{\lambda}{n}\right)^n \left(1 - \frac{\lambda}{n}\right)^{-k} \\ &= \frac{\lambda^k}{k!} \lim_{n \rightarrow \infty} \left(\frac{n(n-1) \dots (n-k+1)}{n^k}\right) \left(1 - \frac{\lambda}{n}\right)^n \left(1 - \frac{\lambda}{n}\right)^{-k} \end{aligned}$$

We now take the limit of each of these expressions individually

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{n(n-1) \dots (n-k+1)}{n^k} &= 1 \\ \lim_{n \rightarrow \infty} \left(1 - \frac{\lambda}{n}\right)^n &= e^{-\lambda} \\ \lim_{n \rightarrow \infty} \left(1 - \frac{\lambda}{n}\right)^{-k} &= 1 \end{aligned}$$

Which gives us

$$\begin{aligned} P_X(k) &= \frac{\lambda^k}{k!} \cdot 1 \cdot 1 \cdot e^{-\lambda} \\ &= \frac{e^{-\lambda} \lambda^k}{k!} \end{aligned}$$

□

As such we can define the Poisson distributions PDF as

$$P_X(k) \begin{cases} \frac{e^{-\lambda} \lambda^k}{k!} & k \in \text{Range}(X) \\ 0 & \text{otherwise} \end{cases}$$

### 3.3 Cumulative distribution function

**Definition 8.** The cumulative distribution functions of a random variable  $X$  is defined as  $F_X(x) = P(X \leq x)$ , for all  $x \in \mathbb{R}$ .

## A Chapter problems

### A.1 Chapter 1

**Problem 1** Suppose the universal set is defined as  $S = \{x \in \mathbb{N} | 1 \leq x \leq 10\}$ ,  $A = \{1, 2, 3\}$ ,  $B = \{x \in S | 2 \leq x \leq 7\}$  and  $C = \{7, 8, 9, 10\}$ .

a) Find  $A \cup B$

$$A \cup B = \{1, 2, 3\} \cup \{2, 3, 4, 5, 6, 7\} = \{1, 2, 3, 4, 5, 6, 7\}$$

b) Find  $(A \cup C) \setminus B$

$$(A \cup C) \setminus B = \{1, 2, 3, 7, 8, 9, 10\} \setminus \{2, 3, 4, 5, 6, 7\} = \{1, 8, 9, 10\}$$

c) Find  $\overline{A} \cup (B \setminus C)$

$$\overline{A} \cup (B \setminus C) = \{4, 5, 6, 7, 8, 9, 10\} \cup \{2, 3, 4, 5, 6\} = \{2, 3, 4, 5, 6, 7, 8, 9, 10\}$$

d) Do  $A$ ,  $B$  and  $C$  form a partition of  $S$ ?

No as  $A \cap B \neq \emptyset$  and  $B \cap C \neq \emptyset$ .

**Problem 2** When working with real numbers, our universal set is  $\mathbb{R}$ . Find each of the following sets a)  $[6, 8] \cup [2, 7[$

$$[2, 8]$$

b)  $[6, 8] \cap [2, 7[$

$$[6, 7[$$

c)  $\overline{[0, 1]}$

$$]-\infty, 0[ \cup ]1, \infty[$$

d)  $[6, 8] \setminus ]2, 7[$

$$[7, 8]$$

**Problem 3** For each of the following Venn diagrams, write the sets denoted by the shaded area

a)

$$A \cup B \setminus A \cap B$$

b)

$$B \setminus C$$



c)

$$A \cap B + A \cap C$$

d)

$$C \cup (A \cap B) \setminus ((C \cap A) \cup (C \cap B))$$

**Problem 4** A coin is tossed twice. Let  $S = \{H, T\} \times \{H, T\}$ . Write the following sets by listing their elements:

a) First toss is heads

$$A = \{(H, T), (H, H)\}$$

b) At least one tails

$$B = \{(T, T), (T, H), (H, T)\}$$

c) Two tosses are different

$$C = \{(H, T), (T, H)\}$$

**Problem 5** Let  $A = \{1, 2, \dots, 100\}$ . For any  $i \in \mathbb{N}$ , define  $A_i$  as the set of numbers in  $A$  that are divisible by  $i$ .

a)

$$|A_1| = \left\lfloor \frac{|S|}{1} \right\rfloor = \left\lfloor \frac{100}{1} \right\rfloor = 100$$

$$|A_2| = \left\lfloor \frac{|S|}{2} \right\rfloor = \left\lfloor \frac{100}{2} \right\rfloor = 50$$

$$|A_3| = \left\lfloor \frac{|S|}{3} \right\rfloor = \left\lfloor \frac{100}{3} \right\rfloor = 33$$

$$|A_4| = \left\lfloor \frac{|S|}{4} \right\rfloor = \left\lfloor \frac{100}{4} \right\rfloor = 25$$

$$|A_5| = \left\lfloor \frac{|S|}{5} \right\rfloor = \left\lfloor \frac{100}{5} \right\rfloor = 20$$

b)

By the distributive property we have that

$$A_2 \cap A_3 \cap A_5 = (A_2 \cap A_3) \cap A_5$$

As the intersection of  $A_2$  and  $A_3$  must be the even factors of 3 we get

$$A_2 \cap A_3 = \{x \in A | x = 6n, n \in \mathbb{N}\}$$

The cardinality of this must be half of the original as every other value is valid

$$|A_2 \cap A_3| = \left\lfloor \frac{33}{2} \right\rfloor = 16$$

As only every fifth factor of 6 is a factor of 5 we get that

$$A_2 \cap A_3 \cap A_5 = \{x \in A | x = 30n, n \in \mathbb{N}\}$$

As this is every fifth value of the previous intersection, the cardinality must be one fifth of the previous

$$|A_2 \cap A_3 \cap A_5| = \left\lfloor \frac{16}{5} \right\rfloor = 3$$

**Problem 6** As  $A_1, A_2, A_3$  form a partition of the universal set, the cardinality of  $B$  must be equal to the sum of the cardinalities in the individual partitions.

$$|B| = \sum_{i=1}^3 |B \cap A_i| = 10 + 20 + 15 = 45$$

**Problem 7** a) As the numbers can be listed in one-to-one correspondance with the natural numbers the set is countable.

b) As the set is made up of the union of 2 countable sets we have that

$$B = \bigcup_{i \in \mathbb{Q}} \bigcup_{j \in \mathbb{Q}} \{a_i + b_j \sqrt{2}\}$$

As such it must be countable as its constituents are.

c) As the set is a subset of an uncountable set (real numbers), it is not countable.

**Problem 8** We take the limit of the upper bound of the interval as  $n \rightarrow \infty$

$$\lim_{n \rightarrow \infty} \frac{n-1}{n} = 1$$

As this limit is the upper bound of the interval, corresponding to the maximal interval included in the union of sets we get that

$$A = \bigcup_{n=1}^{\infty} A_n = [0; 1[$$

**Problem 9** Opposed to the previous problem the smallest set will here define the set as the intersection is limited to the smallest component. We take the limit as  $n \rightarrow \infty$  as the value is inverse proportional to  $n$

$$\lim_{n \rightarrow \infty} \frac{1}{n} = 0$$

As such

$$A = \bigcap_{n=1}^{\infty} A_n = \{0\}$$

**Problem 10** a)

b)

**Problem 11** As the set is given as an interval it is clear that

$$[0, 1[ \subset \mathbb{R}$$

As subsets of uncountable sets are uncountable, it becomes clear that the range is uncountable.

**Problem 12** a)

Reading the function it becomes clear that the domain of the function is given by

$$\{H, T\}^3$$

While the co-domain is given by

$$\mathbb{N} \cup \{0\}$$

b)

As the function is limited by the amount of heads that can appear in the sequence it is clear that

$$\text{Range}(f) = \{0, 1, 2, 3\}$$

c)

Knowing that  $x = 2$  tells us that 2 heads are present in the sequence, and as such 1 tails must also be present, as such the possible events are

$$\{(H, H, T), (H, T, H), (T, H, H)\}$$

**Problem 13** a)

We logically assume that the events are disjoint as two teams cant win, as such it is clear that

$$0.5 + P(b) + 0.25 = 1 \Leftrightarrow P(b) = 1 - 0.5 - 0.25 = 0.25$$

b) As the events are disjoint we determine the probability as

$$P(b \cup d) = 0.25 + 0.25 = 0.5$$

**Problem 14** a)

By inclusion-exclusion principle we have that

$$P(A \cap B) = P(A) + P(B) - P(A \cup B) = 0.4 + 0.7 - 0.9 = 0.2$$

b)

We have that

$$P(\bar{A} \cap B) = P(B \setminus A)$$

Expanding this expression we get that

$$P(\bar{A} \cap B) = P(B) - P(A \cap B) = 0.7 - 0.2 = 0.5$$

c)

Expanding the expression we again get

$$P(A \setminus B) = P(A) - P(A \cap B) = 0.4 - 0.2 = 0.2$$

d)

Expanding the expression we get

$$P(\bar{A} \setminus B) = P(\bar{A}) - P(\bar{A} \cap B) = (1 - 0.4) - 0.5 = 0.1$$

e)

By the inclusion-exclusion principle we have

$$P(\bar{A} \cup B) = P(\bar{A}) + P(B) - P(\bar{A} \cap B) = (1 - 0.4) + 0.7 - 0.5 = 0.8$$

f) By the distributive law

$$A \cap (B \cup \bar{A}) = (A \cap B) \cup (A \cap \bar{A})$$

As

$$A \cap \bar{A} = \emptyset$$

by definition. We get that

$$P(A \cap (B \cup \bar{A})) = P((A \cap B) \cup \emptyset) = 0.2$$

**Problem 15** a)

We assume the rolls are independent, as such

$$P(X_2 = 4) = \frac{1}{6}$$

b)

The sample space is given by

$$\begin{aligned} S = \{ & (1, 1), (1, 2), (1, 3), (1, 4), (1, 5), (1, 6), (2, 1), (2, 2), (2, 3), (2, 4), \\ & (2, 5), (2, 6), (3, 1), (3, 2), (3, 3), (3, 4), (3, 5), (3, 6), (4, 1), (4, 2), \\ & (4, 3), (4, 4), (4, 5), (4, 6), (5, 1), (5, 2), (5, 3), (5, 4), (5, 5), \\ & (5, 6), (6, 1), (6, 2), (6, 3), (6, 4), (6, 5), (6, 6) \} \end{aligned}$$

The outcomes that satisfy the event are

$$\{X_1 + X_2 = 7\} = \{(1, 6), (2, 5), (3, 4), (4, 3), (5, 2), (6, 1)\}$$

As such

$$P(X_1 + X_2 = 7) = \frac{6}{36} = \frac{1}{6}$$

c)

As they are independent we determine the probability as

$$P(X_1 \neq 2 \cap X_2 \geq 4) = P(X_1 \neq 2)P(X_2 \geq 4) = \frac{5}{6} \cdot \frac{3}{6} = \frac{15}{36} = \frac{5}{12}$$

**Problem 16** a)

As the sum of the individual probabilities must equate to 1 we wish to determine a  $c$  that satisfies the equation

$$\sum_{k=1}^{\infty} P(k) = \sum_{k=1}^{\infty} \frac{c}{3^k} = 1$$

We apply the formula for a geometric series

$$\sum_{k=0}^{\infty} cr^k = \frac{a}{1-r} \text{ for } |r| < 1$$

As this starts from 0 we subtract  $c$  as  $P(\{0\}) = \frac{c}{k^0} = c$

$$\begin{aligned} 1 &= -c + \sum_{k=0}^{\infty} c \left(\frac{1}{3}\right)^k \\ &= -c + \frac{c}{1 - \frac{1}{3}} \end{aligned}$$

By isolation for  $c$  we then get that

$$\frac{2}{3} = -\frac{2}{3}c + c \Leftrightarrow \frac{2}{3} = \frac{1}{3}c \Leftrightarrow 2 = c$$

b)

The set is given by the union of the 3, as such

$$\begin{aligned} P(\{2, 4, 6\}) &= P(\{2\} \cup \{4\} \cup \{6\}) \\ &= P(\{2\}) + P(\{4\}) + P(\{6\}) \\ &= \frac{2}{9} + \frac{2}{81} + \frac{2}{729} \\ &= \frac{182}{729} \end{aligned}$$

c) This must equate to the complement of  $P(\{1, 2\})$ , as such

$$\begin{aligned} P(\{3, 4, 5, \dots\}) &= 1 - P(\{1, 2\}) \\ &= 1 - \left(\frac{2}{3} + \frac{2}{9}\right) \\ &= 1 - \frac{8}{9} \\ &= \frac{1}{9} \end{aligned}$$

**Problem 17** We have that

$$\begin{aligned} P(A) &= P(B) \\ P(C) &= 2P(D) \\ P(A \cup C) &= P(A) + P(C) = 0.6 \end{aligned}$$

We rewrite all terms as functions of  $P(A)$ , these must equate to 1 as a team has to win, making it the sample space

$$\begin{aligned} 1 &= P(A) + P(A) + (0.6 - P(A)) + \frac{0.6 - P(A)}{2} \\ &= 0.5P(A) + 0.9 \end{aligned}$$

Isolating for  $P(A)$  we get

$$P(A) = \frac{1 - 0.9}{0.5} = 0.2$$

From this we can determine the rest of the probabilities using the requirements set earlier

$$P(A) = P(B) = 0.2$$

$$P(C) = 0.6 - P(A) = 0.4$$

$$P(D) = 0.5P(C) = 0.2$$

$$\sum P = 0.2 + 0.2 + 0.4 + 0.2 = 1$$

**Problem 18** a)

We insert  $t = 1$

$$P(T \leq 1) = \frac{1}{16}1^2 = \frac{1}{16}$$

b)

This must equate to the complement of it taking 2 hours, as such

$$P(2 > t) = 1 - \left(\frac{1}{16}2^2\right) = \frac{3}{4}$$

c)

This equates to the difference between the probability of the job taking more than 1 hour and more than 3 hours

$$P(1 \leq T \leq 3) = P(T \leq 1) - P(T \leq 3) = \frac{1}{16}3^2 - \frac{1}{16}1^2 = \frac{1}{2}$$

**Problem 19** The problem has the form of the quadratic equation

$$ax^2 + bx + c = 0 \implies x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

For this to have real solutions it is necessary that

$$b^2 - 4ac \geq 0 \vee a \neq 0$$

Translating this to the given equation we wish to determine values  $A, B \in [0, 1]$  that satisfy

$$1^2 - 4AB \geq 0 \vee A \neq 0$$

Seeing this as a function based on the given figure, we write that

$$1^2 - 4xy \geq 0$$

Isolating for  $y$  it becomes apparent that

$$\begin{aligned} 1^2 - 4xy &\geq 0 \\ 1^2 &\geq 4xy \\ \frac{1}{4x} &\geq y \end{aligned}$$

As the maximal  $y$ -value is 1 we determine the intersection

$$1 = \frac{1}{4x} \Leftrightarrow x = \frac{1}{4}$$

Now we can determine the probability as the proportion of the area that fits under this curve in the interval for which it is inside the unit square

$$\begin{aligned} P(1 - 4AB \geq 0) &= \frac{1}{4} + \frac{1}{4} \int_{\frac{1}{4}}^1 \frac{1}{4x} \\ &= \frac{1}{4} + \frac{1}{4} [\ln x]_{\frac{1}{4}}^1 \\ &= \frac{1}{4} + \frac{1}{4} \left( \ln 1 - \ln \frac{1}{4} \right) \\ &\approx \frac{3}{5} \end{aligned}$$

**Problem 20** a)

As every set is a subset of the next, it is clear that  $A_n$  will be the largest or equal to the largest set in the sequence, as such

$$P\left(\bigcup_{i=1}^{\infty} A_i\right) = \lim_{n \rightarrow \infty} P(A_n)$$

As all the sets are subsets of  $A_n$  and therefore their union will create  $A_n$ .

b)

Here it is the opposite where  $A_n$  will be the smallest or equal to the smallest set in the sequence as such

$$P\left(\bigcap_{i=1}^{\infty} A_i\right) = \lim_{n \rightarrow \infty} P(A_n)$$

As the intersection will always be limited by the smallest set in the sequence.



**Problem 21 DO LATER**

**Problem 22** Let  $A$  be the event that the customer purchased a cup of coffee and  $B$  the event they purchased a piece of cake, then

$$P(A) = 0.7$$

$$P(B) = 0.4$$

$$P(A \cap B) = 0.2$$

By conditional probability we have that

$$\begin{aligned} P(A|B) &= \frac{P(A \cap B)}{P(B)} \\ &= \frac{0.2}{0.4} \\ &= 0.5 \end{aligned}$$

**Problem 23 a)**

$$\begin{aligned} P(A|B) &= \frac{P(A \cap B)}{P(B)} \\ &= \frac{0.2}{0.35} \\ &\approx 0.57 \end{aligned}$$

b)

$$\begin{aligned} P(C|B) &= \frac{P(B \cap C)}{P(B)} \\ &= \frac{0.15}{0.35} \\ &\approx 0.43 \end{aligned}$$

c)

$$\begin{aligned} P(B|A \cup C) &= \frac{P(B \cap (A \cup C))}{P(A \cup C)} \\ &= \frac{0.25}{0.7} \\ &\approx 0.36 \end{aligned}$$

d)

$$\begin{aligned}
 P(B|A \cap C) &= \frac{P(B \cap (A \cap C))}{P(A \cap C)} \\
 &= \frac{0.1}{0.2} \\
 &= 0.5
 \end{aligned}$$

**Problem 24** As this is an interval with infinite points, we determine the proportion of the interval that is spanned by the interval were observing, as such

$$P(A \leq X \leq B) = \frac{B - A}{10}$$

a)

$$P(2 \leq X \leq 5) = \frac{5 - 2}{10} = 0.3$$

b)

$$P(X \leq 2 | X \leq 5) = \frac{2}{5} = 0.4$$

c)

$$\begin{aligned}
 P(3 \leq X \leq 8 | X \geq 4) &= \frac{P(3 \leq X \leq 8 \cap X \geq 4)}{P(X \geq 4)} \\
 &= \frac{P(4 \leq X \leq 8)}{0.6} \\
 &= \frac{\frac{8-4}{10}}{0.6} \\
 &= \frac{2}{3}
 \end{aligned}$$

**Problem 25** We define the event  $A$  as getting an A in the course and  $B$  as living on campus

$$\begin{aligned}
 P(A) &= \frac{120}{600} = 0.2 \\
 P(B) &= \frac{200}{600} = \frac{1}{3} \\
 P(A|\bar{B}) &= \frac{80}{400} = 0.2P(A|B) = \frac{40}{200} = 0.2
 \end{aligned}$$

For  $A$  and  $B$  to be independent we would expect

$$P(A|B) = P(A)$$

Which is true, as such, the 2 events are independent.

**Problem 26** Define  $N_1, N_6$  as the number of times out of  $n$  that a 1 or 6 is rolled, and let  $X_i$  be the  $i$ 'th roll, then

$$P(N_1 \geq 1 \cap N_6 \geq 1) = 1 - P(\overline{N_1 \geq 1 \cap N_6 \geq 1})$$

By De Morgans Law we get

$$\begin{aligned} &= 1 - P(N_1 = 0 \cap N_6 = 0) \\ &= 1 - (P(X_1 \neq 1, X_2 \neq 1, \dots, X_n \neq 1) + P(X_1 \neq 6, X_2 \neq 6, \dots, X_n \neq 6) \\ &\quad - P((X_1 \neq 1, X_2 \neq 1, \dots, X_n \neq 1) \cap (X_1 \neq 6, X_2 \neq 6, \dots, X_n \neq 6))) \end{aligned}$$

As the final term is equivalent to

$$\bigcap_{i=1}^n P(X_i \neq 1 \cap X_i \neq 6)$$

We get that

$$\begin{aligned} &= 1 - \left( 2 \left( \frac{5}{6} \right)^n - \left( \frac{4}{6} \right)^n \right) \\ &= 1 - \frac{2 \cdot 5^n - 4^n}{6^n} \end{aligned}$$

**Problem 27** a)

Tree is finished on paper.

b)

We sum the probabilities that result in an error

$$\begin{aligned} P(E) &= P(E|G) + P(E\overline{G}) \\ &= 0.08 + 0.06 \\ &= 0.14 \end{aligned}$$

c)

From the definition of conditional probability we have

$$\begin{aligned} P(G|\overline{E}) &= \frac{P(G \cap \overline{E})}{P(\overline{E})} \\ &= \frac{0.72}{1 - 0.14} \\ &= \frac{0.72}{0.86} \\ &\approx 0.84 \end{aligned}$$

**Problem 28** We define  $D$  as the unit being defective, the probability of picking a defective unit on the first draw is

$$P(D_1) = \frac{5}{100}$$

Assuming that one wasn't defective the next draw has

$$P(D_2|\overline{D_1}) = \frac{5}{99}$$

And the 3rd

$$P(D_3|\overline{D_2}, \overline{D_1}) = \frac{5}{98}$$

We wish to determine the probability of the event

$$P(D = 1) = P(\{(D, \overline{D}, \overline{D}), (\overline{D}, D, \overline{D}), (\overline{D}, \overline{D}, D)\})$$

As these are disjoint we determine them as the sum of their individual probabilities

$$\begin{aligned} P(D = 1) &= P(\{(D, \overline{D}, \overline{D})\}) + P(\{(\overline{D}, D, \overline{D})\}) + P(\{(\overline{D}, \overline{D}, D)\}) \\ &= \frac{5}{100} \cdot \frac{95}{99} \cdot \frac{94}{98} + \frac{95}{100} \cdot \frac{5}{99} \cdot \frac{94}{98} + \frac{95}{100} \cdot \frac{94}{99} \cdot \frac{5}{98} \\ &\approx 0.14 \end{aligned}$$

**Problem 29** a)

As the components are connected in series, the probability of the system being functional is equal to the complement of one or more of the components not working. Let  $F$  be the event that the system is functional and  $P(C_i)$  be the event that the  $i$ 'th component is functional.

$$P(F) = P(C_1 \cap C_2 \cap C_3) = P(C_1)P(C_2)P(C_3)$$

b)

As they're connected in parallel they must all fail for the circuit to be non-functional, as such the probability must be given by the union of them

$$P(F) = P(C_1 \cup C_2 \cup C_3)$$

By the inclusion-exclusion principle

$$\begin{aligned} P(F) &= P(C_1) + P(C_2) + P(C_3) - P(C_1)P(C_2) - P(C_1)P(C_3) - \\ &\quad P(C_2)P(C_3) + P(C_1)P(C_2)P(C_3) \end{aligned}$$

c)

The probability that this circuit works must be given by the probability that either  $C_1$  and  $C_3$  or  $C_2$  and  $C_3$  are functional.

$$\begin{aligned} P(F) &= P((C_1 \cap C_3) \cup (C_2 \cap C_3)) \\ &= P(C_1 \cap C_3) + P(C_2 \cap C_3) - P((C_1 \cap C_3) \cap (C_2 \cap C_3)) \\ &= P(C_1)P(C_3) + P(C_2)P(C_3) - P(C_1 \cap C_2 \cap C_3) \end{aligned}$$

d)

The probability that this circuit works is given by the probability that either  $C_3$  or both  $C_1$  and  $C_2$  work

$$\begin{aligned} P(F) &= P(C_3 \cup (C_1 \cap C_2)) \\ &= P(C_3) + P(C_1 \cap C_2) \\ &= P(C_3) + P(C_1)P(C_2) - P(C_1)P(C_2)P(C_3) \end{aligned}$$

e)

The probability that this circuit works is determined by whether  $C_1, C_2$  and  $C_5$  or  $C_3, C_4$  and  $C_5$  work, therefore


$$\begin{aligned} P(F) &= P(C_1 \cap C_2 \cap C_5) + P(C_3 \cap C_4 \cap C_5) - P(C_1 \cap C_2 \cap C_3 \cap C_4 \cap C_5) \\ &= P(C_1)P(C_2)P(C_5) + P(C_3)P(C_4)P(C_5) - \prod_{i=1}^5 P(C_i) \end{aligned}$$

**Problem 30** a)

Rewriting the set as a piecewise function we get

$$A = \begin{cases} x - y \leq \frac{1}{2} & x \geq y \\ y - x \leq \frac{1}{2} & y \geq x \end{cases} \implies \begin{cases} y \leq x - \frac{1}{2} & x \geq y \\ y \leq \frac{1}{2} + x & y \geq x \end{cases}$$

At the same time  $B$  is equal to the area above the  $45^\circ$  line in the unit square. Drawing this in Geogebra gives us



b)

Using geometry to calculate the areas we get that

$$P(A) = 1 - 2 \cdot 0.5 \cdot 0.5 \cdot 0.5 = \frac{3}{4}$$

$$P(B) = 1 - 0.5 \cdot 1 \cdot 1 = \frac{1}{2}$$

c)

We determine the probability of the intersection using geometry

$$P(A \cap B) = 1 - 0.5 - 0.5 \cdot 0.5 \cdot 0.5 = \frac{3}{8}$$

For them to be independent

$$P(A \cap B) = P(A)P(B) = \frac{3}{4} \cdot \frac{1}{2} = \frac{3}{8}$$

As such they are independent.

**Problem 31** Let  $A$  be the event that an email is spam,  $B$  the event that it contains the word refinance, we then have that

$$P(A) = 0.5$$

$$P(B|A) = 0.01$$

$$P(B|\overline{A}) = 0.00001$$

We wish to determine the probability

$$P(A|B)$$

By Bayes rule we have that

$$\begin{aligned} P(A|B) &= \frac{P(B|A)P(A)}{P(B)} \\ &= \frac{0.01 \cdot 0.5}{0.01 \cdot 0.5 + 0.00001 \cdot 0.5} \\ &\approx 0.99 \end{aligned}$$

**Problem 32** a)

For the path to be open it is clear that  $B_1, B_4$  or  $B_2, B_5$  or  $B_1, B_3, B_5$  or  $B_2, B_3, B_4$  must be open

$$P(A) = P(\{B_1 \cap B_3 \cap B_4\} \cup \{B_1 \cap B_3 \cap B_5\} \cup \{B_2 \cap B_3 \cap B_5\} \cup \{B_2 \cap B_3 \cap B_4\})$$

We rename each path to  $P_i$  where  $i$  is the arbitrary number assigned to the

path. By the inclusion exclusion principle we have that

$$\begin{aligned}
P(A) &= P(P_1) + P(P_2)P(P_3) + P(P_4) - P(P_1)P(P_2) - P(P_1)P(P_3) \\
&\quad - P(P_1)P(P_4) - P(P_2)P(P_3) - P(P_2)P(P_4) - P(P_3)P(P_4) \\
&\quad + P(P_1)P(P_2)P(P_3) + P(P_1)P(P_2)P(P_4) + P(P_1)P(P_3)P(P_4) \\
&\quad + P(P_2)P(P_3)P(P_4) - P(P_1)P(P_2)P(P_3)P(P_4) \\
&= P(B_1 \cap B_4) + P(B_2 \cap B_5) + P(B_1 \cap B_3 \cap B_5) + P(B_2 \cap B_3 \cap B_4) \\
&\quad - P((B_1 \cap B_4) \cap (B_2 \cap B_5)) - P((B_1 \cap B_4) \cap (B_1 \cap B_3 \cap B_5)) \\
&\quad - P((B_1 \cap B_4) \cap (B_2 \cap B_3 \cap B_4)) - P((B_2 \cap B_5) \cap (B_1 \cap B_3 \cap B_5)) \\
&\quad - P((B_2 \cap B_5) \cap (B_2 \cap B_3 \cap B_4)) - P((B_1 \cap B_3 \cap B_5) \cap (B_2 \cap B_3 \cap B_4)) \\
&\quad + P((B_1 \cap B_4) \cap (B_2 \cap B_5) \cap (B_2 \cap B_3 \cap B_5)) \\
&\quad + P((B_1 \cap B_4) \cap (B_2 \cap B_5) \cap (B_2 \cap B_3 \cap B_4)) \\
&\quad + P((B_1 \cap B_4) \cap (B_1 \cap B_3 \cap B_5) \cap (B_2 \cap B_3 \cap B_4)) \\
&\quad + P((B_2 \cap B_5) \cap (B_2 \cap B_3 \cap B_4) \cap (B_2 \cap B_3 \cap B_4)) \\
&\quad - P((B_1 \cap B_4) \cap (B_2 \cap B_5) \cap (B_1 \cap B_3 \cap B_5) \cap (B_2 \cap B_3 \cap B_4)) \\
&= P(B_1 \cap B_4) + P(B_2 \cap B_5) + P(B_1 \cap B_3 \cap B_5) + P(B_2 \cap B_3 \cap B_4) \\
&\quad - P(B_1 \cap B_2 \cap B_4 \cap B_5) - P(B_1 \cap B_3 \cap B_4 \cap B_5) \\
&\quad - P(B_1 \cap B_2 \cap B_3 \cap B_4) - P(B_1 \cap B_2 \cap B_3 \cap B_5) \\
&\quad - P(B_2 \cap B_3 \cap B_4 \cap B_5) - P(B_1 \cap B_2 \cap B_3 \cap B_4 \cap B_5) \\
&\quad + P(B_1 \cap B_2 \cap B_3 \cap B_4 \cap B_5) + P(B_1 \cap B_2 \cap B_3 \cap B_4 \cap B_5) \\
&\quad + P(B_1 \cap B_2 \cap B_3 \cap B_4 \cap B_5) + P(B_1 \cap B_2 \cap B_3 \cap B_4 \cap B_5) \\
&\quad - P(B_1 \cap B_2 \cap B_3 \cap B_4 \cap B_5) \\
&= P_1P_4 + P_2P_5 + P_1P_3P_5 + P_2P_3P_4 - P_1P_2P_4P_5 - P_1P_3P_4P_5 - P_1P_2P_3P_4 \\
&\quad - P_1P_2P_3P_5 - P_2P_3P_4P_5 - P_1P_2P_3P_4P_5 + 4P_1P_2P_3P_4P_5 - P_1P_2P_3P_4P_5 \\
&= P_1P_4 + P_2P_5 + P_1P_3P_5 + P_2P_3P_4 - P_1P_2P_4P_5 - P_1P_3P_4P_5 - P_1P_2P_3P_4 \\
&\quad - P_1P_2P_3P_5 - P_2P_3P_4P_5 + 2P_1P_2P_3P_4P_5 \\
&= P_1P_4(1 - P_2P_5 - P_3P_5 - P_2P_3 + 2P_2P_3P_5) + P_2P_5 \\
&\quad + P_1P_3P_5 + P_2P_3(P_4 - P_1P_5 - P_4P_5)
\end{aligned}$$

b) We use Bayes rule to determine the probability, conditioning  $A$  on  $B_3$  we effectively cancel out the  $P_3$  terms as  $P(P_3|B_3) = 1$

$$\begin{aligned}
P(A|B_3) &= P_1P_4(1 - P_2P_5 - P_5 - P_2 + 2P_2P_5) + P_2P_5 + P_1P_5 \\
&\quad + P_2(P_4 - P_1P_5 - P_4P_5)
\end{aligned}$$



By Bayes rule we have that

$$P(B_3|A) = \frac{P(A|B_3)P_3}{P(A)}$$

Inserting known information

$$P(B_3|A) = \frac{P_1P_3P_4(1 - P_2P_5 - P_5 - P_2 + 2P_2P_5) + P_2P_3P_5 + P_1P_3P_5 + P_2P_3(P_4 - P_1P_5 - P_4P_5)}{P_1P_4(1 - P_2P_5 - P_3P_5 - P_2P_3 + 2P_2P_3P_5) + P_2P_5 + P_1P_3P_5 + P_2P_3(P_4 - P_1P_5 - P_4P_5)}$$

**Problem 33** Before any extra information is given we have that

$$P(H_1) = P(H_2) = P(H_3) = \frac{1}{3}$$

We let  $C_i$  be the event that the host opens the  $i$ 'th door and  $H_i$  be the event that the car is behind the  $i$ 'th door, assuming we opened the 1st door we get the following probabilities

$$P(C_1|H_1) = 0$$

$$P(C_2|H_1) = \frac{1}{2}$$

$$P(C_3|H_1) = \frac{1}{2}$$

$$P(C_1|H_2) = 0$$

$$P(C_2|H_2) = 0$$

$$P(C_3|H_2) = 1$$

$$P(C_1|H_3) = 0$$

$$P(C_2|H_3) = 1$$

$$P(C_3|H_3) = 0$$

If the host choses to open the 3rd door, we wish to determine whether

$$P(H_2|C_3) > P(H_1)$$

As that would put us in an advantageous situation. We apply Bayes rule and get

$$\begin{aligned} P(H_2|C_3) &= \frac{P(C_3|H_2)P(H_2)}{P(C_3|H_1)P(H_1) + P(C_3|H_2)P(H_2) + P(C_3|H_3)P(H_3)} \\ &= \frac{1 \cdot \frac{1}{3}}{\frac{1}{2} \cdot \frac{1}{3} + 1 \cdot \frac{1}{3} + 0 \cdot \frac{1}{3}} \\ &= \frac{2}{3} \end{aligned}$$

As

$$P(H_2|C_3) > P(H_1)$$

It would be advantageous to switch our guess.

**Problem 34** a)

We have that

$$\begin{aligned} P(A) &= \frac{1}{6} \\ P(B) &= \frac{1}{6} \\ P(C) &= \frac{1}{6} \end{aligned}$$

For  $A$  and  $B$  to be independent it must be true that

$$P(A \cap B) = P(A)P(B)$$

The intersection of the 2 is only satisfied by the pair  $(2, 5)$ , as such

$$P(A \cap B) = \frac{1}{36} \stackrel{?}{=} \frac{1}{6} \cdot \frac{1}{6}$$

As the sides are equivalent the two are independent.

b)

Again we determine  $P(A \cap C)$ , this is again a single pair,  $(2, 3)$  as such the rest of the problem is equivalent to the previous. The two are independent.

c)

Again  $P(B \cap C)$  is a single pair  $(4, 3)$ , as such they are once again independent.

d)

As we already know we satisfy 3 of the requirements we must at last determine whether

$$P(A \cap B \cap C) = P(A)P(B)P(C)$$

This however is not true as the intersection is empty due to 2 and 3 never adding up to 7, as such

$$P(A \cap B \cap C) = P(\emptyset) = 0$$

Which is not equal to the product of the individual probabilities as

$$0 \neq \left(\frac{1}{6}\right)^3$$

**Problem 35**

**Problem 36** We define  $H \dots H$  as the event that  $n$  coin tosses resulted in heads, and  $H_{n+1}$  as the event that the next coin toss results in heads and  $F$  be that we picked the fair coin. We wish to determine the probability

$$P(H_{n+1}|H \dots H)$$

From the law of total probability it becomes clear that

$$P(H_{n+1}) = P(H_{n+1}|F)P(F) + P(H_{n+1}|\bar{F})P(\bar{F})$$

Conditioning both sides on  $H \dots H$  we then get

$$P(H_{n+1}|H \dots H) = P(H_{n+1}|F, H \dots H)P(F|H \dots H) + P(H_{n+1}|\bar{F}, H \dots H)P(\bar{F}|H \dots H)$$

The terms with two conditions are conditionally independent on  $H \dots H$  as only the selected coin matters, as such

$$\begin{aligned} P(H_{n+1}|F, H \dots H) &= \frac{1}{2} \\ P(H_{n+1}|\bar{F}, H \dots H) &= 1 \end{aligned}$$

By using Bayes rule we can determine  $P(F|H \dots H)$  as

$$\begin{aligned} P(F|H \dots H) &= \frac{P(H \dots H|F)P(F)}{P(H \dots H|F)P(F) + P(H \dots H|\bar{F})P(\bar{F})} \\ &= \frac{\frac{1}{2}^n \cdot \frac{1}{2}}{\frac{1}{2}^n \cdot \frac{1}{2} + 1 \cdot \frac{1}{2}} \\ &= \frac{\frac{1}{2}^n}{\frac{1}{2}^n + 1} \\ &= \frac{1}{1 + \frac{1}{\frac{1}{2}^n}} \\ &= \frac{1}{2^n + 1} \end{aligned}$$

And  $P(\bar{F}|H \dots H)$  as its complement

$$P(\bar{F}|H \dots H) = 1 - \frac{1}{2^n + 1}$$

The probability of the  $n$ 'th toss being must therefore be the sum of these probabilities

$$\begin{aligned}P(H_{n+1}|H \dots H) &= P(F|H \dots H)P(F) + P(\overline{F}|H \dots H) \\&= \frac{1}{2} \cdot \frac{1}{2^n + 1} + \left(1 - \frac{1}{2^n + 1}\right) \\&= \frac{1}{2(2^n + 1)} + \left(1 - \frac{1}{2^n + 1}\right) \\&= 1 + \frac{-1}{2(2^n + 1)} \\&= 1 - \frac{1}{2(2^n + 1)}\end{aligned}$$

**Problem 37**

**Problem 38**

**Problem 39****A.2 Chapter 2****Problem 1**

**Problem 2** We make use of combinations with  $n = 12, k = 8$

$$\binom{12}{8} = \frac{12!}{8!(12-8)!} = 495$$

**Problem 3** a)

The probability of picking exactly 4 black phones will be equal to

$$P(B = 4) = \frac{|B = 4|}{|S|}$$

As they are not put back we determine  $|S|$  as the amount of 10-element subsets of a 50 element set using combinations

$$|S| = \binom{50}{10} = \frac{50!}{10!(50-10)!} = 10272278170$$

Simultaneously we can determine the amount of sets containing 4 black phones and 6 white phones as

$$|B = 4| = \binom{20}{4} \binom{30}{10-4} = \frac{20!}{4!(20-4)!} \cdot \frac{30!}{6!(30-6)!} = 2876839875$$

As such

$$P(B = 4) = \frac{2876839875}{10272278170} \approx 0.28$$

b) We wish to determine  $P(B \in \{0, 1, 2\})$ , we use the same method as the previous section and get that

$$\begin{aligned} |B \in \{0, 1, 2\}| &= \sum_{i=0}^2 \binom{20}{i} \binom{30}{10-i} \\ &= \binom{20}{0} \binom{30}{10} + \binom{20}{1} \binom{30}{9} + \binom{20}{2} \binom{30}{8} \\ &= 30045015 + 286143000 + 1112055750 \\ &= 1428243765 \end{aligned}$$

And again we determine the probability as

$$P(B \in \{0, 1, 2\}) = \frac{1428243765}{10272278170} \approx 0.14$$

**Problem 4** a)

We once again make use of combinations, the sample space is given by

$$|S| = \binom{52}{5} = \frac{52!}{5!(52-5)!} = 2598960$$

The amount of 5 element sets consisting of 1 ace must be given by

$$|A = 1| = \binom{4}{1} \binom{48}{4} = \frac{4!}{1!(4-1)!} \cdot \frac{48!}{4!(48-4)!} = 778320$$

Determining the probability is then as simple as

$$P(A = 1) = \frac{778320}{2598960} \approx 0.30$$

b) The phrasing of the question suggests that its easier to determine the complement, as such we find the probability of getting no aces using the same method as in the previous section

$$|A = 0| = \binom{4}{0} \binom{48}{5} = \frac{4!}{0!(4)!} \cdot \frac{48!}{5!(48-5)!} = 1712304$$

The probability is then

$$P(A = 0) = \frac{1712304}{2598960} \approx 0.66$$

And we then determine the complement as this is equal to the probability of getting 1 or more aces

$$P(A \in \{1, 2, 3, 4\}) = P(\overline{A = 0}) = 1 - P(A = 0) \approx 0.34$$

**Problem 5** We wish to determine the probability

$$P(A = 2|A \geq 1)$$

As the reverse conditional probability is independent of the condition we know that

$$P(A \geq 1|A = 2) = 1$$

Using Bayes law we then find that

$$P(A = 2|A \geq 1) = \frac{P(A \geq 1|A = 2)P(A = 2)}{P(A \geq 1)}$$

We determine  $|A = 2|$  using the same method as in previous section

$$|A = 2| = \binom{4}{2} \binom{48}{3} = \frac{4!}{2!(4-2)!} \cdot \frac{48!}{3!(48-3)!} = 103776$$

And then make use of this to determine the probability of getting 2 aces

$$P(A = 2) = \frac{103776}{2598960} \approx 0.04$$

As such we have that

$$P(A = 2|A \geq 1) = \frac{1 \cdot 0.04}{0.34} \approx 0.12$$

**Problem 6** Assuming the cards are dealt in bulk we can determine that

$$n_{\text{cards}} = 52 - 26 = 26$$

$$n_{\text{spades}} = 13 - 7 = 6$$

As such 26 cards containing 6 spades will be dealt to player  $C$ , giving us that the sample space is

$$|S| = \binom{26}{13} = \frac{26!}{13!(26-13)!}$$

Whilst the amount of decks containing exactly 4 spades is

$$|C = 4| = \binom{6}{4} \binom{20}{13-4} = \frac{6!}{4!(6-4)!} \cdot \frac{20!}{9!(20-9)!}$$

As such the probability is

$$P(C = 4) = \frac{\binom{6}{4} \binom{20}{9}}{\binom{26}{13}} \approx 0.24$$

**Problem 7** The sample space is given by

$$|S| = \binom{50}{15} = \frac{50!}{15!(50-15)!}$$

By inclusion exclusion we have that

$$P(Y \cup J) = P(Y) + P(J) - P(Y \cap J)$$

The amount of sets containing either you or Joe is

$$|Y| = |J| = \binom{1}{1} \binom{49}{14} = \frac{1!}{1!(1-1)!} \cdot \frac{49!}{14!(49-14)!}$$

The amount of sets containing both you and your friend Joe is

$$|Y, J| = \binom{2}{2} \binom{48}{13} = \frac{2!}{2!(2-2)!} \cdot \frac{48!}{13!(48-13)!}$$

As such the probability of both you and Joe being in group is

$$P(Y \cup J) = \frac{\binom{2}{2} \binom{48}{13}}{\binom{50}{15}} \approx 0.09$$

Inserting in the inclusion exclusion principle we then get

$$P(Y \cup J) = 2 \frac{\binom{1}{1} \binom{49}{14}}{\binom{50}{15}} - \frac{\binom{2}{2} \binom{48}{13}}{\binom{50}{15}} \approx 0.51$$

**Problem 8** For a set with  $n$  elements of which  $r$  are unique we have that there are

$$N = \frac{n!}{n_1! n_2! \dots n_r!}$$

Where  $n_i$  is the amount of repetitions of the given element, in Massachusetts we observe that

$$n_m = 1$$

$$n_a = 2$$

$$n_s = 4$$

$$n_c = 1$$

$$n_h = 1$$

$$n_u = 1$$

$$n_e = 1$$

$$n_t = 2$$

As such

$$N = \frac{13!}{2!4!2!} = 64864800$$



**Problem 9** a)

We use the binomial theorem and as such have that

$$P(k) = \binom{n}{k} p^k (1-p)^{n-k}$$

The probability of it containing 8 heads and 12 tails must simply be given by inserting  $k = 8$  and determining the amount of possible sets using combinations

$$P(8) = \binom{20}{8} p^8 (1-p)^{20-8}$$

b)

Observing more than 8 heads and more than 8 tails is only possible when observing either 9, 10 or 11 heads, as such the probability must be given by

$$P(H \geq 8, T \geq 8) = \sum_{k=9}^{11} \binom{20}{k} p^k (1-p)^{20-k}$$

**Problem 10** As every path will be described by some scrambled sequence of 20 movements to the right and 10 movements up, as such the problem boils down to finding out how many ways you can move right 20 times and up 10 times, which is given by

$$\frac{30!}{20!10!} = 30045015$$

**Problem 11** We have the sample size from the previous problem, we determine the amount of paths that end up at  $(10, 5)$  by determining the amount of paths that end at that point, this is given by

$$\frac{15!}{10!5!} = 3003$$

Afterwards there is once again 10 rights and 5 ups to reach the final point, giving us the same amount of paths once again, as such

$$P((10, 5) \in \text{Path}) = \frac{\left(\frac{15!}{10!5!}\right)^2}{\frac{30!}{20!10!}} \approx 0.30$$

**Problem 12** Using the binomial theorem we have that

$$P(k) = \binom{n}{k} p_a^k (1 - p_a)^{n-k}$$

As  $p_a$  denotes the probability of moving up for it to reach the wanted point we must have that  $k = 5$  and  $n = 15$ , therefore

$$P(5) = \binom{15}{5} p_a^5 (1 - p_a)^{15-5}$$

**Problem 13** a)

We make use of the law of total probability as well as the binomial theorem to write that

$$\begin{aligned} P(H \geq 3) &= \sum_{i=1}^2 P(H = 3|C_i)P(C_i) + P(H = 4|C_i)P(C_i) + P(H = 5|C_i)P(C_i) \\ &= 0.5 \sum_{i=1}^2 \binom{5}{3} p_i^3 (1 - p_i)^2 + \binom{5}{4} p_i^4 (1 - p_i)^1 + \binom{5}{5} p_i^5 \\ &\approx 0.35 \end{aligned}$$

b)

We wish to determine the probability

$$P(C_2|H \geq 3)$$

We can easily determine the reverse conditional probability using the binomial theorem

$$\begin{aligned} P(H \geq 3|C_2) &= \binom{5}{3} p^3 (1 - p)^2 + \binom{5}{4} p^4 (1 - p)^1 + \binom{5}{5} p^5 \\ &\approx 0.21 \end{aligned}$$

As we know  $P(H \geq 3)$  from the previous question, we apply Bayes rule and get that

$$\begin{aligned} P(C_2|H \geq 3) &= \frac{P(H \geq 3|C_2)P(C_2)}{P(H \geq 3)} \\ &= \frac{0.21 \cdot 0.5}{0.35} \\ &= 0.3 \end{aligned}$$

**Problem 14**

**Problem 15**

**Problem 16**

**Problem 17**

**Problem 18**

**Problem 19** We apply that the number of possible solutions for an equation equating to  $k$  with  $n$  distinct elements is given by

$$\binom{k+n-1}{k}$$

As the  $x_i$ 's are limited to the natural numbers we rewrite such that we are working in the domain  $\{0, 1, 2, \dots\}$  by stating that

$$y_i = x_i - 1 \Leftrightarrow x_i = y_i + 1$$

Rewriting the equation we then get

$$y_1 + y_2 + y_3 + y_4 + y_5 = 95$$

As such the number of solutions is given by

$$\binom{95+5-1}{k} = \frac{99!}{95!} = 3764376$$

**Problem 20** We can determine that as the sum of solutions of the equation

$$x_2 + x_3 + x_4 = 100 - x_1$$

We rewrite  $x_1$  as  $i$  and thus this can be expressed as

$$x_2 + x_3 + x_4 = 100 - i$$

By sum we then have

$$\begin{aligned} n_{\text{solutions}} &= \sum_{i=0}^{10} \binom{100-i+3-1}{100-i} \\ &= \binom{102}{100} + \binom{101}{99} + \dots + \binom{92}{90} \\ &= 51271 \end{aligned}$$

**Problem 21****A.3 Chapter 3**