


Ants on Manifolds

A CS Theory Seminar


Oliver Chubet, 2/21/25



Do space filling curves really exist?




What do we mean by *curve*?



What do we mean by *curve*?


Let X be some (*compact*) space.



What do we mean by *curve*?

Let X be some (*compact*) space.

A curve is an embedding $f : [0,1] \rightarrow X$.




What do we mean by *curve*?




Let X be some (*compact*) space.

A curve is an embedding $f : [0,1] \rightarrow X$.

(*No crossings, continuous*)



Cantor and Cardinality



“In 1878, Georg Cantor demonstrated that any two finite-dimensional smooth manifolds, no matter what their dimensions, have the same cardinality...

... and Mathematics has never been the same since.”

- Hans Sagan

Netto's Theorem

Netto's Theorem

A bijection $\phi : [0,1] \rightarrow [0,1]^2$

cannot be continuous.

Netto's Theorem

A bijection $\phi : [0,1] \rightarrow [0,1]^2$

cannot be continuous.

(More generally, manifolds of different dimensions cannot be homeomorphic.)

Corollary:


Corollary:

Space filling curves don't exist.

The background of the image features a repeating pattern of a Hilbert curve, a continuous fractal curve that fills a square. The curve is rendered in a light gray color, with its segments having a slight 3D effect through shading. The pattern is centered and covers the entire area of the image.


The Hilbert Curve

***(ith-order)* Hilbert Curves**





***(ith-order)* Hilbert Curves**




***(ith-order)* Hilbert Curves**







***(ith-order)* Hilbert Curves**




(ith-order) Hilbert Curves




***(ith-order)* Hilbert Curves**




(ith-order) Hilbert Curves




$$i = 0$$




$$i = 1$$



$i = 2$




$i = 3$




$i = 4$



$i = 5$



$i = 5$



The Hilbert Curve

The Hilbert Curve

Let $h_i : [0,1] \rightarrow [0,1]^2$ denote the order i th-order Hilbert curve.

The Hilbert Curve

Let $h_i : [0,1] \rightarrow [0,1]^2$ denote the order i th-order Hilbert curve.

The Hilbert curve is

The Hilbert Curve

Let $h_i : [0,1] \rightarrow [0,1]^2$ denote the order i th-order Hilbert curve.

The Hilbert curve is

$$h(x) := \lim_{i \rightarrow \infty} h_i(x)$$

The Hilbert Curve

Let $h_i : [0,1] \rightarrow [0,1]^2$ denote the order i th-order Hilbert curve.

The Hilbert curve is

$$h(x) := \lim_{i \rightarrow \infty} h_i(x)$$

Q: is h surjective?

The Hilbert Curve

Let $h_i : [0,1] \rightarrow [0,1]^2$ denote the order i th-order Hilbert curve.

The Hilbert curve is

$$h(x) := \lim_{i \rightarrow \infty} h_i(x)$$

Q: is h surjective?

Q: is h injective?

The Hilbert Curve

Let $h_i : [0,1] \rightarrow [0,1]^2$ denote the order i th-order Hilbert curve.







The Hilbert curve is

$$h(x) := \lim_{i \rightarrow \infty} h_i(x)$$

Q: is h surjective?

Q: is h injective?

Q: is h continuous?



***(i*th-order) Cantor Set**

***(i*th-order) Cantor Set**

$[0,1]$

(ith-order) Cantor Set




$[0,1]$




C_1




(ith-order) Cantor Set




(ith-order) Cantor Set




(ith-order) Cantor Set




(ith-order) Cantor Set



(ith-order) Cantor Set



(ith-order) Cantor Set



Defining the Cantor Set

“The Set of Excluded Middle Thirds”

Defining the Cantor Set

“The Set of Excluded Middle Thirds”

The *Cantor set* is $\mathcal{C} := \bigcap C_i$

Defining the Cantor Set

“The Set of Excluded Middle Thirds”

The **Cantor set** is $\mathcal{C} := \bigcap C_i$

Equivalently, \mathcal{C} is the set of all x such that

Defining the Cantor Set

“The Set of Excluded Middle Thirds”

The **Cantor set** is $\mathcal{C} := \bigcap C_i$

Equivalently, \mathcal{C} is the set of all x such that

$$x = \frac{2t_1}{3} + \frac{2t_2}{3^2} + \frac{2t_3}{3^3} + \frac{2t_4}{3^4} \dots, \text{ where } \forall i, t_i \in \{0,1\}$$

Defining the Cantor Set

“The Set of Excluded Middle Thirds”

The **Cantor set** is $\mathcal{C} := \bigcap C_i$

Equivalently, \mathcal{C} is the set of all x such that

$$x = \frac{2t_1}{3} + \frac{2t_2}{3^2} + \frac{2t_3}{3^3} + \frac{2t_4}{3^4} \dots, \text{ where } \forall i, t_i \in \{0,1\}$$

Or in **ternary**:

Defining the Cantor Set

“The Set of Excluded Middle Thirds”

The **Cantor set** is $\mathcal{C} := \bigcap C_i$

Equivalently, \mathcal{C} is the set of all x such that

$$x = \frac{2t_1}{3} + \frac{2t_2}{3^2} + \frac{2t_3}{3^3} + \frac{2t_4}{3^4} \dots, \text{ where } \forall i, t_i \in \{0,1\}$$

Or in **ternary**:

$$x = 0.(2t_1)(2t_2)(2t_3)(2t_4)\dots$$

Cardinality of \mathcal{C} vs. $[0,1]$

Cardinality of \mathcal{C} vs. $[0,1]$

Define $\eta : \mathcal{C} \rightarrow [0,1]$ by the map

Cardinality of \mathcal{C} vs. $[0,1]$

Define $\eta : \mathcal{C} \rightarrow [0,1]$ by the map

$$0.(2t_1)(2t_2)(2t_3)(2t_4)\dots \mapsto 0.t_1t_2t_3t_4\dots$$

Cardinality of \mathcal{C} vs. $[0,1]$

Define $\eta : \mathcal{C} \rightarrow [0,1]$ by the map

$$0.(2t_1)(2t_2)(2t_3)(2t_4)\dots \mapsto 0.t_1t_2t_3t_4\dots$$

- η is injective so the cardinality of \mathcal{C} is at least that of $[0,1]$

Cardinality of \mathcal{C} vs. $[0,1]$

Define $\eta : \mathcal{C} \rightarrow [0,1]$ by the map

$$0.(2t_1)(2t_2)(2t_3)(2t_4)\dots \mapsto 0.t_1t_2t_3t_4\dots$$

- η is injective so the cardinality of \mathcal{C} is at least that of $[0,1]$
- $\mathcal{C} \subset [0,1]$ so the cardinality of $[0,1]$ is at least that of \mathcal{C}

From the Cantor Set to $[0,1]^2$

From the Cantor Set to $[0,1]^2$

Define $\varphi : \mathcal{C} \rightarrow [0,1]^2$ by the mapping

From the Cantor Set to $[0,1]^2$

Define $\varphi : \mathcal{C} \rightarrow [0,1]^2$ by the mapping

$$0.(2t_1)(2t_2)(2t_3)(2t_4)\dots(2t_i) \mapsto \begin{bmatrix} 0.t_1t_3t_5\dots \\ 0.t_2t_4t_6\dots \end{bmatrix}$$

From the Cantor Set to $[0,1]^2$

Define $\varphi : \mathcal{C} \rightarrow [0,1]^2$ by the mapping

$$0.(2t_1)(2t_2)(2t_3)(2t_4)\dots(2t_i) \mapsto \begin{bmatrix} 0.t_1t_3t_5\dots \\ 0.t_2t_4t_6\dots \end{bmatrix}$$

Q: is φ surjective?

From the Cantor Set to $[0,1]^2$

Define $\varphi : \mathcal{C} \rightarrow [0,1]^2$ by the mapping

$$0.(2t_1)(2t_2)(2t_3)(2t_4)\dots(2t_i) \mapsto \begin{bmatrix} 0.t_1t_3t_5\dots \\ 0.t_2t_4t_6\dots \end{bmatrix}$$

Q: is φ surjective?

Q: is φ injective?

From the Cantor Set to $[0,1]^2$

Define $\varphi : \mathcal{C} \rightarrow [0,1]^2$ by the mapping

$$0.(2t_1)(2t_2)(2t_3)(2t_4)\dots(2t_i) \mapsto \begin{bmatrix} 0.t_1t_3t_5\dots \\ 0.t_2t_4t_6\dots \end{bmatrix}$$

Q: is φ surjective?

Q: is φ injective?

Q: is φ continuous?




The Lebesgue Curve


***(ith-order)* Lebesgue Curve**

Lebsegue extends φ continuously to $[0,1]$ by linear interpolation
(for each i th-order approximation)


Eg. \mathcal{L}_2




$i = 1$




$i = 2$




$i = 3$




$i = 4$



$i = 5$



$i = 5$



Properties of the Lebesgue Curve

Properties of the Lebesgue Curve

- z is *surjective* onto $[0,1]^2$

Properties of the Lebesgue Curve

- z is ***surjective*** onto $[0,1]^2$
- z is ***not injective***

Properties of the Lebesgue Curve

- z is ***surjective*** onto $[0,1]^2$
- z is ***not injective***
- z is a ***continuous*** map

Properties of the Lebesgue Curve

- z is ***surjective*** onto $[0,1]^2$
- z is ***not injective***
- z is a ***continuous*** map
- z is ***almost everywhere differentiable***

How to Fill Space


(A paradigm)

1. Net

2. Order

3. Recurse

Example:



Example: Quadtrees

